

arXiv:2511.05372v1 [quant-ph] 7 Nov 2025

# Adaptive quantum phase estimation can be better than non-adaptive

Noah Linden* Ronald de Wolf†

November 10, 2025

**Abstract**

Quantum phase estimation is one of the most important tools in quantum algorithms. It can be made non-adaptive (meaning all applications of the unitary $U_\phi$ happen simultaneously) without using more applications of $U_\phi$, albeit at the expense of using many more qubits. It is also known that there is no advantage for adaptive algorithms in the case where the phase that needs to be estimated is arbitrary or is uniformly random. Here we give examples of a special case of phase estimation, with a promise on the values that the unknown phase can take, where adaptive methods *are* provably better than non-adaptive methods by a factor of nearly 2 in the number of uses of $U_\phi$. We also prove some upper bounds on the maximum advantage that adaptive algorithms for phase estimation can achieve over non-adaptive ones.

## 1 Introduction

### 1.1 Phase estimation

In phase estimation we have the ability to apply a unitary $U_\phi$ (possibly controlled by another qubit) and are given a copy of an eigenstate $|\Psi\rangle$ of $U_\phi$ with eigenvalue $e^{2\pi i\phi}$. The task is to estimate the phase $\phi \in [0,1)$. Kitaev [Kit95] showed how to estimate $\phi$ with additive error $\delta$ (modulo 1) using roughly $1/\delta$ applications of $U_\phi$. The standard circuit for this is as follows. We use a set of $k = \lceil \log(1/\delta) \rceil$ auxiliary qubits starting in state $|0\rangle$ which will end up containing a $k$-bit estimate of $\phi$. These $k$ qubits are first put into a uniform superposition over all $j = 0, \ldots, 2^k - 1$, which controls how many times $U_\phi$ is applied to $|\Psi\rangle$. Because $|j\rangle \otimes U_\phi^j |\Psi\rangle = e^{2\pi i j\phi}|j\rangle \otimes |\Psi\rangle$, we obtain the following $k$-qubit state (tensored with our one copy of $|\Psi\rangle$)

$$\frac{1}{\sqrt{2^k}} \sum_{j=0}^{2^k-1} e^{2\pi i j\phi}|j\rangle. \tag{1}$$

By writing the binary expansion of $j$ as a sum of powers of 2, this can be nicely factored as

$$\frac{1}{\sqrt{2}}(|0\rangle + e^{2\pi i 2^{k-1}\phi}|1\rangle) \otimes \frac{1}{\sqrt{2}}(|0\rangle + e^{2\pi i 2^{k-2}\phi}|1\rangle) \otimes \cdots \otimes \frac{1}{\sqrt{2}}(|0\rangle + e^{2\pi i\phi}|1\rangle). \tag{2}$$

*School of Mathematics, University of Bristol. Partially supported by the UK Engineering and Physical Sciences Research Council through grant EP/T001062/1. n.linden@bristol.ac.uk

†QuSoft, CWI and University of Amsterdam, the Netherlands. Partially supported by the Dutch Research Council (NWO) through Gravitation-grant Quantum Software Consortium, 024.003.037. rdewolf@cwi.nl