

# Quantum simulation of the Dicke model in a two-dimensional ion crystal

Bryce Bullock[1,*], Sean R. Muleady[2,3,*], Jennifer F. Lilieholm[1], Yicheng Zhang[4], Arghavan Safavi-Naini[5,6], Robert J. Lewis-Swan[4], John J. Bollinger[1], Ana Maria Rey[7,8] and Allison L. Carter[1]

[1]NIST, Boulder, Colorado, USA. [2]QuICS, NIST and University of Maryland, USA. [3]JQI, NIST and University of Maryland, USA. [4]University of Oklahoma, USA. [5]University of Amsterdam, Netherlands. [6]QuSoft, Netherlands. [7]JILA, NIST and University of Colorado, Boulder, USA. [8]Center for Theory of Quantum Matter, University of Colorado, Boulder, USA.

[*]These authors contributed equally. Correspondence: sean.muleady@gmail.com; arey@jilau1.colorado.edu.

arXiv:2602.06114v3 [quant-ph] 18 Oct 2026

**Abstract**

Quantum many-body systems driven far from equilibrium can exhibit phase transitions, chaos and entanglement, but probing these phenomena in large, nearly closed systems remains challenging [1, 2]. The Dicke model, which couples an ensemble of two-level systems collectively to a single bosonic mode, provides a fundamental framework for exploring these phenomena in light–matter interactions [3, 4]. Here we show that a two-dimensional crystal of approximately 100 trapped ions realizes this model and accesses integrable dynamical phase transitions, signatures of chaotic dynamics and fluctuation-driven spin–phonon excitation growth. Adiabatic elimination of the vibrational mode reveals ferromagnetic and paramagnetic dynamical spin phases, whereas strong spin–phonon coupling produces irregular trajectories and local relaxation consistent with quantum thermalization. Following a quench from an unstable fixed point, correlated excitation growth precedes generalized vacuum Rabi collapses and revivals; calibrated calculations identify two-mode squeezing 2.6 dB below the standard quantum limit, supported by reconstruction from spin measurements. By connecting collective dynamical instabilities to experimentally accessible signatures of quantum correlations, these results establish large ion crystals as a scalable platform for investigating non-equilibrium light–matter dynamics and generating resources for quantum-enhanced sensing [5, 6].

## 1 Introduction

Understanding how classical dynamical instabilities and chaos persist or break down as quantum correlations develop is a central question in non-equilibrium many-body physics. Quantum simulators offer a way to address this question through unitary dynamics, which can produce dynamical phase transitions, entanglement growth and thermalization without an external bath. However, accessing these phenomena experimentally remains challenging because the exponential growth of Hilbert space makes coherent control increasingly difficult as system size grows.

Spin-boson models provide a natural setting for this exploration. Relevant throughout condensed-matter physics, quantum optics, and molecular and chemical dynamics, they couple discrete quantum spins to bosonic modes that can become highly occupied and mediate relaxation, transport, and collective behavior. A paradigmatic example is the Dicke model [3, 4], in which a collective spin couples to a single bosonic mode, supporting a crossover between integrable and chaotic dynamics [2, 7–11]. Despite the model's broad relevance, experiments have yet to access its quantum dynamics at scale [11–24], with features such as pair production, two-mode spin–boson squeezing, and generalized vacuum Rabi collapses and revivals remaining elusive.

Here we realize the Dicke model in a two-dimensional crystal of approximately 100 trapped $^9Be^+$ ions, whose internal states form a collective spin coupled to the center-of-mass vibrational mode (Fig. 1a). The long coherence time of this mode enables controlled, nearly unitary spin–phonon evolution across a broad range of dynamical regimes.

When the bosonic mode can be adiabatically eliminated, we observe a dynamical phase transition between ferromagnetic (trapped) and paramagnetic (untrapped) dynamics (Fig. 1b), distinguished by whether the average magnetization remains confined below the equator of the Bloch sphere [1, 25–32].