

arXiv:2605.16494v1 [cond-mat.quant-gas] 15 May 2026

# Universal dynamics from a single-particle dark state

Ruben Daraban,[1] Arghavan Safavi-Naini,[2,3] Johannes Schachenmayer,[1] and Mohammad Maghrebi[4,*]

[1] *CESQ/ISIS (UMR 7006), CNRS and Université de Strasbourg, 67000 Strasbourg, France*
[2] *Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*
[3] *QuSoft, Science Park 123, 1098 XG Amsterdam, The Netherlands*
[4] *Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan 48824 USA*

Open quantum systems can host dark or subradiant states whose decay is highly suppressed. While these states have been extensively studied in the few-excitation regime, their impact on the many-body dynamics remains largely unexplored. Here, we study a spin chain subject to correlated dissipation on neighboring sites, which admits a single-particle dark state at zero momentum. We show that the single-particle dark state qualitatively alters the many-body dynamics at long times, and identify its distinct universal behavior. While the zero-momentum mode is dark at the single-particle level, it decays slowly as $1/\log t$ as it becomes dressed by other modes through a dissipation-induced nonlinearity. We demonstrate that the momentum distribution takes a universal scaling form in $k\sqrt{t}$, and the total density decays as $1/\sqrt{t}\log t$. Our results further elucidate the origin of the conflicting results in recent works. Finally, we corroborate the analytics with matrix product state simulations and show that the same universal behavior persists for soft-core interactions, underscoring the universality of the emergent dynamics.

Many-body open quantum systems have attracted much attention in the last two decades. On one hand, they are very relevant to existing quantum simulators which emulate many-body physics of interest but also unavoidably come with noise and decoherence [1]. On the other hand, tailored dissipation can be a resource itself, for example, for quantum state preparation [2–4]. Furthermore, dissipation competes with Hamiltonian dynamics in driven open quantum systems, giving rise to genuinely novel forms of non-equilibrium physics [5].

Dissipative systems can host subradiant or *dark* states that are long-lived or do not decay. Subradiance arises when photons emitted by atoms interfere destructively suppressing their decay [6, 7]. Integrating out photons then leads to correlated long-range atomic dissipation, resulting in momentum-dependent dissipation which could be highly suppressed for subradiant states, a scenario that naturally occurs in waveguide QED [8]. While most studies of subradiance have focused on states with only a few atomic excitations [9–18], the many-body regime, with a few exceptions [19, 20], remains largely unexplored. Interestingly, states with few excitations are shown to exhibit a kind of *fermionization* [9, 19, 21–23], which also leaves a distinct fingerprint on the long-time dynamics [24]. In fact, even a single-particle dark state can drastically affect many-body dynamics since it results in a slow mode. A minimal such model, in which a spin chain is subject to correlated dissipation on neighboring sites, has recently attracted considerable attention [25–27]. This model admits a single-particle dark state, which leads to slow dynamics at long times. However, different works report conflicting results and even different critical exponents. At a technical level, while the Hamiltonian itself can be mapped to free fermions, dissipation renders the model highly nonlinear [26, 27].





FIG. 1. (a) Schematic representation of a spin chain (alternatively, hard- or soft-core bosons) subject to correlated dissipation on neighboring sites featuring a single-particle dark state at zero momentum ($\Gamma_{k=0} = 0$). (b) The momentum distribution at long times exhibits a diffusion-like behavior with the width decaying as $\sim 1/\sqrt{t}$, as if governed by the single-particle decay rate $\Gamma_k \sim k^2$. However, the $k = 0$ mode also decays, albeit slowly, as it becomes dressed with other modes due to a dissipation-induced nonlinearity. This decay takes a universal form of an inverse logarithm.