

# Stronger Welch Bounds and Optimal Approximate $k$-Designs

Riccardo Castellano,[1] Dmitry Grinko,[2,3,4] Sadra Boreiri,[1] Nicolas Brunner,[1] and Jef Pauwels[1,5]

[1] *Department of Applied Physics, University of Geneva, Switzerland*
[2] *QuSoft, Amsterdam, The Netherlands*
[3] *Institute for Logic, Language and Computation, University of Amsterdam, The Netherlands*
[4] *Korteweg-de Vries Institute for Mathematics, University of Amsterdam, The Netherlands*
[5] *Constructor University, 28759 Bremen, Germany*

The question of how well a finite ensemble of quantum states can approximate Haar randomness is relevant in many areas of quantum physics. Here we consider this problem from a geometrical perspective, via the notion of Welch bounds. The latter are a set of inequalities, originally derived in the context of signal processing, which precisely characterize exact $k$-designs, i.e. a finite ensemble of states that exactly reproduces the $k$-th moment of the Haar measure. We derive stronger Welch bounds which enable a quantitative and operationally meaningful characterisation of approximate $k$-designs. Our new bounds are often tight, allowing us to characterize optimal approximate designs in regimes where the cardinality of the ensemble is small. A notable example is complete sets of mutually unbiased bases (MUBs), which form an optimal approximate 3-design among 1-designs of their cardinality. This leads to a new variational approach to determine the existence of complete sets of MUBs, which we explore numerically in dimension six. We highlight a connection to shadow tomography, relating our bounds to the shadow norm. We comment on how the method extends to other types of designs, including toric and unitary designs. A key technical ingredient, is the computation of the complete spectrum of the partially transposed symmetric-subspace projector, including multiplicities and eigenvectors, which may find further applications.

arXiv:2602.13099v3 [quant-ph] 29 Sep 2026

## I. INTRODUCTION

How uniformly can a finite set of unit vectors be distributed in a complex vector space? This geometric question arises across mathematics, physics, and engineering. It is exemplified by the well-known Thomson problem, where one aims to place charged particles on the unit sphere as far apart from each other as possible, in order to minimize the total electrostatic potential energy. The question is also central in information theory, notably in coding theory, signal design, and compressed sensing. When signals are represented by complex vectors, there is a fundamental trade-off between the number of signals, the dimension of the vector space, and the cross-correlations between the signals. Formally, this is captured by a family of inequalities derived by Welch in 1974 [1], commonly referred to as Welch bounds, which have been extensively studied [2–5].

Welch bounds are also directly relevant to quantum information, where many problems require the construction of sets of states that are as uniformly distributed as possible. Consider a set of $N$ pure quantum states, represented by complex unit vectors in a Hilbert space of dimension $d$: $\chi := \{|\psi_i\rangle\}_{i=1}^N \subset \mathbb{C}^d$. The Welch inequality of order $k$ reads

$$\sum_{i,j=1}^{N} \left|\langle\psi_i|\psi_j\rangle\right|^{2k} \geq \frac{N^2}{\binom{d+k-1}{k}}, \qquad k \in \mathbb{N}^+.$$

The quantity on the left-hand side, called the unnormalized $k$-frame potential, measures moments of the pairwise overlaps. For each $k$, it is bounded from below by a constant depending on $d$ and $N$.

Of particular interest are cases in which the Welch inequality is saturated. For $k = 1$, equality is attained by tight frames, whose rescaled projectors define a generalized quantum measurement given by a positive-operator-valued measure (POVM) with rank-one elements. More generally, tight frames are spanning sets of vectors that may be overcomplete, with redundancy that can be exploited to enable stable data recovery over noisy or lossy signal transfer.

For a general $k$, the Welch inequality is saturated solely by sets of states featuring a remarkable property, namely forming a $k$-design [6, 7]. The latter have become central in quantum information, as they provide finite and highly symmetric surrogates of Haar-random states [8]. More precisely, a complex projective $k$-design is a finite ensemble of pure states whose average reproduces the Haar average of every polynomial of degree $k$ in the state amplitudes and degree $k$ in their complex conjugates. The Haar measure describes the uniform distribution over all pure states. State designs play a key role in quantum-state estimation problems [9]. The most well-known examples are: (i) complete sets of mutually unbiased bases (MUBs) with $N = d(d+1)$ (for prime-power dimensions) [10] and (ii) symmetric informationally complete (SIC) POVMs with $N = d^2$ elements [11]. Both families are 2-designs, and are central in optimal quantum state estimation [9, 10], quantum cryptography [12, 13], entanglement detection [14], and quantum foundations [15].

The saturation of Welch bounds thus provides a complete characterization of finite sets of states reproducing Haar moments up to order $k$, forming exact designs. This requires, however, very specific parameters; in particular, the cardinality $N$ of the set must be large enough, and the required cardinality grows very quickly with the dimension $d$ and $k$.

