arXiv:2603.21873v2 [quant-ph] 3 Aug 2026

Accurate ground state energy estimation with noise and imperfect state preparation

Alicja Dutkiewicz,[1,2] Thomas E. O'Brien,[3,2] and Stefano Polla[2,4]

[1] *QuSoft & CWI, Amsterdam, the Netherlands*
[2] *⟨aQa^L⟩ & Instituut-Lorentz, Universiteit Leiden, the Netherlands*
[3] *Google Quantum AI, Munich, Germany*
[4] *QuSoft, HIMS & IvI, University of Amsterdam, the Netherlands*

We introduce a classical estimator for the post-processing of quantum phase estimation (QPE) data when a single target phase is isolated within a known interval, as is typical of ground state energy estimation of gapped systems. Our estimator filters the QPE signal within this promise region and recovers the phase through a moment-projection routine, which is robust to both external spurious phases and experimental noise. In the noiseless case this achieves an exponential suppression of bias with respect to a naive mean estimator. In the presence of global depolarizing noise the bias is exponentially small in the circuit depth t, and the variance is $O(t^{-2}F^{-2})$ for circuit fidelity F. This improves by a factor of t^2 over a naive shifted-and-rescaled-mean approach. To mitigate realistic circuit-level noise, we combine our method with the explicit unbiasing scheme described in Ref. [1]. This yields an overhead interpolating between the F^{-4} scaling typical of explicitly unbiased error mitigation and a reduced F^{-2} scaling when the noise samples fall outside the promise interval. We validate our estimators on a small-scale simulation of the Ising model, observing better-than-expected performance for a global depolarizing noise approximation. This robustness to both multiple eigenvalues and realistic noise makes limited-depth phase estimation practical for early fault tolerant quantum experiments.

CONTENTS

I. Introduction 2
I. A. Summary of key results 2
II. Definitions 4
III. Background 7
III. A. QFT-based phase estimation 7
III. B. QSP-based phase estimation 9
IV. Results 10
IV. A. Moment projection estimator 10
IV. B. Noise unbiasing for moment projection 12
IV. C. Moment projection for phase estimation 13
IV. D. Gaussian kernel and no noise 16
IV. E. Gaussian kernel and global depolarising noise 18
IV. F. Gaussian kernel and arbitrary noise with noise unbiasing 21
V. Numerical demonstration 22
VI. Conclusion and outlook 25
Author contributions and acknowledgments 26
References 26
A. Proofs of Lemmas 17, 18, 20, and 21 29
B. Proofs of Lemmas 27 and 29 32