

Optimal Lower Bound for Ground-State Energy Estimation with a Guiding State

Rolando D. Somma* Ronald de Wolf†

arXiv:2608.24493v2 [quant-ph] 30 Sep 2026

Abstract

Suppose we can apply the unitary $U = e^{iH}$ for some Hamiltonian H, and are given access to a unitary that prepares a guiding state promised to have overlap at least $\gamma > 0$ with the ground space of H. Our goal is to estimate the ground-state energy of H within additive error $\delta > 0$ and success probability at least $1 - \varepsilon$, $\varepsilon > 0$. How many applications of U and its inverse U^{-1} are necessary and sufficient? This quantity corresponds to the total Hamiltonian-simulation time needed. An upper bound $O(\log(1/\varepsilon)\log(1/\gamma)/\gamma\delta)$ was known, and was improved to $O(\log(1/\varepsilon)/\gamma\delta)$ very recently [JW26]. A matching lower bound was known whenever one of the three parameters $\delta, \gamma, \varepsilon$ was held constant [MdW26]. In this paper we prove the joint lower bound $\Omega(\log(1/\varepsilon)/\gamma\delta)$ with the tight ε-dependence provided the dimension of H is at least $\log(1/\varepsilon)/\gamma^2$. Furthermore, we show that this same lower bound (with slightly larger dimension) holds for both the special case in which the ground state is guaranteed to be unique and H has a gap of δ between its first and second eigenvalue; and for ground-state preparation, where δ denotes the spectral gap and ε now is the approximation error. The lower bounds also apply when the Hamiltonian can be accessed via its block-encoding, and when fractional powers of U are allowed, as in continuous-time Hamiltonian simulation. Lastly, improved upper bounds are known when H is nonnegative and presented as a sum of squares; and our results imply the lower bound $\Omega(\log(1/\varepsilon)/\gamma\sqrt{\delta})$ for this case.

1 Introduction

1.1 Ground-state energy estimation with guiding states

One of the core problems in quantum chemistry is the following: given a classical description of some Hamiltonian H (e.g., an electronic structure Hamiltonian), estimate its *ground-state energy*, which is its smallest eigenvalue $\lambda_{\min}$. If H is normalized such that its eigenvalues are all in $[0, 2\pi - 2\delta)$ (the "-2δ" is to avoid the issue that 0 and 2π are the same around the unit circle) and we define the unitary $U = e^{iH}$ (which has the same eigenvectors as H, with eigenvalue λ of H becoming eigenvalue $e^{i\lambda}$ for U), then estimating the ground-state energy of H is equivalent to estimating the smallest eigen*phase* of U.

If we are additionally given a *ground state* (i.e., an eigenstate of H corresponding to $\lambda_{\min}$), then the well-known phase estimation algorithm [Kit95] is tailor-made to estimate the ground-state

*Google Quantum AI, Venice, CA 90291, United States.

†Google Quantum AI, Venice, CA 90291, United States. Also QuSoft, CWI and University of Amsterdam, the Netherlands. Partially supported by the Dutch Research Council (NWO) through Gravitation-grant Quantum Software Consortium, 024.003.037.

