



Optimal Quantum Algorithm for Ground-State Energy Estimation with a Guiding State

Stacey Jeffery[1,2] and Freek Witteveen[1]

[1]QuSoft & CWI, Amsterdam
[2]QLever & University of Amsterdam

arXiv:2608.24494v1 [quant-ph] 25 Aug 2026

Abstract

In the problem of ground-state energy estimation, one aims to estimate the smallest eigenvalue of a Hamiltonian, often given a guiding state, with some promised overlap γ with the ground space. The main approach to this problem is to simulate its evolution, and estimate the smallest (or equivalently, largest) eigenphase of the resulting unitary U. We give a quantum algorithm that estimates the largest eigenphase of a unitary U in this guided setting using a factor of $\log \frac{1}{\gamma}$ fewer queries to U than the previous best approach. The result matches an existing lower bound, and answers an open question from Mande and de Wolf. The algorithm is based on transducers, which often allow composition of quantum algorithms without overhead from error reduction.

1 Introduction

One of the most promising applications of quantum computers is the simulation of complex quantum systems that are relevant to physics and chemistry. A central task in this domain is to estimate the ground state energy of a Hamiltonian. There are various approaches to solving this on a quantum computer. For fault-tolerant quantum computation, the standard paradigm for computing ground state energies is based on quantum phase estimation (QPE), [Kit96, AL99], see [BBMC20, MEAG+20] for a discussion of the application to computational physics and chemistry. Here, we consider the problem where we would like to know the ground state energy E_0 of a Hamiltonian H to precision δ, and we assume that we are given the ability to prepare a state $|\psi\rangle$ that has overlap γ with the ground state. The computational problem is the *guided ground state energy problem*:

> Given a Hamiltonian H with eigen-decomposition
>
> $$H = \sum_{k=0}^{K-1} E_k \Pi_k, \qquad E_0 < E_1 < \cdots < E_{K-1}$$
>
> and a state-preparation unitary A that prepares a state $|\psi\rangle$ such that $\|\Pi_0|\psi\rangle\| \geq \gamma > 0$, estimate E_0 to precision $\delta > 0$.

Here, we have not specified how we have access to the Hamiltonian. What matters is that one can implement unitary time evolution on the quantum computer (which is possible in various access models to H), or encode the spectrum of H in a different way in a unitary operator. If we let $U = e^{-iH}$ be the time-evolution unitary, or alternatively if we let U be the qubitized walk operator associated to H, the guided ground state energy problem directly reduces to the following *max eigenphase estimation* problem: