

Optimal cloning of mixed states

Marco Fanizza* Dmitry Grinko† Thilo Scharnhorst‡ Jack Spilecki‡

arXiv:2608.27298v1 [quant-ph] 27 Aug 2026

Abstract

We consider the problem of *approximate cloning* of quantum states: given n copies of an unknown state $\rho \in \mathbb{C}^{d\times d}$, prepare an $(n+k)$-copy state with high fidelity to $\rho^{\otimes(n+k)}$. Werner's pure state cloner is the optimal channel for the pure state case, and shows that $n = \Theta(kd/\varepsilon)$ copies are necessary and sufficient to clone k additional copies of an unknown pure state to fidelity $1-\varepsilon$. The random purification channel gives a straightforward extension of Werner's cloner to mixed state inputs: given n copies of a mixed state, randomly purify your input, apply Werner's channel in the larger Hilbert space, and then trace out the auxiliary registers. This gives a mixed state cloner using $n = O(krd/\varepsilon)$ copies to clone rank-r states. Can one do any better? We show that the answer is no: one must use $n = \Omega(krd/\varepsilon)$ copies. We prove our lower bound by studying the special case of *projector cloning*, in which the input state ρ is promised to be of the form P/r, where P is a rank-r orthogonal projector.

As a further application of our techniques, we consider the closely related problem of *approximate transposition* of quantum states, where one seeks to convert $\rho^{\otimes n}$ to a k-copy state with high fidelity to $(\rho^T)^{\otimes k}$. Here, we again show $n = \Theta(krd/\varepsilon)$ copies are necessary and sufficient for this task.



*Inria, Télécom Paris – LTCI, Institut Polytechnique de Paris, Palaiseau, France. marco.fanizza@inria.fr

†QuSoft & Institute for Logic, Language and Computation & Korteweg-de Vries Institute for Mathematics, University of Amsterdam, The Netherlands. d.grinko@uva.nl

‡UC Berkeley. {thilo,jspilecki}@berkeley.edu