



Quantum Fourier transform toolbox

Carli Bruinsma[1,2,*], Pietro M. Posta[3,*], Joppe Stokvis[1,4,*],
Dmitry Grinko[1,2,5], Maris Ozols[1,2,5]

[1]QuSoft, Amsterdam, The Netherlands
[2]Institute for Logic, Language and Computation, University of Amsterdam, The Netherlands
[3]Department of Mathematical Sciences, University of Copenhagen, Denmark
[4]Mathematical Institute, Leiden University, The Netherlands
[5]Korteweg-de Vries Institute for Mathematics, University of Amsterdam, The Netherlands

arXiv:2608.28573v2 [quant-ph] 30 Sep 2026

Abstract

Quantum Fourier transforms (QFTs) are essential primitives in quantum algorithms. While abelian groups admit efficient QFT circuits, with circuit size polynomial in the logarithm of the group order, efficient constructions are known for relatively few non-abelian families. We develop two new approaches to QFT circuit construction, based on Mackey theory and Clifford theory, respectively, and use them to show exponential improvement in circuit cost for specific group families. Using the Mackey-theoretic approach, we obtain explicit quantum circuits for the QFT over $\mathrm{GL}_2(\mathbb{F}_q)$ that scale polynomially in $\log q$, rather than polynomially in q. Using the Clifford-theoretic approach, we obtain QFT circuits for wreath products $F \wr S_n$, whose cost depends on the cost of a QFT over F and the size of its representation registers. This removes the restriction $|F| = \mathrm{poly}(n)$ required by previous generic constructions and can yield exponential improvements when F itself has an efficient QFT. Together, these methods provide new systematic tools to construct QFTs for broad classes of finite groups.

Contents

1 Introduction **2**
1.1 Previous work . . . 3
1.2 Main results . . . 7
1.3 Organization of the paper . . . 8

2 Preliminaries and notation **9**
2.1 Representations, labels, and dual actions . . . 9
2.2 Regular representations and the Fourier transform . . . 9
2.3 Restriction and induction . . . 11
2.4 Successors and predecessors of a representation . . . 12

3 QFT using subgroups **12**
3.1 Bratteli diagram and Gelfand–Tsetlin basis . . . 13
3.2 Fourier transform for a group and a subgroup . . . 13
3.3 Induction along a multiplicity-free chain . . . 16

* These authors contributed equally
Email addresses: carlibruinsma@gmail.com, pmp@math.ku.dk, j.a.stokvis@math.leidenuniv.nl, d.grinko@uva.nl, and marozols@gmail.com.

