



A Complete and Natural Rule Set for Multi-Qudit Clifford Circuits in All Odd Prime Dimensions

Xiaoning Bian[1], Sarah Meng Li[2,3], Neil J. Ross[4], John van de Wetering[2,3], and Yuming Zhao[5]

[1]Tsinghua University, Beijing, China
[2]University of Amsterdam, Amsterdam, Netherlands
[3]QuSoft, Amsterdam, Netherlands
[4]Dalhousie University, Halifax, Canada
[5]University of Copenhagen, Copenhagen, Denmark

arXiv:2609.40106v2 [quant-ph] 5 Oct 2026

We present a complete set of 16 rewrite rules for multi-qudit Clifford circuits in all odd prime dimensions. Completeness means that any two Clifford circuits representing the same linear map can be rewritten into each other using these rules. Each rule involves at most three qudits and admits an intuitive interpretation. To establish completeness, we first work at the symplectic level, using the isomorphism between the symplectic group $\mathrm{Sp}(2n, \mathbb{Z}_p)$ and the quotient of the Clifford group by the Pauli group. We construct a circuit normal form that captures the stabiliser tableau of a Clifford operator and is unique up to Pauli correction. Using this normal form, we derive a complete set of symplectic relations, which we then lift to Clifford relations by incorporating Pauli corrections. We also formally verify completeness in the Agda proof assistant.

Contents

1 Introduction **2**
1.1 Main Results . . . 3
1.2 Proof Overview . . . 5
1.3 Paper Organisation . . . 7

2 Preliminaries **7**
2.1 Rings . . . 7
2.2 Operators . . . 8
2.2.1 Pauli Operators . . . 9
2.2.2 Clifford Operators . . . 9
2.2.3 Clifford Conjugation of Pauli Generators . . . 11
2.2.4 Symplectic Structure of the Clifford Group . . . 13
2.2.5 Derived Generators . . . 13

Xiaoning Bian: bian@dal.ca
Sarah Meng Li: sarah.li@uva.nl
Neil J. Ross: neil.jr.ross@dal.ca
John van de Wetering: john@vdwetering.name
Yuming Zhao: yuming@math.ku.dk

