

2022–2023



Testing bosonic and fermionic Gaussian states with a constant number of copies

Jop Briët[*] Davi Castro-Silva[†] Michael Walter[‡] Freek Witteveen[§]

arXiv:2610.06794v1 [quant-ph] 5 Oct 2026

Abstract

Bosonic and fermionic Gaussian quantum states are families of quantum states that play a central role in quantum physics and quantum information processing. We study the problem of testing whether a given state is either Gaussian, or has bounded overlap with any Gaussian state. Previous analyses of this testing problem led to tests that require a number of copies of the state that scales with the number of modes (the system size). In this work we show that a *constant* number of copies of the state suffices. We derive these results based on the symmetries of identical copies of Gaussian states, and by bounding the distance to the set of Gaussian states through a gradient flow and a spectral bound studied previously for the Kac random walk on a sphere.

1 Introduction

Gaussian quantum states are an important set of parametrized quantum states. They are relevant as ground states of quadratic Hamiltonians in quantum optics and material science, as a tractable set of quantum states for approximation or simulation by classical computation, and are often used in quantum communication protocols. As the other side of the coin, *non-Gaussian* states also play a central role: they appear as ground states of more complicated Hamiltonians, and as a resource for quantum computation beyond classical simulability. This motivates the following basic question: given copies of a state $|\psi\rangle$, can one do a measurement that determines whether the state is Gaussian, or has bounded maximal overlap with all Gaussian states? This question has been studied in previous works [1–4]. These works led to protocols that resolve this question, provided access to a number of copies of the state that scales with the number of bosonic or fermionic modes.

The present work shows that a *constant* number of copies suffices.

Theorem 1.1 (Testing Gaussianity with constant sample complexity)**.** *There are tests that use $O(\frac{1}{\varepsilon}\log\frac{1}{\delta})$ copies of a state $|\psi\rangle$, and determine, with probability of failure at most δ, whether $|\psi\rangle$ is a fermionic or bosonic Gaussian state, or $|\langle\phi|\psi\rangle|^2 \leq 1-\varepsilon$ for all Gaussian states $|\phi\rangle$.*

We also give tests for mean-zero bosonic states, and for coherent states. The tests act on two copies at a time when testing fermionic, mean-zero bosonic Gaussian states, or coherent states. The test for bosonic Gaussian states uses four copies at a time.

The test is based on a well-known characterization of Gaussian states, based on their invariance properties under phase space rotations. For fermionic Gaussian states and bosonic mean-zero Gaussian states, we consider a two-copy test with accepting operator P, which projects onto states which are invariant under rotations (beam-splitters of arbitrary angles) between the two copies. This measurement can be efficiently implemented with a quantum circuit: [3, 5] gives a measurement scheme in terms of Bell sampling, and [6] gives an implementation in the bosonic case. For the case of non-zero mean, there is an easy trick to get a new state from two copies that has zero mean, by applying a beam-splitter and discarding one copy. This

[*]CWI & QuSoft, Amsterdam, University of Leiden j.briet@cwi.nl
[†]Department of Computer Science, University of Warwick, davi.de-castro-silva@warwick.ac.uk
[‡]LMU Munich & MCQST and University of Amsterdam & QuSoft, michael.walter@lmu.de
[§]CWI & QuSoft, Amsterdam, f.witteveen@cwi.nl

