

Sustainable trees

Ferenc Bencs* Maarten van Dijk† Krystal Guo‡

October 12, 2026

arXiv:2610.17078v1 [math.CO] 12 Oct 2026

Abstract

Let X be a connected graph and let $\lambda_1(X)$ denote the spectral radius of its adjacency matrix. An edge e of X is *sustainable* if it can be deleted and replaced by a non-edge so that the resulting graph is connected and its spectral radius does not decrease; X is *sustainable* if every edge is. We study this edgewise notion of spectral robustness for trees, where each replacement is an elementary exchange between trees on the same vertex set. We establish basic properties of sustainability and exhibit infinite families of sustainable and non-sustainable trees. Our main results concern trees with a vertex of large degree: if T is a tree on n vertices with $\Delta(T) \geq n/2$, then every vertex of maximum degree is a principal vertex, that is, a vertex at which the Perron vector is maximal. Building on this, we prove that every tree with $\Delta(T) \geq n/2 + 1$ is non-sustainable, and we classify the sustainable trees in the boundary case $\Delta(T) = n/2$: for $n \geq 4$, the only such trees are the double broom $B(2, n/2, n/2 - 1)$ and one exceptional tree on eight vertices. We conclude with open problems on the generalized tree-shift order and the asymptotic proportion of sustainable trees.

Keywords: spectral radius, trees, Perron vector, principal vertex.

MSC2020: Primary 05C50; Secondary 05C05, 15A18.

1 Introduction

Let X be a connected graph, and let $\lambda_1(X)$ denote the spectral radius of its adjacency matrix. For an edge $e \in E(X)$, consider the operation of deleting e and inserting a non-edge $f \in E(\overline{X})$, subject to the requirement that the resulting graph remains connected. We say that the edge e is*sustainable* if at least one such replacement satisfies

$$\lambda_1(X - e + f) \geq \lambda_1(X),$$

and we call X *sustainable* if every edge of X is sustainable. Thus a non-sustainable graph contains a spectrally indispensable edge: an edge for which every admissible one-edge replacement strictly decreases the spectral radius.

*Centrum Wiskunde & Informatica, Amsterdam, The Netherlands. Supported by the Netherlands Organisation of Scientific Research (NWO): VI.Veni.222.303., Email: f.bencs@gmail.com

†Email: m.z.j.vandijk00@gmail.com

‡Korteweg-de Vries Institute for Mathematics and QuSoft, University of Amsterdam, Amsterdam, The Netherlands. Email: k.guo@uva.nl

