

2015–
2020



Bounds on entanglement dimensions and quantum graph parameters via noncommutative polynomial optimization

Sander Gribling* David de Laat† Monique Laurent‡

arXiv:1708.09696v2 [math.OC] 9 Jan 2018

Abstract

In this paper we study optimization problems related to bipartite quantum correlations using techniques from tracial noncommutative polynomial optimization. First we consider the problem of finding the minimal entanglement dimension of such correlations. We construct a hierarchy of semidefinite programming lower bounds and show convergence to a new parameter: the minimal average entanglement dimension, which measures the amount of entanglement needed to reproduce a quantum correlation when access to shared randomness is free. Then we study optimization problems over synchronous quantum correlations arising from quantum graph parameters. We introduce semidefinite programming hierarchies and unify existing bounds on quantum chromatic and quantum stability numbers by placing them in the framework of tracial optimization.

1 Introduction

1.1 Bipartite quantum correlations

One of the distinguishing features of quantum mechanics is quantum entanglement, which allows for nonclassical correlations between spatially separated parties. In this paper we consider the problems of quantifying the advantage entanglement can bring (first investigated through Bell inequalities in the seminal work [3]) and quantifying the minimal amount of entanglement necessary for generating a given correlation (initiated in [5] and continued, e.g., in [38, 54, 47]).

Quantum entanglement has been widely studied in the bipartite correlation setting (for a survey, see, e.g., [39]). Here we have two parties, Alice and Bob, where Alice receives a question s taken from a finite set S and Bob receives a question t taken from a finite set T. The parties do not know each other's questions, and after receiving the questions they do not communicate. Then, according to some predetermined protocol, Alice returns an answer a from a finite set A and Bob returns an answer b from a finite set B. The probability that the parties answer (a, b) to questions (s, t) is given by a *bipartite correlation* $P(a, b|s, t)$, which satisfies $P(a, b|s, t) \geq 0$ for all $(a, b, s, t) \in \Gamma$ and $\sum_{a,b} P(a, b|s, t) = 1$ for all $(s, t) \in S \times T$. Throughout we set $\Gamma = A \times B \times S \times T$. Which bipartite correlations $P = (P(a, b|s, t)) \in \mathbb{R}^\Gamma$ are possible depends on the additional resources available to the two parties Alice and Bob.

If the parties do not have access to additional resources, then the correlation is *deterministic*, which means it is of the form $P(a, b|s, t) = P_A(a|s)\, P_B(b|t)$, with $P_A(a|s)$ and $P_B(b|t)$ taking values in $\{0, 1\}$ and $\sum_a P_A(a|s) = \sum_b P_B(b|t) = 1$ for all s, t. If the parties have access to local randomness, then P_A and P_B take values in $[0, 1]$. If the parties have access to shared randomness, then the resulting correlation is a convex combination of deterministic correlations

*CWI and QuSoft, Amsterdam, the Netherlands. Supported by the Netherlands Organization for Scientific Research, grant number 617.001.351. gribling@cwi.nl

†CWI and QuSoft, Amsterdam, the Netherlands. Supported by the Netherlands Organization for Scientific Research, grant number 617.001.351, and by the ERC Consolidator Grant QPROGRESS 615307. mail@daviddelaat.nl

‡CWI and QuSoft, Amsterdam, and Tilburg University, Tilburg, the Netherlands. laurent@cwi.nl