

2016–
2021

Many-body effects in quantum metrology

Jan Czajkowski,[1] Krzysztof Pawłowski,[2] and Rafał Demkowicz-Dobrzański[3]

[1] *QuSoft, University of Amsterdam, Institute for Logic, Language and Computation (ILLC), P.O. Box 94242, 1090 GE Amsterdam, The Netherlands*
[2] *Center for Theoretical Physics, Polish Academy of Sciences, Al. Lotników 32/46, 02-668 Warsaw, Poland*
[3] *Faculty of Physics, University of Warsaw, ul. Pasteura 5, PL-02-093 Warszawa, Poland*

We study the impact of many-body effects on the fundamental precision limits in quantum metrology. On the one hand such effects may lead to non-linear Hamiltonians, studied in the field of non-linear quantum metrology, while on the other hand they may result in decoherence processes that cannot be described using single-body noise models. We provide a general reasoning that allows to predict the fundamental scaling of precision in such models as a function of the number of atoms present in the system. Moreover, we describe a computationally efficient approach that allows for a simple derivation of quantitative bounds. We illustrate these general considerations by a detailed analysis of fundamental precision bounds in a paradigmatic atomic interferometry experiment with standard linear Hamiltonian but with *both* single and two-body losses taken into account—a model which is motivated by the most recent Bose-Einstein Condensate (BEC) magnetometry experiments. Using this example we also highlight the impact of the atom number super-selection rule on the possibility of protecting interferometric protocols against decoherence.

arXiv:1810.03651v2 [quant-ph] 17 Oct 2018

I. INTRODUCTION

The research effort in the field of quantum metrology [1–4] goes along two main directions. The first is the experimental one, with the ultimate goal of designing metrological setups that harness quantum behavior of light and matter while at the same time minimize unwanted effects of decoherence in order to reach the precision regime inaccessible within the classical paradigm [5–9]. The theoretical effort, on the other hand, aims at identifying new promising proposals for quantum metrological protocols as well as fundamental bounds on precision that can in principle be achieved in quantum systems. The bounds are derived using simplified models that are intended to capture the essential features of the relevant physical systems and take into account the dominant sources of decoherence [10–17].

Typically, a significant discrepancy is observed between the fundamental bounds arising from theoretical considerations and the performance of practical setups. This is due to either oversimplified theoretical modeling of important physical effects present in the experiment or experimental inability of preparing the optimal quantum states and implementing the optimal quantum operations required by the theory. A noticeable exception from this rule is the case of quantum enhanced optical interferometry, where the performance of squeezed-light enhanced gravitation wave detector devices [18, 19] approaches closely the fundamental limit predicted by the theory [20].

Atomic systems are inherently more complex than purely optical ones, and hence the theoretical analysis of fundamental metrological bounds is much more challenging. While in optical interferometry the dominant source of decoherence is loss, which has single-body character, in atomic systems interactions between atoms naturally lead to various many-body effects. In cold atom experiments, apart from single-particle losses, which in principle might be described using a similar formalism as in the optical case, two- and three-body losses play a crucial role in the system dynamics. This poses a serious challenge to theorists, as the most powerful theoretical methods developed in the field of quantum metrology are naturally tailored to deal with uncorrelated, or in other words, single-body noise [12, 13, 21].

In this paper, we substantially develop the ideas from [16], where it was proposed to view the evolution of an N-particle permutationally invariant system with at most n-body interactions as equivalent to an application series of n-particle quantum channels to n-element subsets of N particles. In this way, one is able to make use of theoretical methods, developed with single-body noise models in mind, in many-body noise cases, without the need to consider the full N-particle space but rather a much smaller n-particle space. In what follows we will call this the *reduced particle number* (RPN) approach. In [16] this idea was used to obtain scalings of precision bounds in cases of a k-body Hamiltonian and l-body noise processes. Here we generalize these considerations to a situation where both Hamiltonian and the noise part contains many terms with different level of non-linearity and provide a clear recipe for how to proceed in order to obtain the eventual fundamental precision scaling. Furthermore, we discuss the way to obtain explicit *quantitative* bounds using this approach, and apply it to find limits on precision of atomic interferometry in presence of both