


2016–
2021

Conditions for superdecoherence

Joris Kattemölle[1,2] and Jasper van Wezel[1,2]

[1]Institute for Theoretical Physics, University of Amsterdam, Science Park 904, Amsterdam, The Netherlands

[2]QuSoft, CWI, Science Park 123, Amsterdam, The Netherlands

arXiv:1912.04266v5 [quant-ph] 7 May 2020

Decoherence is the main obstacle to quantum computation. The decoherence rate per qubit is typically assumed to be constant. It is known, however, that quantum registers coupling to a single reservoir can show a decoherence rate per qubit that increases linearly with the number of qubits. This effect has been referred to as superdecoherence, and has been suggested to pose a threat to the scalability of quantum computation. Here, we show that superdecoherence is absent when the spectrum of the single reservoir is continuous, rather than discrete. The reason of this absence, is that, as the number of qubits is increased, a quantum register inevitably becomes susceptible to an ever narrower bandwidth of frequencies in the reservoir. Furthermore, we show that for superdecoherence to occur in a reservoir with a discrete spectrum, one of the frequencies in the reservoir has to coincide exactly with the frequency the quantum register is most susceptible to. We thus fully resolve the conditions that determine the presence or absence of superdecoherence. We conclude that superdecoherence is easily avoidable in practical realizations of quantum computers.

1 Introduction

In principle, quantum computers can solve problems that are intractable on any classical computer. The largest obstacle to bringing this in practice is decoherence [1], and it is essential to understand the sources and effects of decoherence under practical circumstances encountered in actual quantum computers. As we inch towards full-scale quantum computing, where we are already facing systems with on the order of a hundred qubits [2–4], the system size dependence of decoherence becomes of increasing importance.

Decoherence is commonly studied in a simplified spin-boson model, where only the dephasing effects of the bosonic bath are taken into account [5–14]. Henceforth we will refer to this model as simply 'the dephasing model'. This model is exactly solvable, and at the same time broadly relevant because dephasing times are typically much shorter than relaxation times [7, 15, 12]. It should be noted, however, that there are situations where it does not accurately describe the decoherence process because of non-perturbative effects [15, 12]. If, in the dephasing model, each qubit is assumed to couple to its own, independent reservoir, the decoherence rate per qubit is constant. If, on the other hand, the qubits couple to single reservoir, the decoherence rate per qubit scales linearly with the number of qubits for certain states [6, 16, 17, 7, 8, 18, 19]. This effect has been referred to as superdecoherence, in analogy with superradiance.

Superdecoherence has been predicted in Refs. [7, 15], and has been observed experimentally in an ion-trap quantum computer [20]. Although some states suffer superdecoherence, the probability of running into such a state during the course of an actual algorithm may be extremely small [17]. Additionally, if the decoherence is dominated by relaxation, rather than dephasing, is has been shown that superdecoherence does not occur for the Greenberger-Horne-Zeilinger (GHZ) and the Hadamard state [21]. Also the particular model of solid-state qubits coupling to a single phonon reservoir has been shown not to give rise to superdecoherence [22]. The latter approach focuses on a specific setting of the dephasing model: the geometry of the quantum register is assumed to be a linear array, and the phonon reservoir is assumed to be three-dimensional and thermal, with a continuous spectrum and a linear dispersion relation. Therefore, it is unable to reveal the general underlying physical reasons for the absence of superdecoherence. The reason why superdecoherence emerges in other settings of the single-reservoir dephasing model remained unknown.

Here, we fully resolve the physical conditions that determine the presence or absence of superdecoherence in the dephasing model, with all qubits coupling to a single bath. We do not make any assumptions about the geometry of the quantum register, the dimension d, the reservoir dispersion relation, or the directional dependence of the spin-boson interaction. For the reservoir state, we assume a very general initial condition that applies to practically relevant situations. In this general setting, we find that the (im)possibility of superdecoherence due to a single reservoir is completely determined by the boundedness of the *spectral density* and the *occupation density* of the reservoir.

The spectral density is the density of modes at a