



QuSoft is named on page 18 of the paper.

The border support rank of two-by-two matrix multiplication is seven

Markus Bläser, Matthias Christandl and Jeroen Zuiddam

arXiv:1705.09652v1 [cs.CC] 26 May 2017

Abstract

We show that the border support rank of the tensor corresponding to two-by-two matrix multiplication is seven over the complex numbers. We do this by constructing two polynomials that vanish on all complex tensors with format four-by-four-by-four and border rank at most six, but that do not vanish simultaneously on any tensor with the same support as the two-by-two matrix multiplication tensor. This extends the work of Hauenstein, Ikenmeyer, and Landsberg. We also give two proofs that the support rank of the two-by-two matrix multiplication tensor is seven over any field: one proof using a result of De Groote saying that the decomposition of this tensor is unique up to sandwiching, and another proof via the substitution method. These results answer a question asked by Cohn and Umans. Studying the border support rank of the matrix multiplication tensor is relevant for the design of matrix multiplication algorithms, because upper bounds on the border support rank of the matrix multiplication tensor lead to upper bounds on the computational complexity of matrix multiplication, via a construction of Cohn and Umans. Moreover, support rank has applications in quantum communication complexity.

1. Introduction

Multiplication of two $n \times n$ matrices over a field $\mathbb{F}$ is an $\mathbb{F}$-bilinear map $\mathbb{F}^{n\times n} \times \mathbb{F}^{n\times n} \to \mathbb{F}^{n\times n}$ called the matrix multiplication map. The matrix multiplication map corresponds naturally to the following structure tensor. Let $[n]$ be the set $\{1, 2, \ldots, n\}$ and let $\{e_{ij} : i, j \in [n]\}$ be the standard basis for the vector space $\mathbb{F}^{n\times n}$ of $n \times n$ matrices. Define the structure tensor of the matrix multiplication map as

$$\langle n, n, n\rangle \coloneqq \sum_{i,j,k\in[n]} e_{ij} \otimes e_{jk} \otimes e_{ki} \in \mathbb{F}^{n\times n} \otimes \mathbb{F}^{n\times n} \otimes \mathbb{F}^{n\times n}.$$

(Technically, this is the structure tensor of the trilinear map that computes the trace of a product of three matrices.) Let V_1, V_2, and V_3 be vector spaces. The tensor rank of a tensor $t \in V_1 \otimes V_2 \otimes V_3$ is the smallest number r such that t can be written as a sum of r simple tensors $v_1 \otimes v_2 \otimes v_3 \in V_1 \otimes V_2 \otimes V_3$.