



QuSoft is named on page 11 of the paper.

arXiv:1804.00431v2 [math.RT] 23 Jan 2019

HORN INEQUALITIES AND QUIVERS

V. BALDONI, M. VERGNE, AND M. WALTER

Abstract. Let G be a complex reductive group acting on a finite-dimensional complex vector space $\mathcal{H}$. Let B be a Borel subgroup of G and let T be the associated torus. The Mumford cone is the polyhedral cone generated by the T-weights of the polynomial functions on $\mathcal{H}$ which are semi-invariant under the Borel subgroup. In this article, we determine the inequalities of the Mumford cone in the case of the linear representation of the group $G = \prod_x \mathrm{GL}(n_x)$ associated to a quiver and a dimension vector $n = (n_x)$. We give these inequalities in terms of filtered dimension vectors, and we directly adapt Schofield's argument to inductively determine the dimension vectors of general subrepresentations in the filtered context. In particular, this gives one further proof of the Horn inequalities for tensor products.

1. A generalized Horn condition

Let G be a connected complex reductive algebraic group, K a maximal compact subgroup of G and T be a maximal torus of K. Let $\mathfrak{t}$ denote the Lie algebra of T, and $\Lambda \subset i\mathfrak{t}^*$ be the lattice of weights of T. Let us choose a Borel subgroup B of G which contains T. We denote the set of associated positive roots by R_G^+. This determines a closed Weyl chamber $i\mathfrak{t}^*_{\geq 0}$ and the set P_G of dominant weights. We denote by V_λ the irreducible representation of G associated to $\lambda \in P_G$.

Now let $\Pi\colon G \to \mathrm{GL}(\mathcal{H})$ be a linear representation of G on a finite-dimensional complex vector space $\mathcal{H}$. We denote by π the corresponding infinitesimal representation of the Lie algebra of G. We assume that the action of G in the space $R(\mathcal{H})$ of polynomial functions on $\mathcal{H}$ decomposes with finite multiplicities. This is the case, e.g., when the action includes the homotheties. Consider the *Mumford cone* $\mathrm{Cone}(G, \mathcal{H}) \subset i\mathfrak{t}^*$, defined as the cone generated by the dominant weights λ such that $V_\lambda \subseteq R(\mathcal{H})$. The equation $V_\lambda \subseteq R(\mathcal{H})$ means that there exists a nonzero polynomial function on $\mathcal{H}$, which is semi-invariant under the action of B and of weight λ under T. The cone $\mathrm{Cone}(G, \mathcal{H})$ is a polyhedral cone determined by a finite number of inequalities. By Mumford's theorem [7], it can also be described using the moment map.

Example 1.1. *Consider $\mathcal{H} = \bigoplus_{i=1}^s \mathrm{Hom}(V_{x_i}, V_y)$, where V_{x_i}, V_y are n-dimensional vector spaces. It has a natural action of $G = \prod_{i=1}^s \mathrm{GL}(n) \times \mathrm{GL}(n)$. The cone $\mathrm{Cone}(G, \mathcal{H})$ is generated by the tuples of dominant weights $(\lambda_1, \dots, \lambda_s, \mu)$ associated to Young diagrams such that $V_\mu \subseteq$*