


2014–
2018



QuSoft is named on page 22 of the paper.

arXiv:1905.03148v1 [math.CO] 8 May 2019

THE ASYMPTOTIC INDUCED MATCHING NUMBER OF HYPERGRAPHS: BALANCED BINARY STRINGS

SRINIVASAN ARUNACHALAM, PÉTER VRANA, AND JEROEN ZUIDDAM

Abstract. We compute the asymptotic induced matching number of the k-partite k-uniform hypergraphs whose edges are the k-bit strings of Hamming weight $k/2$, for any large enough even number k. Our lower bound relies on the higher-order extension of the well-known Coppersmith–Winograd method from algebraic complexity theory, which was proven by Christandl, Vrana and Zuiddam. Our result is motivated by the study of the power of this method as well as of the power of the Strassen support functionals (which provide upper bounds on the asymptotic induced matching number), and the connections to questions in tensor theory, quantum information theory and theoretical computer science.

Phrased in the language of tensors, as a direct consequence of our result, we determine the asymptotic subrank of any tensor with support given by the aforementioned hypergraphs. In the context of quantum information theory, our result amounts to an asymptotically optimal k-party stochastic local operations and classical communication (slocc) protocol for the problem of distilling GHZ-type entanglement from a subfamily of Dicke-type entanglement.

Keywords. k-partite k-uniform hypergraphs, asymptotic induced matchings, higher-order Coppersmith–Winograd method

1. Introduction

1.1. **Problem.** We study in this paper an asymptotic parameter of k-partite k-uniform hypergraphs: the asymptotic induced matching number. For $k \in \mathbb{N}$, a *k-partite k-uniform hypergraph*, or *k-graph* for short, is a tuple of finite sets $V_1, \dots, V_k$ together with a subset Φ of their cartesian product:

$$\Phi \subseteq V_1 \times \cdots \times V_k.$$

Whenever possible we will leave the vertex sets V_i implicit and refer to the k-graph by its edge set Φ. For any $k \in \mathbb{N}$ we use the notation $[k] \coloneqq \{1, 2, \dots, k\}$. Let Φ be a k-graph. We say a subset Ψ of Φ is *induced* if $\Psi = \Phi \cap (\Psi_1 \times \cdots \times \Psi_k)$ where for each $i \in [k]$ we define the marginal set $\Psi_i \coloneqq \{a_i : a \in \Psi\}$. We call Ψ a *matching* if any two distinct elements $a, b \in \Psi$ are distinct in all k coordinates, that is, $\forall i \in [k] : a_i \neq b_i$. The *subrank*[1] or *induced matching number* $\mathrm{Q}(\Phi)$ is defined as the size of the largest subset Ψ of Φ that is an induced matching, that is,

$$\mathrm{Q}(\Phi) \coloneqq \max\{|\Psi| : \Psi \subseteq \Phi, \Psi = \Phi \cap (\Psi_1 \times \cdots \times \Psi_k), \forall a \neq b \in \Psi \ \forall i \in [k]\ a_i \neq b_i\}.$$

For example, consider the 3-graph $\Phi = \{(1,1,1), (2,2,2), (3,3,3)\} \subseteq [3] \times [3] \times [3]$. Here Φ is itself an induced matching, and so $\mathrm{Q}(\Phi) = 3$. Next, let $\Phi = \{(1,1,1), (2,2,2), (3,3,3), (1,2,3)\}$. Now the subset $\{(1,1,1), (2,2,2)\} \subseteq \Phi$ is an induced matching and there is no larger induced matching in Φ, and so $\mathrm{Q}(\Phi) = 2$.

We define the *Kronecker product* of two k-graphs $\Phi \subseteq V_1 \times \cdots \times V_k$ and $\Psi \subseteq W_1 \times \cdots \times W_k$ as the k-graph

$$\Phi \boxtimes \Psi \coloneqq \big\{\big((a_1, b_1), \dots, (a_k, b_k)\big) : a \in \Phi, b \in \Psi\big\}$$

Date: May 9, 2019.

[1]The term subrank originates from an analogous parameter in the theory of tensors, see Section 1.4.1.