



QuSoft is named on page 20 of the paper.

arXiv:2004.14872v2 [math.AG] 24 Oct 2022

MINIMAL LENGTH IN AN ORBIT CLOSURE AS A SEMICLASSICAL LIMIT

COLE FRANKS AND MICHAEL WALTER

Abstract. Consider the action of a connected complex reductive group on a finite-dimensional vector space. A fundamental result in invariant theory states that the orbit closure of a vector v is separated from the origin if and only if some homogeneous invariant polynomial is nonzero on v, i.e. v is not in the null cone. Thus, efficiently finding the minimum distance between the orbit closure and the origin can lead to deterministic algorithms for null cone membership, an important polynomial identity testing problem including the non-commutative Edmonds problem. This connection to optimization has recently led to efficient algorithms for many problems in invariant theory.

Here we explore a refinement of the famous duality between orbit closures and invariant polynomials, which holds that the following two quantities coincide: (1) the logarithm of the Euclidean distance between the orbit closure and the origin and (2) the rate of exponential growth of the *invariant part* of $v^{\otimes k}$ in the semiclassical limit as k tends to infinity. This result can be deduced from work of S. Zhang (*Geometric reductivity at Archimedean places*, 1994), which uses sophisticated tools in arithmetic geometry. We provide a new and independent elementary proof inspired by the Fourier-analytic proof of the local central limit theorem. We generalize the result to projections onto highest weight vectors and isotypical components, and explore connections between such semiclassical limits and the asymptotic behavior of multiplicities in representation theory, large deviations theory in classical and quantum statistics, and the Jacobian conjecture as reformulated by Mathieu. Our formulas imply that they can be computed, in many cases efficiently, to arbitrary precision.

1. Introduction

Consider a vector v in a finite-dimensional representation space V of a connected complex reductive group G. Let V carry an inner product invariant under a maximally compact subgroup $K \subseteq G$. A fundamental result in geometric invariant theory states that the orbit closure of v under G contains the zero vector (v is *unstable*) if and only if every nonconstant homogeneous G-invariant polynomial vanishes on v (v is in the *null cone*) [27, 37]. In this manuscript we prove quantitative sharpenings and extensions of this relationship.

We quantify the evaluations of invariant polynomials in degree k by the *invariant part* of $v^{\otimes k}$, that is, its image under the orthogonal projection Π_k onto the invariant subspace of $V^{\otimes k}$. The sequence $\|\Pi_k v^{\otimes k}\|$ is super-multiplicative, so it is natural to study the following semiclassical limit:[1]

$$\limsup_{k\to\infty} \|\Pi_k v^{\otimes k}\|^{\frac{1}{k}} \tag{1.1}$$

Because $v^{\otimes k}$ is an element of the space of symmetric tensors, we may think of $\|\Pi_k v^{\otimes k}\|^2$ as a sum of squares $\sum_p |p(v)|^2$ of evaluations of an orthonormal basis $\{p\}$ of G-invariant homogeneous polynomials of degree k. Equivalently, it is equal to $\mathbb{E}|P(v)|^2$ where P is a random homogeneous G-invariant polynomial of degree k drawn from the complex standard Gaussian distribution. Thus, Eq. (1.1) measures the asymptotic behavior of *random* invariant polynomials.

[1] Considering the lim sup rather than the limit is necessary only because the sequence may vanish outside an additive subsemigroup of $\mathbb{N}$. By Fekete's lemma, there is $a \in \mathbb{N}$ such that restricted to the subsemigroup $\{ka : k \in \mathbb{N}\}$ the sequence $\|\Pi_k v^{\otimes k}\|^{\frac{1}{k}}$ has a limit and outside of which $\|\Pi_k v^{\otimes k}\|^{\frac{1}{k}}$ eventually vanishes. See also Lemma 2.3.