
2018–
2020


2019–
2023





QuSoft is named on page 33 of the paper.

arXiv:2008.12110v1 [math.OC] 27 Aug 2020

INTERIOR-POINT METHODS FOR UNCONSTRAINED GEOMETRIC PROGRAMMING AND SCALING PROBLEMS

PETER BÜRGISSER, YINAN LI, HAROLD NIEUWBOER, AND MICHAEL WALTER

Abstract. We provide a condition-based analysis of two interior-point methods for unconstrained geometric programs, a class of convex programs that arise naturally in applications including matrix scaling, matrix balancing, and entropy maximization. Our condition numbers are natural geometric quantities associated with the Newton polytope of the geometric program, and lead to diameter bounds on approximate minimizers. We also provide effective bounds on the condition numbers both in general and under combinatorial assumptions on the Newton polytope. In this way, we generalize the iteration complexity of recent interior-point methods for matrix scaling and matrix balancing. Recently, there has been much work on algorithms for certain optimization problems on Lie groups, known as capacity and scaling problems. For commutative groups, these problems reduce to unconstrained geometric programs, which serves as a particular source of motivation for our work.

1. Introduction

Geometric programming is an optimization paradigm that generalizes linear programming and has a wide range of applications [15, 6]. In this paper, we are concerned with *unconstrained geometric programs*. These are optimization problems of the form

$$\begin{aligned} &\text{minimize} && f(z) \\ &\text{subject to} && z \in \mathbb{R}^n,\ z > 0, \end{aligned} \tag{1.1}$$

where $f(z)$ is a *posynomial* in positive real variables $z_1, \dots, z_n$. That is,

$$f(z) = \sum_{i=1}^{k} q_i z^{\omega_i} = \sum_{i=1}^{k} q_i \prod_{j=1}^{n} z_j^{\omega_{i,j}}, \tag{1.2}$$

where the coefficients q_i are positive and the exponents $\omega_{i,j}$ are real numbers. In a general geometric program (GP), one adds posynomial inequality and monomial equality constraints. Although posynomials are non-convex in general, they are convex in x after the change of variables $z = e^x$. As such, they are the simplest family of *geodesically convex* programming problems. It is well-known that standard convex programming techniques, like the ellipsoid method or interior-point methods, can solve GP in polynomial time, see for instance [36, 29, 4, 7, 24, 39, 40, 34, 27]. Furthermore, it has been observed that interior-point methods are also efficient and robust in practice (cf. [7, 6]). However, explicit iteration complexity bounds are not readily available in the current literature.

In this paper, we present a detailed analysis of two interior-point algorithms for unconstrained GP in terms of natural geometric condition numbers. Our first algorithm applies to instances that (roughly speaking) have a well-conditioned Newton polytope, while our second algorithm has no such assumption but instead relies on