

2022–
2023

QuSoft is named on page 47 of the paper.

arXiv:2102.10018v3 [math.CO] 9 May 2023

Geometrical sets with forbidden configurations

Davi Castro-Silva

May 10, 2023

Abstract

Given finite configurations $P_1, \dots, P_n \subset \mathbb{R}^d$, let us denote by $\mathbf{m}_{\mathbb{R}^d}(P_1, \dots, P_n)$ the maximum density a set $A \subseteq \mathbb{R}^d$ can have without containing congruent copies of any P_i. We will initiate the study of this geometrical parameter, called the *independence density* of the considered configurations, and give several results we believe are interesting. For instance we show that, under suitable size and non-degeneracy conditions, $\mathbf{m}_{\mathbb{R}^d}(t_1P_1, t_2P_2, \dots, t_nP_n)$ progressively 'untangles' and tends to $\prod_{i=1}^n \mathbf{m}_{\mathbb{R}^d}(P_i)$ as the ratios t_{i+1}/t_i between consecutive dilation parameters grow large; this shows an exponential decay on the density when forbidding multiple dilates of a given configuration, and gives a common generalization of theorems by Bourgain and by Bukh in geometric Ramsey theory. We also consider the analogous parameter $\mathbf{m}_{\mathbb{S}^d}(P_1, \dots, P_n)$ in the more complicated framework of sets on the unit sphere $\mathbb{S}^d$, obtaining the corresponding results in this setting.

1 Introduction

The general problem we consider in this paper can be phrased by the following question: how large can a set be if it does not contain a given geometrical configuration?

The simplest and most well-studied instance of this problem concerns forbidden configurations of only two points on $\mathbb{R}^d$, which are then characterized by their distance; since there clearly exist unbounded sets on $\mathbb{R}^d$ which do not span a given distance, the appropriate notion of 'largeness' must take into account their density rather than their cardinality or measure. Define the *upper density* $\overline{d}(A)$ of a measurable set $A \subseteq \mathbb{R}^d$ by

$$\overline{d}(A) = \limsup_{T \to \infty} \frac{\mathrm{vol}(A \cap [-T, T]^d)}{\mathrm{vol}([-T, T]^d)},$$

where vol denote the Lebesgue measure. Our general problem in this case becomes: what is the maximum upper density that a subset of $\mathbb{R}^d$ can have if it does not contain pairs of points at distance 1?[1]

This extremal density is commonly denoted $m_1(\mathbb{R}^d)$, and it is associated to the measurable chromatic number[2] $\chi_m(\mathbb{R}^d)$ of the Euclidean space by the simple inequality $\chi_m(\mathbb{R}^d) \geq 1/m_1(\mathbb{R}^d)$. Indeed, if no colour class contains pairs of points at unit distance, then each of them has upper density at most $m_1(\mathbb{R}^d)$, and it takes at least $1/m_1(\mathbb{R}^d)$ such classes to cover the whole space. The parameter $m_1(\mathbb{R}^d)$ is many times studied in the context of providing lower bounds for the measurable chromatic number.

[1] Note that this problem is dilation invariant, so there is no loss of generality in assuming the forbidden distance to be 1.

[2] The measurable chromatic number of $\mathbb{R}^d$ is the minimum number of measurable sets needed to partition $\mathbb{R}^d$ so that no two points belonging to the same part are at distance 1 from each other.