
2015–
2020


2018–
2020

QuSoft is named on page 33 of the paper.

arXiv:2107.02567v2 [quant-ph] 19 May 2022

On a tracial version of Haemers bound

Li Gao* Sander Gribling† Yinan Li‡

May 20, 2022

Abstract

We extend upper bounds on the quantum independence number and the quantum Shannon capacity of graphs to their counterparts in the commuting operator model. We introduce a von Neumann algebraic generalization of the fractional Haemers bound (over $\mathbb{C}$) and prove that the generalization upper bounds the commuting quantum independence number. We call our bound the tracial Haemers bound, and we prove that it is multiplicative with respect to the strong product. In particular, this makes it an upper bound on the Shannon capacity. The tracial Haemers bound is incomparable with the Lovász theta function, another well-known upper bound on the Shannon capacity. We show that separating the tracial and fractional Haemers bounds would refute Connes' embedding conjecture.

Along the way, we prove that the tracial rank and tracial Haemers bound are elements of the (commuting quantum) asymptotic spectrum of graphs (Zuiddam, Combinatorica, 2019). We also show that the inertia bound (an upper bound on the quantum independence number) upper bounds the commuting quantum independence number.

1 Introduction

1.1 Independence number from nonlocal games

Two players, Alice and Bob, attempt to convince a referee that a given graph G contains an independent set of size d. The referee sends two integers $i_A, i_B \in [d]$ to Alice and Bob, respectively, who then response vertices g_A and g_B satisfying that

- if $i_A = i_B$ then $g_A = g_B$;
- if $i_A \neq i_B$ then $\{g_A, g_B\} \in E(\overline{G})$.

Alice and Bob can decide a strategy beforehand, but they cannot communicate during the game. Such a nonlocal game is known as the independent set game [MR16, MRV16]. A classical strategy corresponds to the case that Alice and Bob only use shared (classical) randomness. A perfect classical strategy enables Alice and Bob to win the independent set game for graph G with certainty. It can be shown that a perfect classical strategy exists if and only if the graph G has an independent set of size d. Thus, the largest possible d for a graph G where a perfect classical strategy for the independent set game exists is exactly the independence number $\alpha(G)$ of G.

One may also consider quantum strategies, where Alice and Bob are allowed to share quantum entanglement and perform measurements on their own systems. A perfect quantum strategy enables

*Department of Mathematics, University of Houston, Houston, TX, 77204, USA. lgao12@uh.edu
†IRIF, Université de Paris, CNRS, Paris, France. gribling@irif.fr
‡Graduate school of mathematics, Nagoya University, Nagoya, Japan. Yinan.Li@maths.nagoya-u.ac.jp