



QuSoft is named on page 27 of the paper.

Submitted to the Annals of Statistics

arXiv:2110.07583v3 [math.ST] 23 Oct 2025

NEAR OPTIMAL SAMPLE COMPLEXITY FOR MATRIX AND TENSOR NORMAL MODELS VIA GEODESIC CONVEXITY

BY COLE FRANKS[1,a], RAFAEL OLIVEIRA[2,b], AKSHAY RAMACHANDRAN[3,c] AND MICHAEL WALTER[4,d]

[1]*Department of Mathematics, Massachusetts Institute of Technology,* [a]*franks@mit.edu*

[2]*Cheriton School of Computer Science, University of Waterloo,* [b]*rafael@uwaterloo.ca*

[3]*Computer Science Department, University of British Columbia,* [c]*aramach@cs.ubc.ca*

[4]*Ludwig-Maximilians-Universität München,* [d]*michael.walter@lmu.de*

The matrix normal model, i.e., the family of Gaussian matrix-variate distributions whose covariance matrices are the Kronecker product of two lower dimensional factors, is frequently used to model matrix-variate data. The tensor normal model generalizes this family to Kronecker products of three or more factors. We study the estimation of the Kronecker factors of the covariance matrix in the matrix and tensor normal models.

For the above models, we show that the maximum likelihood estimator (MLE) achieves *nearly optimal nonasymptotic sample complexity* and *nearly tight error rates* in the Fisher-Rao and Thompson metrics. In contrast to prior work, our results do not rely on the factors being well-conditioned or sparse, nor do we need to assume an accurate enough initial guess. For the matrix normal model, all our bounds are minimax optimal up to logarithmic factors, and for the tensor normal model our bounds for the largest factor and for overall covariance matrix are minimax optimal up to constant factors provided there are enough samples for any estimator to obtain constant Frobenius error. In the same regimes as our sample complexity bounds, we show that the flip-flop algorithm, a practical and widely used iterative procedure to compute the MLE, converges linearly with high probability.

Our main technical insight is that, given enough samples, the negative log-likelihood function is *strongly geodesically convex* in the geometry on positive-definite matrices induced by the Fisher information metric. This strong convexity is determined by the expansion of certain random quantum channels.

CONTENTS

1 Introduction 2
1.1 Formal definitions and our results 6
1.2 Results on sample complexity & error bounds 8
1.3 Results on the flip-flop algorithm for MLE estimation 10
2 Geodesic convexity, sample complexity & error bounds 12
2.1 Geodesic convexity 12
2.2 Proof outline 15
2.3 Bounding the gradient 16
2.4 Strong convexity from expansion 17
2.5 Tensor normal model: sample complexity & error bounds 19
3 Matrix normal model: improved sample complexity & error bounds 20

MSC2020 subject classifications: Primary 62F12; secondary 62F30.

Keywords and phrases: Covariance estimation, matrix normal model, tensor normal model, maximum likelihood estimation, geodesic convexity, operator scaling, quantum expansion.