


2022–
2023



QuSoft is named on page 17 of the paper.

arXiv:2304.03234v3 [math.CO] 5 Nov 2024

ON THE THRESHOLD FOR SZEMERÉDI'S THEOREM WITH RANDOM DIFFERENCES

JOP BRIËT AND DAVI CASTRO-SILVA

Abstract. Using recent developments on the theory of locally decodable codes, we prove that the critical size for Szemerédi's theorem with random differences is bounded from above by $N^{1-\frac{2}{k}+o(1)}$ for length-k progressions. This improves the previous best bounds of $N^{1-\frac{1}{\lceil k/2\rceil}+o(1)}$ for all odd k.

1. Introduction

Szemerédi [20] proved that dense sets of integers contain arbitrarily long arithmetic progressions, a result which has become a hallmark of additive combinatorics. Multiple proofs of this result were found over the years, using ideas from combinatorics, ergodic theory and Fourier analysis over finite abelian groups.

Furstenberg's ergodic theoretic proof [14] opened the floodgates to a series of powerful generalizations. In particular, it led to versions of Szemerédi's theorem where the common differences for the arithmetic progressions are restricted to very sparse sets. We say that a set $D \subseteq [N]$ is *t-intersective* if any positive-density set $A \subseteq [N]$ contains a $(t+1)$-term arithmetic progression with common difference in D. Szemerédi's theorem implies that, for large enough N_0, the set $\{0, 1, \ldots, N_0\}$ is t-intersective for all $N \geq N_0$. Non-trivial examples include a result of Bergelson and Leibman [3] showing that the perfect squares (and more generally, images of integer polynomials with zero constant term) are t-intersective for every t, and a special case of a result of Wooley and Ziegler [23] showing the same for the prime numbers minus one.

The existence of such sparse intersective sets motivated the problem of showing whether, in fact, random sparse sets are typically intersective. The task of making this quantitative falls within the scope of research on threshold phenomena. We say that a property of subsets of $[N]$, given by a family $\mathcal{F} \subseteq 2^{[N]}$, is *monotone* if $A \in \mathcal{F}$ and $A \subseteq B \subseteq [N]$ imply $B \in \mathcal{F}$. The *critical size* $m^* = m^*(N)$ of a property is the least m such that a uniformly random m-element subset of $[N]$ has the property with probability at least $1/2$. (This value exists if $\mathcal{F}$ is non-empty and monotone, as this probability then increases

This work was supported by the Dutch Research Council (NWO) as part of the NETWORKS programme (grant no. 024.002.003). An extended abstract of this work appeared in the proceedings of EUROCOMB'23 [7].