

Entanglement Wedge Reconstruction via Universal Recovery Channels

Jordan Cotler,[1] Patrick Hayden,[1] Geoffrey Penington,[1] Grant Salton,[1] Brian Swingle,[1,2,3] and Michael Walter[1,4]

[1]*Stanford Institute for Theoretical Physics, Stanford University, Stanford CA 94305, USA*
[2]*Department of Physics, Harvard University, Cambridge MA 02138, USA*
[3]*Martin Fisher School of Physics, Brandeis University, Waltham MA 02453, USA*
[4]*Korteweg-de Vries Institute for Mathematics, Institute for Theoretical Physics, Institute for Logic, Language and Computation & QuSoft, University of Amsterdam, The Netherlands*

We apply and extend the theory of universal recovery channels from quantum information theory to address the problem of entanglement wedge reconstruction in AdS/CFT. It has recently been proposed that any low-energy local bulk operators in a CFT boundary region's entanglement wedge can be reconstructed on that boundary region itself. Existing work arguing for this proposal relies on algebraic consequences of the exact equivalence between bulk and boundary relative entropies, namely the theory of operator algebra quantum error correction. However, bulk and boundary relative entropies are only approximately equal in bulk effective field theory, and in similar situations it is known that predictions from exact entropic equalities can be qualitatively incorrect. The framework of universal recovery channels provides a robust demonstration of the entanglement wedge reconstruction conjecture in addition to new physical insights. Most notably, we find that a bulk operator acting in a given boundary region's entanglement wedge can be expressed as the response of the boundary region's modular Hamiltonian to a perturbation of the bulk state in the direction of the bulk operator. This formula can be interpreted as a noncommutative version of Bayes' rule that attempts to undo the noise induced by restricting to only a portion of the boundary, and has an integral representation in terms of modular flows. To reach these conclusions, we extend the theory of universal recovery channels to finite-dimensional operator algebras and demonstrate that recovery channels approximately preserve the multiplicative structure of the operator algebra.

arXiv:1704.05839v4 [hep-th] 4 Sep 2018

The AdS/CFT correspondence is a duality between a gravitational theory in $d+1$-dimensional asymptotically AdS space and a conformal field theory in one fewer spatial dimensions [1–5]. The CFT lives on the boundary of the bulk AdS space, and the quantum state of the boundary CFT is dual to the state of the quantum gravity theory in the bulk. Certain CFT states correspond to classical geometries in the bulk, and this important class of states occupies much of our attention. For such states, there is truly an emergent spatial direction in the bulk theory and understanding how locality with respect to the bulk geometry arises in the boundary theory is a longstanding problem. One way to approach the problem is to ask which regions of the bulk are completely described by a given region of the boundary.

The question above has been phrased in various forms over the years [6, 7], but a natural version is to identify all local bulk operators that can be expressed in terms of boundary operators with support only on the boundary subregion. Significant progress has been made on that problem, starting with the so-called HKLL prescription [2]. This method works in certain cases, but falls short in general; there are bulk operators that should be expressible on the boundary subregion that are inaccessible to this technique. A natural example is the operator corresponding to the area of the minimal surface in the bulk that calculates the entropy of the boundary subregion, the so-called Ryu-Takayanagi surface [8, 9]. For a single interval in empty AdS, HKLL and related techniques provide the answer, but even for a disconnected subregion of the boundary with total size greater than half, the minimal surface in the bulk falls outside of the purview of HKLL. We will review these techniques in detail later.

The problem of finding the dual to a boundary subregion is at the heart of the subject of bulk reconstruction: given an operator in the bulk, can one find a representation of this operator acting on a subregion of the boundary? Recently it has been proposed that, for a given subregion A of the conformal boundary, any low-energy bulk operator acting on the *entanglement wedge* of A (defined to be the bulk domain of dependence of any achronal bulk surface bounded by A and its associated covariant Ryu-Takayanagi minimal surface) can be reconstructed using only information in A [6, 10, 11]. The entanglement wedge reconstruction conjecture was strengthened in [12] and established in tensor network toy models of holography [12–14]. Very recently, it was proved in [15, 16] under the condition that the bulk and boundary relative entropies be *exactly* equal.

At large but finite N, within the framework of bulk effective field theory, one only expects approximate equality of the bulk and boundary relative entropies. When similar situations were studied in the quantum information theory literature, it was found that algebraic consequences of exact entropic equalities often do not provide qualitatively correct predictions in the approximate case. An important and, in fact, closely related example is the exact saturation of strong subadditivity of the von Neumann entropy, which is known to imply that the underlying state is a quantum Markov chain [17]. The associated algebraic structure slightly generalizes operator algebra quantum error correction, precisely the structure relevant in [15]. Near-saturation of strong subadditivity, however, *fails* to imply proximity to a quantum Markov chain state for systems of large Hilbert space dimension [18, 19].

Indeed the most direct generalization of the reconstruction theorem from [15] to approximate relative entropy equalities does not lead to full approximate quantum error correction (which is what we establish in this paper), but rather to a significantly weaker condition known as zerobits [20, 21]. Ignoring