

Stimulated Raman adiabatic passage-like protocols for amplitude transfer generalize to many bipartite graphs

Koen Groenland
QuSoft and CWI, Science Park 123, 1098 XG Amsterdam, the Netherlands
& Institute of Physics, University of Amsterdam,
*Science Park 904, 1098 XH Amsterdam, the Netherlands**

Carla Groenland
Mathematical Institute, University of Oxford,
Andrew Wiles Building, Radcliffe Observatory Quarter (550),
Woodstock Road, Oxford OX2 6GG, United Kingdom

Reinier Kramer
Max-Planck-Institut für Mathematik, Vivatsgasse 7, 53111 Bonn, Germany

(Dated: 17 June, 2020)

Adiabatic passage techniques, used to drive a system from one quantum state into another, find widespread application in physics and chemistry. We focus on techniques to spatially transport a quantum amplitude over a strongly coupled system, such as STImulated Raman Adiabatic Passage (STIRAP) and Coherent Tunnelling by Adiabatic Passage (CTAP). Previous results were shown to work on certain graphs, such as linear chains, square and triangular lattices, and branched chains. We prove that similar protocols work much more generally, in a large class of (semi-)bipartite graphs. In particular, under random couplings, adiabatic transfer is possible on graphs that admit a perfect matching both when the sender is removed and when the receiver is removed. Many of the favorable stability properties of STIRAP/CTAP are inherited, and our results readily apply to transfer between multiple potential senders and receivers. We numerically test transfer between the leaves of a tree, and find surprisingly accurate transfer, especially when straddling is used. Our results may find applications in short-distance communication between multiple quantum computers, and open up a new question in graph theory about the spectral gap around the value 0.

arXiv:1904.09915v2 [quant-ph] 17 Jun 2020

* koen.groenland@gmail.com