


Disorder enhanced quantum many-body scars in Hilbert crystals

Bart van Voorden,[1] Matteo Marcuzzi,[2] Kareljan Schoutens,[1,3] and Jiří Minář[1,3]

[1] *Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*

[2] *School of Physics and Astronomy, University of Nottingham, Nottingham, NG7 2RD, United Kingdom*

[3] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*

(Dated: December 11, 2020)

We consider a model arising in facilitated Rydberg chains with positional disorder which features a Hilbert space with the topology of a d-dimensional hypercube. This allows for a straightforward interpretation of the many-body dynamics in terms of a single particle one on the Hilbert space and provides an explicit link between the many-body and single particle scars. Exploiting this perspective, we show that an integrability-breaking disorder enhances the scars followed by inhibition of the dynamics due to strong localization of the eigenstates in the large disorder limit. Next, mapping the model to the spin-1/2 XX Heisenberg chain offers a simple geometrical perspective on the recently proposed Onsager scars [PRL **124**, 180604 (2020)], which can be identified with the scars on the edge of the Hilbert space. This makes apparent the origin of their insensitivity to certain types of disorder perturbations.

arXiv:2012.05310v1 [cond-mat.stat-mech] 9 Dec 2020

Introduction. The understanding of thermalization and relaxation dynamics is at the forefront of research on quantum many-body systems out-of-equilibrium. Since the formulation of the eigenstate thermalization hypothesis[1–3], predicting fast thermalization following a quench from most many-body states, many exceptions to this behaviour have been identified. The prominent examples are integrable[4,5] and many-body localized (MBL) systems[6–14]. A recently added category are quantum many-body scars (QMBS)[15,16], which are particular eigenstates responsible for slow decay and oscillatory behaviour of observables following a quantum quench from certain initial states, typically close to a product state, as observed in Ref.[17] realizing the so-called PXP model[18]. This has triggered a great interest in QMBS in settings ranging from constrained to driven[19–64] and recently also disordered systems[65].

QMBS owe their name to the single-particle quantum scars[66,67] which were in turn inspired by particle motion in classical billiards. In both the quantum and classical cases, it is the shape of the billiard boundary, such as the celebrated Bunimovich stadium or cardioid shape[68,69], which causes the motion of the particle to be generically ergodic. The exception to this rule is a set of periodic trajectories, around which the density of certain wavefunctions - the scars - is enhanced in the quantum case.

Here we analyze a model of spins-1/2, which describes a chain of Rydberg atoms with open boundaries under a facilitation condition[70]. Representing the Hilbert space as a graph, we show that it corresponds to a truncated hypercube with the dimension given by the number of spin clusters (cf. below for definition).

This allows us to identify the QMBS as single particle scars on the Hilbert space. Building on the graph representation of the Hilbert space, an approach also exploited in the studies of MBL[6,7,31,71–76], we demonstrate that the scar signatures are *enhanced* in the presence of disorder, naturally emerging from the positional disorder of the atoms. Finally, exploiting the mapping of the present model to the Heisenberg spin-1/2 XX chain[77], we identify the recently proposed Onsager scars[78,79] with scars corresponding to sparse eigenstates residing at the "edge" of the Hilbert space. This provides intriguing connections between QMBS and single-particle scars and highlights the utility of a graph-theoretical approach to many-body dynamics, which has been advocated also in the studies of quantum chaos[80–84], integrability[85], QMBS[86] and fermionic and exchange models[87,88].

The model. We consider a one dimensional chain of M Rydberg atoms along the z-axis, with open boundaries and spaced by r_0. We denote the ground and excited (Rydberg) state as $|\downarrow\rangle$, $|\uparrow\rangle$. The corresponding Hamiltonian reads

$$H_{\rm Ry} = \sum_k \frac{\Omega}{2}\sigma_k^x + \Delta n_k + \sum_{l>k} V(|\mathbf{r}_k - \mathbf{r}_l|)\, n_k n_l, \qquad (1)$$

where $\sigma_k^x = |\uparrow_k\rangle\langle\downarrow_k| + |\downarrow_k\rangle\langle\uparrow_k|$, $n_k = |\uparrow_k\rangle\langle\uparrow_k|$, and $V(r) = C_\alpha/r^\alpha$, $r = |\mathbf{r}|$. C_α, which we take to be positive, is the interaction strength coefficient with $\alpha = 3\,(6)$ for dipole-dipole (Van der Waals) interaction. The positions of the atoms are $\mathbf{r}_k = (0, 0, (k-1)r_0) + \delta\mathbf{r}_k$, where $\delta\mathbf{r}_k$ describes the disorder which induces the disorder in energy. Denoting $V_{\rm NN} = V(r_0)$ and $V_{\rm NNN} = V(2r_0)$, we define an energy shift for a pair of nearest neighbours $\delta V_k = V_{\rm NN} - V(|\mathbf{r}_{k+1} - \mathbf{r}_k|)$.

It has been shown in[70] that under the facilitation condition $\Delta = -V_{\rm NN}$ and in the regime $V_{\rm NN} \gg \Omega, \delta V_k$ the Hamiltonian (1) effectively reduces to

$$H_{\rm eff} = \Delta N_{\rm cl} + \sum_k \frac{\Omega}{2}\sigma_k^x P_{\langle k\rangle} + \delta V_k n_k n_{k+1} + V_{\rm NNN} n_k n_{k+2}, \qquad (2)$$

where $P_{\langle k\rangle} = n_{k-1} + n_{k+1} - 2n_{k-1}n_{k+1}$ and $N_{\rm cl} = \sum_k n_k(1-n_{k+1})$, $n_0 = n_{N+1} = 0$, denotes the number of clusters, which are blocks of consecutive spin excitations (e.g. the configuration $\downarrow\downarrow\boxed{\uparrow\uparrow}\downarrow\boxed{\uparrow\uparrow\uparrow}$ contains two clusters highlighted by boxes). The projector $P_{\langle k\rangle}$ ensures