

2021–
2022

arXiv:2207.12960v2 [quant-ph] 17 Mar 2023

Projective measurements can probe non-classical work extraction and time-correlations

Santiago Hernández-Gómez,[1,2,3,*] Stefano Gherardini,[4,1,†] Alessio Belenchia,[5,6] Matteo Lostaglio,[7,‡] Amikam Levy,[8] and Nicole Fabbri[1,3]

[1] *European Laboratory for Non-linear Spectroscopy (LENS), Università di Firenze, I-50019 Sesto Fiorentino, Italy*
[2] *Dipartimento di Fisica e Astronomia, Università di Firenze, I-50019, Sesto Fiorentino, Italy*
[3] *Istituto Nazionale di Ottica del Consiglio Nazionale delle Ricerche (CNR-INO), I-50019 Sesto Fiorentino, Italy*
[4] *Istituto Nazionale di Ottica del Consiglio Nazionale delle Ricerche (CNR-INO), Science Park, Basovizza, I-34149 Trieste, Italy*
[5] *Institut für Theoretische Physik, Eberhard-Karls-Universität Tübingen, 72076 Tübingen, Germany*
[6] *Centre for Theoretical Atomic, Molecular and Optical Physics, School of Mathematics and Physics, Queen's University Belfast, Belfast BT7 1NN, United Kingdom*
[7] *Korteweg-de Vries Institute for Mathematics and QuSoft, University of Amsterdam, The Netherlands*
[8] *Department of Chemistry and Center for Quantum Entanglement Science and Technology, Bar-Ilan University, Ramat-Gan 52900, Israel*

We demonstrate an experimental technique to characterize genuinely nonclassical multi-time correlations using projective measurements with no ancillas. We implement the scheme in a nitrogen-vacancy center in diamond undergoing a unitary quantum work protocol. We reconstruct quantum-mechanical time correlations encoded in the Margenau-Hills quasiprobabilities. We observe work extraction peaks five times those of sequential projective energy measurement schemes and in violation of newly-derived stochastic bounds. We interpret the phenomenon via anomalous energy exchanges due to the underlying negativity of the quasiprobability distribution.

There is no unique way of defining a joint probability for the multi-time statistics of quantum mechanical quantities since even a single quantum observable does not always commute with itself at different times. Nevertheless, across the quantum sciences, multi-time fluctuations of quantum-mechanical quantities, especially energy, play a crucial role. Correlation functions between events at different times resemble joint probability distributions for the eigenvalues of the observables, but they are in general neither real nor positive. In fact, they are represented by *quasiprobabilities*, akin to the well-known Wigner function in quantum optics [1], but associated with a process rather than a state. The standard definition of a temporal correlation function between two events described by projectors $\Pi_i(0)$ and $\Pi_f(t)$,

$$q_{if}^{\mathrm{KD}} = \mathrm{Tr}\left[\rho\Pi_i(0)\Pi_f(t)\right], \tag{1}$$

coincides with a Kirkwood-Dirac quasiprobability (KDQ) [2–4], and the same goes for multi-time extensions.

As we survey in [5], the centrality of the KDQ and its real part, the Margenau-Hill quasiprobability (MHQ), has only recently come to be fully appreciated. These quasiprobabilities underpin analysis from perturbation theory [5] to information scrambling [6, 7]. Weak values [5, 8, 9] are conditional KDQ [5, 7] and generalized *anomalous weak values* are in one-to-one correspondence with non-classical (negative or complex) KDQ quasiprobabilities [10, 11]. Negative values of the MHQ are linked to quantum metrological advantages in both local and postselected setups [12–14] and to power output advantages in quantum thermodynamics [14].

Contrary to investigations of multi-time processes via sharp measurements [15–18], experimental investigations with quasiprobabilities have mostly focused on characterizing states rather than processes. This has a long tradition in quantum optics, where phase-space quasiprobabilities have been used extensively for tomographic scopes [19–22]. To our knowledge, experimental access to the KDQ and MHQ has been limited to weak measurement schemes and, even then, mostly to characterize states [23–27].

In such a context, the aim of our work is two-fold: First, pave the way to the experimental study of the role of non-commutativity in temporal correlations via the MHQ. We do so by providing the first experimental demonstration of a *weak two-point measurement* (wTPM) scheme [5, 28], a protocol that – in contrast to weak measurement schemes – requires neither ancillae nor fine-tuned system-ancilla couplings. In fact, we reconstruct the back-reaction-free limit encapsulated by quasiprobabilities not by a weak measurement, but by linearly combining different projective measurement schemes in such a way that different back-reactions cancel. Conceptually our idea can be seen as a twist on probabilistic error cancellation techniques in quantum computing, where several noisy circuits are sampled from to reconstruct an ideal error-free limit [29]. This technique can be deployed in quantum thermodynamics beyond the scope of our study.

Second, we lay down the theoretical and experimental ground for the study of non-classical energetic processes via the MHQ. We measure work extraction peaks in a driven three-level system up to five times those of the