

2022–
2024

Classical shadow tomography for continuous variables quantum systems

Simon Becker, Nilanjana Datta, Ludovico Lami, and Cambyse Rouzé

arXiv:2211.07578v2 [quant-ph] 1 Mar 2023

***Abstract*—In this article we develop a continuous variable (CV) shadow tomography scheme with wide ranging applications in quantum optics. Our work is motivated by the increasing experimental and technological relevance of CV systems in quantum information, quantum communication, quantum sensing, quantum simulations, quantum computing and error correction. We introduce two experimentally realisable schemes for obtaining classical shadows of CV (possibly non-Gaussian) quantum states using only randomised Gaussian unitaries and easily implementable Gaussian measurements such as homodyne and heterodyne detection. For both schemes, we show that $N = O(\mathrm{poly}(\frac{1}{\epsilon}, \log(\frac{1}{\delta}), M_n^{r+\alpha}, \log(m)))$ samples of an unknown m-mode state ρ suffice to learn the expected value of any r-local polynomial in the canonical observables of degree α, both with high probability $1-\delta$ and accuracy ϵ, as long as the state ρ has moments of order $n > \alpha$ bounded by M_n. By simultaneously truncating states and operators in energy and phase space, we are able to overcome new mathematical challenges that arise due to the infinite-dimensionality of CV systems. We also provide a scheme to learn nonlinear functionals of the state, such as entropies over any small number of modes, by leveraging recent energy-constrained entropic continuity bounds. Finally, we provide numerical evidence of the efficiency of our protocols in the case of CV states of relevance in quantum information theory, including ground states of quadratic Hamiltonians of many-body systems and cat qubit states. We expect our scheme to provide good recovery in learning relevant states of 2D materials and photonic crystals.**

I. Introduction

Obtaining classical descriptions of states of quantum-mechanical systems is a fundamental ingredient of quantum computing. It is useful for storing and transmitting quantum information and essential for verification and benchmarking of quantum devices. However, the underlying quantum nature of such systems provides a huge hurdle in obtaining such a classical description: to learn anything about a quantum state one needs to measure it, but measurements in quantum mechanics are inherently destructive and furthermore probabilistic, entailing that individual measurement outcomes only give limited information about the state of the system. Consequently, in order to obtain a classical representation of a quantum state, one requires multiple identical copies of the state on which successive, appropriate (possibly adaptive) single-copy measurements can be performed.

This is what is done in the traditional method of learning an unknown quantum state, known as *quantum state tomography*. It is the process of inferring a quantum state by using suitable measurements on many identical copies of the state. There is, however, a huge practical limitation in using quantum state tomography for many-body quantum systems. This is due to the so-called "curse of dimensionality": the number of parameters needed to fully specify the state of a quantum-mechanical system grows exponentially with the system size. The exponential number of measurements needed to infer these parameters makes quantum state tomography infeasible for large systems. Consequently, full quantum state tomography has only been realised in systems with few components, in particular, in a system of ten qubits, which too required millions of measurements. In a nutshell, obtaining a classical description of a d-dimensional quantum mixed state ρ, given many copies of it, via quantum state tomography, can be shown to require $\Omega(d^2)$ copies of the state. (More recently, it has been shown [1], [2] that $O(d^2)$ copies also suffice.) However, this number grows exponentially with the number n of qubits (since $d = 2^n$), and the problem becomes rapidly intractable.

In 2018 Aaronson [3] pointed out that for certain concrete tasks, obtaining a complete classical characterisation of the quantum state is unnecessary. Instead it is often sufficient to accurately predict many useful properties of the state. This led him to propose a novel task called *shadow tomography*, the aim of which is not to learn a complete description of the unknown quantum state but instead to simultaneously estimate the outcome probabilities associated with a list of M two-outcome measurements, $E_1, \ldots, E_M$, performed on the state, up to a desired accuracy (say, ε). For an unknown d-dimensional quantum mixed state ρ, this requires the prediction of M expectation values, $\mathrm{Tr}[E_i\rho]$ $i = 1, 2, \ldots, M$, to within an additive error ε. Aaronson showed that, remarkably, the number of copies of the

Simon Becker is with the Department of Mathematics, ETH Zürich, Rämistrasse 101, 8092 Zürich, Switzerland. Email: simon.becker@math.ethz.ch

Nilanjana Datta is with the Department of Applied Mathematics and Theoretical Physics, Centre for Mathematical Sciences, University of Cambridge, Cambridge CB3 0WA, United Kingdom. Email: n.datta@damtp.cam.ac.uk

Ludovico Lami is with the Korteweg–de Vries Institute for Mathematics, the Institute for Theoretical Physics, and QuSoft, University of Amsterdam, Science Park 123, 1098 XG Amsterdam. Part of this work was conducted while he was at the Institute for Theoretical Physics, University of Ulm, Albert-Einstein-Allee 11, D-89069 Ulm, Germany. Email: ludovico.lami@gmail.com

Cambyse Rouzé is with the Zentrum Mathematik, Technische Universität München, 85748 Garching, Germany. Email: rouzecambyse@gmail.com