

2022–
2024

arXiv:2302.03075v3 [quant-ph] 21 May 2024

# Testing the quantum nature of gravity without entanglement

Ludovico Lami,[1,2,3,4,*] Julen S. Pedernales,[4,†] and Martin B. Plenio[4,‡]

[1] *QuSoft, Science Park 123, 1098 XG Amsterdam, The Netherlands*
[2] *Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, The Netherlands*
[3] *Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*
[4] *Institut für Theoretische Physik und IQST, Universität Ulm, Albert-Einstein-Allee 11, D-89069 Ulm, Germany*

Given a unitary evolution $U$ on a multi-partite quantum system and an ensemble of initial states, how well can $U$ be simulated by local operations and classical communication (LOCC) on that ensemble? We answer this question by establishing a general, efficiently computable upper bound on the maximal LOCC simulation fidelity — what we call an 'LOCC inequality'. We then apply our findings to the fundamental setting where $U$ implements a quantum Newtonian Hamiltonian over a gravitationally interacting system. Violation of our LOCC inequality can rule out the LOCCness of the underlying evolution, thereby establishing the non-classicality of the gravitational dynamics, which can no longer be explained by a local classical field. As a prominent application of this scheme we study systems of quantum harmonic oscillators initialised in coherent states following a normal distribution and interacting via Newtonian gravity, and discuss a possible physical implementation with torsion pendula. One of our main technical contributions is the analytical calculation of the above LOCC inequality for this family of systems. As opposed to existing tests based on the detection of gravitationally mediated entanglement, our proposal works with coherent states alone, and thus it does not require the generation of largely delocalised states of motion nor the detection of entanglement, which is never created at any point in the process.

## I. INTRODUCTION

The weakest force of Nature, gravity, is also the cruellest. The desire to understand its elusive character has often stirred the very roots of science, breeding so many of the greatest achievements of human thought; at the same time, the extreme weakness of the gravitational interaction, compared to all of the other forces, has placed us in the bitter situation of not, yet, being able to investigate its ultimate nature. The question of whether gravity is fundamentally a classical or a quantum field, or perhaps an entity of an altogether different type, remains so far unanswered.

While probing phenomena where the quantum degrees of freedom of gravity could directly be tested remains prohibitive with current or near-term technologies, determining whether spacetime could exist in a superposition of distinct classical configurations is a substantially easier, if still challenging, endeavour. A first proposal in that direction was put forth by Feynman [1], who presented a Gedankenexperiment in which a mass capable of generating a gravitational field is placed in a superposition of different spatial locations; such a mass could induce — via the gravitational interaction — a superposition in a second test mass, and the physical existence of such quantum coherences (mediated by gravity) can be tested, at least in principle.

In quantum as well as post-quantum theories, correlated superpositions naturally lead to the notion of entanglement [2]. Therefore, further developing the above idea leads to the natural prediction that no classical gravitational field can entangle two quantum mechanical systems [3–5]. Recently, proposals seeking to detect gravitationally generated entanglement [3, 6–15] have prompted a discussion of the conclusions that could be drawn from these experiments regarding the quantum nature of gravity [16–24]. The concept underlying these proposals is the creation of coherently delocalised quantum states of two masses, in such a way that different branches of the wavefunction of one mass have different distances to those of the other mass. As a result of the gravitational interaction, evolution in time will assign diverging phases to those components, effectively entangling the two systems. Such entanglement can then be detected experimentally in various ways. For an introduction to these topics, we refer the reader to the recent review [25].

It is worth noting that besides the impossibility of entanglement generation, a classical gravity would be subjected to several other constraints that may also have observable consequences. For example, the continuous measurement to which a quantum source should be subjected by hand of a classical gravity should produce a decoherence effect [3, 26–31]. Other proposed experiments seek to exploit the spatial superposition of source masses [6, 7, 32–34], or aim to detect the possible non-Gaussian signatures of a genuine quantum theory of gravity [35].

For the sake of this discussion, let us focus our attention on the gravitational entanglement proposals. In lay-

* ludovico.lami@gmail.com
† julen.pedernales@uni-ulm.de
‡ martin.plenio@uni-ulm.de