2022–
2023

arXiv:2305.03012v1 [math.CO] 4 May 2023

Quasirandom additive sets and Cayley hypergraphs

Davi Castro-Silva

CWI & QuSoft

May 5, 2023

Abstract

We study the interplay between notions of quasirandomness for additive sets and for hypergraphs. In particular, we show a strong connection between the notions of Gowers uniformity in the additive setting and discrepancy-type measures of quasirandomness in the hypergraph setting. Exploiting this connection, we provide a long list of disparate quasirandom properties regarding both additive sets and Cayley-type hypergraphs constructed from such sets, and show that these properties are all equivalent (in the sense of Chung, Graham and Wilson) with polynomial bounds on their interdependences.

1 Introduction

Quasirandom properties can be informally thought of as certificates of randomness for the object in consideration. Given some class of combinatorial objects, such as graphs or 3-uniform hypergraphs, we say that a (deterministic) property of these objects is a *quasirandom property* if it satisfies two conditions: a uniformly random object from that class satisfies this property with high probability; and any object which satisfies this property will also behave in many other ways like a random object. In such cases, just by knowing that an object satisfies some quasirandom property, one gleans a wealth of information about its behaviour in many respects; such objects are then said to be *quasirandom*.

The notion of quasirandomness was originally introduced in the setting of graphs, in a seminal paper of Chung, Graham and Wilson [10]. These authors considered several natural properties typically satisfied by random graphs – such as having the expected subgraph counts, having uniform edge-distribution over vertex cuts or having large spectral gap – and showed that all of them are (in a specific sense) equivalent to each other. Following their work, the study of quasirandom properties has been extended to several other combinatorial classes of objects; we refer the reader to Chung's website [4] for a long list of references. In this paper we will consider hypergraphs and additive sets (i.e. subsets of additive groups), focusing on the relationships between their respective quasirandom properties.