


arXiv:2311.02941v2 [physics.atom-ph] 23 Feb 2024

Optical pumping of $5s4d\,^1D_2$ strontium atoms for laser cooling and imaging

Jens Samland [1,2], Shayne Bennetts [1], Chun-Chia Chen (陳俊嘉) [1], Rodrigo González Escudero [1], Florian Schreck [1,3] and Benjamin Pasquiou [1,3,4,5,*]

[1] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, Science Park 904, 1098XH Amsterdam, The Netherlands*
[2] *Physik-Department, Technical University of Munich, James-Franck-Str.1, 85748 Garching, Germany*
[3] *QuSoft, Science Park 123, 1098XG Amsterdam, The Netherlands*
[4] *CNRS, UMR 7538, LPL, F-93430, Villetaneuse, France*
[5] *Laboratoire de Physique des Lasers, Université Sorbonne Paris Nord, F-93430, Villetaneuse, France*

(Dated: February 26, 2024)

We present a faster repumping scheme for strontium magneto-optical traps operating on the broad $5s^2\,^1S_0 - 5s5p\,^1P_1$ laser cooling transition. Contrary to existing repumping schemes, we directly address lost atoms that spontaneously decayed to the $5s4d\,^1D_2$ state, sending them back into the laser cooling cycle by optical pumping on the $5s4d\,^1D_2 - 5s8p\,^1P_1$ transition. We thus avoid the $\sim 100\,\mu s$-slow decay path from $5s4d\,^1D_2$ to the $5s5p\,^3P_{1,2}$ states that is part of other repumping schemes. Using one low-cost external-cavity diode laser emitting at 448 nm, we show our scheme increases the flux out of a 2D magneto-optical trap by 60 % compared to without repumping. Furthermore, we perform spectroscopy on the $5s4d\,^1D_2 - 5s8p\,^1P_1$ transition and measure its frequency $\nu_{^{88}\mathrm{Sr}} = (668917515.3 \pm 4.0 \pm 25)$ MHz. We also measure the frequency shifts between the four stable isotopes of strontium and infer the specific mass and field shift factors, $\delta\nu^{88,86}_{\mathrm{SMS}} = -267(45)$ MHz and $\delta\nu^{88,86}_{\mathrm{FS}} = 2(42)$ MHz. Finally, we measure the hyperfine splitting of the $5s8p\,^1P_1$ state in fermionic strontium, and deduce the magnetic dipole and electric quadrupole coupling coefficients $A = -4(5)$ MHz and $B = 5(35)$ MHz. Our experimental demonstration shows that this simple and very fast scheme could improve the laser cooling and imaging performance of cold strontium atom devices, such as quantum computers based on strontium atoms in arrays of optical tweezers.

I. INTRODUCTION

Laser cooling and trapping techniques such as magneto-optical traps (MOTs) play a pivotal role in cold atom and molecule experiments. As a basic component they enable a wide range of quantum technologies such as optical clocks [1, 2], atom interferometers [3–5], quantum simulators and quantum computers, e.g. using arrays of optical tweezers [6]. The success of these experiments relies on a combination of the right species with the right laser cooling technique. In recent years, laser-cooled alkaline-earth elements, and in particular strontium, have become widely used platforms for next-generation clocks, interferometers, and quantum computers [7–10]. Their success can be attributed to their atomic level structure featuring a broad optical transition perfect for fast laser cooling, and narrow transitions between singlet and triplet states. These provide clock transitions as well as laser-cooling to µK temperatures and even down to quantum degeneracy [11–13]. Continued development of new quantum technologies relies on the continued improvement of laser cooling techniques.

In the case of strontium, the broad 30 MHz-wide $5s^2\,^1S_0 - 5s5p\,^1P_1$ transition used for the first laser cooling stages is not fully closed. Each cycle through this transition has a probability of one out of 50000 [14–17] or one out of 20000 [9, 18, 19] for an atom in the excited $5s5p\,^1P_1$ state to decay into the $5s4d\,^1D_2$ state, leading to losses without an additional repumping scheme. After some time spent in $5s4d\,^1D_2$, the atom further decays into the $5s5p\,^3P_{1,2}$ states, from which one typically repumps it back into the $5s^2\,^1S_0 - 5s5p\,^1P_1$ laser cooling cycle. Unfortunately, the lifetime of the $5s4d\,^1D_2$ state is rather long, 300 µs [20–23]. Such a long time spent in an unaddressed state is potentially detrimental when a fast cycling transition is required in order not to lose the atom, e.g. for a Zeeman slower, a 2D MOT, or a fast and reliable single atom detection [24, 25].

Taking the example of fluorescence imaging single strontium atoms trapped in optical tweezers, each atom needs to scatter many photons in order to achieve a high readout fidelity [26]. However, an atom that decays into the $5s4d\,^1D_2$ state has enough time to escape the tweezers region, if the tweezers is not confining for $5s4d\,^1D_2$. Strikingly, several experiments with tweezers have chosen to use light at $515 - 532$ nm [9, 17], because of the available laser power, the small wavelength and since 515 nm allows to create traps with magic wavelength for the $5s^2\,^1S_0 - 5s5p\,^3P_1$ transition [9, 27]. Moreover, the polarizability at such short wavelength can be suitable for trapping strontium atoms excited to Rydberg states [28–30]. However, tweezers at these wavelengths create a repulsive potential for the $5s4d\,^1D_2$ state that expels the atoms. Even when the tweezers wavelength is chosen to be confining, the decay toward the $5s4d\,^1D_2$ state can remain an inconvenience as one needs to wait for the atoms to decay to the $5s5p\,^3P_{1,2}$ states. For example, fast single atom fluorescence detection, which is useful for quantum

* SrRepump448nm@strontiumBEC.com