

On compatibility of binary qubit measurements

Dmitry Grinko[1,2] and Roope Uola[3]

[1] *QuSoft, University of Amsterdam, Amsterdam, The Netherlands*
[2] *Department of Mathematics, University of Geneva, 1211 Geneva, Switzerland*
[3] *Department of Applied Physics, University of Geneva, 1211 Geneva, Switzerland*

(Dated: 11th July 2024)

Deciding which sets of quantum measurements allow a simultaneous readout is a central problem in quantum measurement theory. The problem is relevant not only from the foundational perspective, but also has direct applications in quantum correlation problems that are fueled by the presence of incompatible measurements. Although central, only few analytical criteria exist for deciding the incompatibility of general sets of measurements. In this work, we approach the problem through functions defined on the Boolean hypercube and their Fourier transformations. We show that this reformulation of the problem leads to a complete geometric characterisation of joint measurability of any finite set of unbiased binary qubit measurements, and gives a necessary condition for the biased case. We discuss our results in the realm of quantum steering, where they translate into a family of steering inequalities. When certain unbiasedness conditions are fulfilled, these criteria are tight, hence fully characterizing the steering problem, when the trusted party holds a qubit and the untrusted party performs any finite number of binary measurements. We further discuss how our results point towards a second-order cone programming approach to measurement incompatibility, and compare this to the predominantly used semi-definite-programming-based techniques. We use our approach to falsify an existing conjecture on measurement incompatibility of special sets of measurements.

arXiv:2407.07711v1 [quant-ph] 10 Jul 2024

INTRODUCTION

Quantum measurement incompatibility forms a cornerstone of quantum information theory. In addition to its importance in describing foundational aspects of quantum theory in, e.g., the Heisenberg microscope [1] and fundamental interferometric bounds [2], incompatibility has recently found various applications in quantum information theory [3]. To mention a few, incompatibility is known to fully characterize quantum steering [4–7], instances of Bell non-locality [8–11], as well as preparation contextuality [12–14], and it works as a prerequisite for the quantum advantage in various prepare-and-measure scenarios [15–24].

The fundamental interest on and the wide applicability of incompatibility have motivated a long line of research on characterizing and quantifying this non-classical phenomenon. This includes the development of semi-definite programming techniques [6, 8, 25–28], quantification techniques motivated by resource theoretic measures [6, 29–31], (semi-)device independent witnesses [32–35], geometric methods [36–46], Banach space techniques [47–49], and formulations of the concept in generalized probabilistic theories [50–52], as well as experimental quantification of the effect [53]. We note that all the mentioned works are in the finite-dimensional and discrete outcome setting, and refer to the work of Busch et al. for a collection of results in the continuous realm [54].

In this manuscript, we introduce an analytic method for characterizing incompatibility of yes-no questions, i.e. binary measurements, based on Fourier transformations of functions defined on the Boolean hypercube. This amounts to representing the existence problem of joint measurements through Fourier transformations of their Bloch representation, and constructing a solution for this dual problem. By utilizing the theory of Fourier analysis [55, 56], we derive a collection of necessary criteria for compatibility of any finite number of binary qubit measurements. We show that when the involved measurements are unbiased, i.e. projective measurements subject to white noise, the criteria give a full analytical characterization of incompatibility. This generalizes various known special instances of the geometric approach to compatibility and as such, we give a brief overview of the existing results on this research line. As direct applications, we give a characterization of the related quantum steering problems, discuss the possibility of using second-order cone programming for solving the problem of joint measurability, and falsify a conjecture on measurement incompatibility presented in [45].

JOINT MEASURABILITY

We model quantum measurements as positive operator valued measures (POVMs). These are collections of positive semi-definite Hermitian operators $\{J(\nu)\}_{\nu\in\Omega}$ acting on a finite-dimensional Hilbert space summing to the identity operator, i.e. $\sum_{\nu\in\Omega} J(\nu) = \mathbb{1}$, where Ω is the set of outcome labels. We denote a measurement assemblage by $\mathcal{M} = \{J_i(\mu_i)\}_{\mu_i,i}$, where $i \in \{1,\dots,N\}$ labels the choice of a measurement, i.e. $\{J_i(\mu_i)\}_{\mu_i\in\Omega_i}$ is a POVM for each i. For an assemblages of N POVMs measurement compatibility is defined through the concept of joint measurability [3, 54, 57]. An assemblage $\mathcal{M}$ is