

# Mutual information fluctuations and non-stabilizerness in random circuits

Arash Ahmadi,[1,*] Jonas Helsen,[2] Cagan Karaca,[1] and Eliska Greplova[1]

[1] *QuTech and Kavli Institute of Nanoscience, Delft University of Technology, Delft, the Netherlands*
[2] *QuSoft and CWI, Amsterdam, The Netherlands*

The emergence of quantum technologies has brought much attention to the characterization of quantum resources as well as the classical simulatability of quantum processes. Quantum resources, as quantified by non-stabilizeress, have in one theoretical approach been linked to a family of entropic, monotonic functions. In this work, we demonstrate both analytically and numerically a simple relationship between non-stabilizerness and information scrambling using the fluctuations of an entropy-based quantifier. Specifically, we find that the non-stabilizerness generated by a random quantum circuit is proportional to fluctuations of mutual information. Furthermore, we explore the role of non-stabilizerness in measurement-induced entanglement phase transitions. We find that the fluctuations of mutual information decrease with increasing non-stabilizerness yielding potentially easier identification of the transition point. Our work establishes a key connection between quantum resource theory, information scrambling and measurement-induced entanglement phase transitions.

arXiv:2408.03831v2 [quant-ph] 12 Aug 2024

## I. INTRODUCTION

Since the formulation of quantum theory, entanglement has been known to be one of the unique signatures of quantum theory [1]. While entanglement is one of the key features of distinguishing the quantum world from the classical, it is known that entanglement alone is insufficient for having an advantage from the *computational* point of view. In particular, the stabilizer states are a set of states that can be highly entangled and at the same time can be simulated efficiently on classical computers [2–7].

The missing feature for quantum computation to have an advantage over classical computation is a concept known as non-stabilizerness or *magic* [8]. In quantum circuit language, in the fault-tolerant regime, the gates that are known to be "cheap" gates [8] are Clifford gates, and magic is injected into the state by adding non-Clifford gates to the circuit. It is also known that the Clifford operations could have easier implementation both at the experimental level and for quantum error correction [2, 4, 9, 10]. This feature makes magic the resource for quantum computation.

Various measures for non-stabilizerness, or magic, have been proposed. Some of the notable ones are magical cross-entropy, mana [8], robustness of magic [11, 12], stabilizer entropies (SE) [13] and others [14–18]. Due to the favourable scaling of the stabilizer entropies these measures received much attention from the community to study the properties of non-stabilizerness of quantum systems [19–25]. Additionally, there has been a significant effort to scale up the computability of stabilizer entropies [26–30].

Diverse studies confirmed there exists a relationship between non-stabilizerness and information scrambling [13, 16, 31]. Information scrambling describes the spread of local information in a generic quantum system [32]. Information scrambling has been shown to be one of the most powerful measures for various quantum properties of quantum systems, from black holes [33–36] to many-body quantum systems [37] to quantum circuits [31, 38]. There are a number of known methods for measuring information scrambling. One of the well-known approaches is based on the measurement of correlator functions, namely Out-of-Time Ordered Correlators (OTOCs) [33, 36, 38–41]. Another approach for studying information scrambling are entropy-based measures [32, 42–48].

In this paper, we describe numerically and analytically a general relationship between non-stabilizerness and *fluctuations* of entropy-based measures of information scrambling. A specific instance of this relation has been observed earlier by connecting fluctuations of OTOCs to non-stabilizerness [13, 31]. In the present work, we generalize this behaviour to fluctuations of generic mutual information measures of disjoint regions of the quantum circuit on t-doped circuits. The observed link between non-stabilizerness and mutual information creates a bridge to the theory of phase transitions in measurement-induced random quantum circuits, where mutual information plays a crucial role [49–81]. Here we show that non-stabilizerness is an ingredient in reducing the spread of entanglement phase transition in random quantum circuits and also reduces the fluctuations of the measured mutual information.

## II. RANDOM CIRCUITS AND MUTUAL INFORMATION

### A. Definitions and notation

The structure of the T-doped circuits that we use in this study consists of blocks of random Clifford operations followed by single T-gates on random qubits in the circuit. The blocks of random Clifford consist of single Clifford gates drawn randomly from the set $\{I, X, Y, Z, H, S\}$ followed by three CNOT gates on two

* a.ahmadi-1@tudelft.nl