

2022–
2024

A solution of the generalised quantum Stein's lemma

Ludovico Lami[1,2,3,4,*]

[1]*Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa, Italy*

[2]*QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*

[3]*Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands*

[4]*Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*

arXiv:2408.06410v3 [quant-ph] 25 Jul 2025

We solve the generalised quantum Stein's lemma, proving that the Stein exponent associated with entanglement testing, namely, the quantum hypothesis testing task of distinguishing between n copies of an entangled state ρ_{AB} and a generic separable state $\sigma_{A^n:B^n}$, equals the regularised relative entropy of entanglement. Not only does this determine the ultimate performance of entanglement testing, but it also establishes the reversibility of all quantum resource theories under asymptotically resource non-generating operations, with the regularised relative entropy of resource governing the asymptotic transformation rate between any two quantum states. As a by-product, we prove that the same Stein exponent can also be achieved when the null hypothesis is only approximately i.i.d., in the sense that it can be modelled by an 'almost power state'. To solve the problem we introduce two techniques. The first is a procedure that we call 'blurring', which, informally, transforms a permutationally symmetric state by making it more evenly spread across nearby type classes. Blurring alone suffices to prove the generalised Stein's lemma in the fully classical case, but not in the quantum case. Our second technical innovation, therefore, is to perform a second quantisation step to lift the problem to an infinite-dimensional bosonic quantum system; we then solve it there by using techniques from continuous-variable quantum information. Rather remarkably, the second-quantised action of the blurring map corresponds to a pure loss channel. A careful examination of this second quantisation step is the core of our quantum solution.

I. INTRODUCTION

Quantum entanglement, initially regarded as a mathematical oddity of quantum theory [1, 2], is increasingly recognised as a fundamental concept of quantum mechanics [3], and as one of the sources of its technological power. It fuels quantum teleportation [4] and superdense coding [5], it enables the violation of Bell inequalities [6], it plays a critical role in quantum key distribution [7, 8], and it enhances quantum communication on noisy channels [9, 10]. Recently, it has been proposed that its detection in a gravitationally interacting system would provide evidence in favour of the quantisation of gravity [11, 12].

A. Entanglement testing

All of the above applications require the detection of entanglement in unknown states, as this is a necessary step, e.g., in certifying the devices that produce it. This leads us to the key operational task that will be the central focus of this paper: ***entanglement testing***. Imagine that you bought

* ludovico.lami@gmail.com