

Many-body enhancement of energy storage in a waveguide-QED quantum battery

Salvatore Tirone,[1,2,3,*] Gian Marcello Andolina,[4] Giuseppe Calajò,[5] Vittorio Giovannetti,[1] and Davide Rossini[6]

[1] *Scuola Normale Superiore, I-56126 Pisa, Italy*
[2] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[3] *Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands*
[4] *JEIP, UAR 3573 CNRS, Collège de France, PSL Research University, 11 Place Marcelin Berthelot, F-75321 Paris, France*
[5] *Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Padova, I-35131 Padova, Italy*
[6] *Dipartimento di Fisica dell'Università di Pisa and INFN, Largo Pontecorvo 3, I-56127 Pisa, Italy*

arXiv:2410.14555v3 [quant-ph] 23 Jul 2025

Quantum batteries have demonstrated remarkable charging properties, showing that a quantum advantage is possible in the realm of quantum thermodynamics. However, finding an effective strategy to store energy for long periods remains crucial in these systems. Here, we investigate different configurations of a waveguide-QED system acting as a quantum battery and show that, in this context, collective effects can slow down the self-discharging time of the battery, thus improving the storage time. Specifically, when the artificial atoms of the array are arranged randomly, the energy and ergotropy of the optical system are shown to decay at a subexponential rate over long periods, in contrast to the energy decay of a single atom in a waveguide. In the case of atoms arranged in an ordered lattice, collective effects slow down dissipative discharging only for a specific lattice spacing. Thus, in both configurations, collective effects can be used to boost the energy-protection properties of optical systems.

I. INTRODUCTION

In the last decade, quantum technologies have been a topic of intense research and development. These technologies take advantage of the unique properties of quantum physics, such as coherence and entanglement, for practical applications. This has led to significant advances in areas like quantum computing [1], sensing [2], communication, and simulation [3]. However, until recently [4], the crucial area of energy production and management has been only partially addressed within the realm of quantum technologies. As an example, quantum memories have been extensively investigated within the field of quantum information as devices capable of storing quantum states, while preserving their fragile quantum properties [5–7]. In contrast, the concept of quantum batteries (QBs)—devices where energy can be injected, stored, and eventually recovered [8]—has only recently garnered increasing interest within the field of quantum thermodynamics.

A QB [8, 9] is composed of multiple identical units, also called quantum cells, often modeled as two-level systems (TLSs). Energy can be stored in these units by exciting them to a higher energy state. Such energy can then be released by allowing the units to relax back to their lower energy state, with the energy difference being used to perform work. The interest in these systems stems from the fact that entanglement (or many-body interactions) between the units of the battery could allow for a super-extensive charging power [10–19] where the battery could be charged faster collectively. Due to this property, most previous studies mainly focused on the charging step, with the aim of increasing the charging power by means of collective effects.

However, a truly efficient QB should be capable of more than just quick charging; it must also be able to store energy for a significant amount of time. In fact, a quantum device that can absorb energy quickly but fails to retain it for a long period does not fit the definition of a "battery", but rather, it might be more appropriate to term it as an "absorber". Various works in the literature deal with the storage phase [20–43]. Our approach, however, introduces a novel direction by utilizing the collective effects of an optical system that exhibits many-body localization properties alongside other distinctive features, as elaborated in the following. Moreover, this system not only holds the potential for developing a QB with a large number of sites, but it is also well-suited for experimental investigation in a waveguide-quantum electrodynamics (QED) setup [44, 45], offering a realistic pathway to test and validate our results.

In this paper, we consider the experimentally relevant platform of waveguide QED, which has emerged in the latest years as a powerful and promising light-matter interface. Waveguide QED broadly refers to scenarios where one or more quantum emitters are coupled to the light confined into a one-dimensional photonic channel [44–47]. This setting has been implemented in the optical regime, with cold atoms or quantum dots coupled to optical fibers and photonic crystal waveguides [48–52], and at microwave frequencies with superconducting circuits, where artificial atoms are strongly coupled to on-chip microwave resonators or transmission lines [53–56]. The 1D confinement allows for highly controlled interactions between light and matter, which can be used for var-

* s.tirone@uva.nl