

Mediating gates between polar molecules using microwave-dressed Rydberg atoms

Bas Gerritsen,[1,2] Sean R. Muleady,[3,4] Arghavan Safavi-Naini,[1,2] Alexey V. Gorshkov,[3,4] and Jeremy T. Young[1,2]

[1] *Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*
[2] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[3] *Joint Center for Quantum Information and Computer Science, NIST/University of Maryland, College Park, Maryland 20742, USA*
[4] *Joint Quantum Institute, NIST/University of Maryland, College Park, Maryland 20742, USA*

(Dated: September 25, 2026)

We propose a scheme for mediating many simultaneous, fast entangling gates between pairs of polar molecules using Rydberg atoms. By using microwave drives, the dipolar interactions between Rydberg atoms are nullified. Using the freedom left in the microwave dressing parameters, the Rydberg van der Waals interactions are minimized and the Rydberg-molecule interaction is tuned into resonance, allowing for the mediation of a modified iSWAP gate between molecules. In the example of mediating gates between ^{23}Na^{133}Cs molecules with ^{133}Cs atoms, the gate is more than two orders of magnitude faster than an equivalent direct molecule-molecule gate. We model the decay of the dressed Rydberg states and the motion of the atom and molecules and obtain a leading-order estimate of the resulting gate infidelities. Finally, we show that by detecting the state of the Rydberg atom after a gate, a subset of gate errors can be converted into erasure errors and mitigated by post-selection or quantum error-correction.

arXiv:2609.28694v1 [quant-ph] 23 Sep 2026

Polar molecules are a promising platform for applications in quantum simulation and quantum computation [1–3]. These molecules offer both a rich variety of long-lived internal states as well as long-range dipole-dipole interactions. Simultaneously, Rydberg atoms exhibit even stronger interactions, both dipole-dipole and van der Waals, albeit with more susceptibility to dissipation. Tweezer arrays of neutral atoms have positioned themselves as one of the leading platforms for quantum science [3–8]. In light of recent advances in the capabilities of tweezer arrays of polar molecules [9–24], hybrid arrays of both atoms and molecules represent a promising new direction.

By choosing suitable Rydberg states, strong interactions between polar molecules and Rydberg atoms can be realized [17, 25, 26]. These interactions between atoms and molecules can be used for a variety of applications, such as generating entanglement between atoms and polar molecules [26], state-insensitive cooling [27] of polar molecules, non-destructive readout of polar molecules [26, 28–31] and mediation of entangling gates [29, 30] or interactions [32] between molecules. We propose a method for mediating a gate between two polar molecules via an intermediary Rydberg atom. In contrast to related prior proposals [29, 30], we use microwave dressing [31, 33] to minimize unwanted Rydberg-Rydberg interactions, thereby enabling many simultaneous molecular gates.

Interactions between molecules are mediated via a pair of Rydberg states $|a\rangle, |b\rangle$ which have a transition frequency close to that of the transition between two rotational states $|0\rangle, |1\rangle$ of a polar molecule, realizing resonant dipolar flip-flop interactions between atoms and molecules as shown in Fig. 1. By positioning a Rydberg atom between each pair of molecules, the resonant dipolar interaction with the atom mediates a modified iSWAP gate [34] between the two molecules, returning the atom to its original state. In order to simultaneously apply many Rydberg-mediated gates, the Rydberg-molecule interaction needs to be significantly stronger than the other interactions. As the transition dipole moments in a Rydberg atom are typically a factor of $\sim 10^3$ larger than the dipole moment of a polar molecule, Rydberg-Rydberg interactions dominate. To reduce these interactions, we will make use of microwave dressing to engineer dressed Rydberg states which have a negligible dipole-dipole interaction with other Rydberg atoms [31, 33]. Furthermore, through careful tuning of the drive parameters,




FIG. 1. (a) Relevant molecular and Rydberg atom energy levels. When tuned to $E_{ab} = E_m$, the molecules can resonantly exchange excitations with the atom at rate J. (b) Schematic drawing and plot of state populations under exchange gate dynamics. After gate time $t_G = \pi/(J\sqrt{2})$, the molecules have exchanged excitations while the Rydberg atom is returned to its original state.