

REVIEW ARTICLE

Julie R. Ingelfinger, M.D., *Editor*

Cochlear Implantation in Adults

Matthew L. Carlson, M.D.

HEARING LOSS IS ONE OF THE MOST PREVALENT AND UNDERTREATED disabilities worldwide.^{1,2} Often simply considered one of the stigmata of senescence, hearing loss has become increasingly recognized as an important health issue that may lead to increased risks of social isolation,^{3,4} depression,⁵ loss of autonomy,⁶ reduced employability,⁷ and neurocognitive dysfunction,^{3,4,8} in addition to its association with many pragmatic safety concerns.⁹⁻¹¹ In view of these consequences, a growing emphasis has been placed on the prevention, early detection, and treatment of hearing loss.¹²

Although hearing aids suffice for many persons with hearing loss, a subset of the population with greater hearing impairment may benefit from cochlear implantation, a relatively low-risk outpatient procedure that generally leads to improvements in speech understanding and quality of life.¹³ Yet it is estimated that less than 10% of adults in the United States and other developed countries who meet the current criteria for cochlear implantation actually receive this treatment.^{14,15} Multiple factors contribute to the widespread underuse of cochlear implants, including limited awareness of the benefits and risks of cochlear implantation and lack of awareness of the current criteria approved by the Food and Drug Administration (FDA).^{14,16} This review addresses the clinical features of hearing loss, current criteria for cochlear implantation in adults, modern cochlear-implant surgery, and anticipated surgical and audiologic outcomes.

From the Department of Otolaryngology–Head and Neck Surgery and the Department of Neurologic Surgery, Mayo Clinic, Rochester, MN. Address reprint requests to Dr. Carlson at the Dept. of Otolaryngology–Head and Neck Surgery, Mayo Clinic, 200 1st St. SW, Rochester, MN 55905, or at carlson.matthew@mayo.edu.

N Engl J Med 2020;382:1531-42.

DOI: 10.1056/NEJMr1904407

Copyright © 2020 Massachusetts Medical Society.

CLASSIFICATION, EPIDEMIOLOGY, AND THERAPEUTIC OPTIONS

The type and degree of hearing loss are currently classified with the use of pure-tone and speech audiometry. Pure-tone audiometry is a standardized behavioral test that ascertains the quietest decibel threshold a person seated in a sound-attenuating booth can perceive through each ear across a range of frequencies. In contrast, word-recognition testing, which is a primary component of speech audiometry, evaluates a person's ability to recognize a list of monosyllabic words (e.g., car or boat) presented to each ear at a comfortably audible level; the resulting score is reported as the percentage of correct answers.

At least 1.2 million adults in the United States and 50 million adults worldwide are living with severe or profound hearing loss — a level of impairment that is not sufficiently corrected with hearing aids — and might benefit from cochlear implantation.^{14,16-18} The prevalence of hearing loss increases with age, and approximately two thirds of people 70 years of age or older in the United States have hearing loss.^{19,20} The proportion of the population that is older than 65 years of age is projected to double within the next three decades, and a proportional increase in the prevalence of hearing loss is likely.²¹

In the United States, advanced age, recreational or occupational noise exposure, hereditary factors, and exposure to ototoxic medications collectively are the most




Figure 1 (facing page). Classification and Pathogenesis of Hearing Loss.

Hearing loss is broadly categorized as conductive or sensorineural. Conductive hearing loss results primarily from any condition that impedes the transmission of acoustic energy through the external or middle ear to the cochlea (e.g., tympanic-membrane perforation, middle-ear effusion, or ossicular-chain erosion). Conductive hearing loss rarely causes substantial deterioration in word recognition and is almost exclusively managed by surgical reconstruction or conventional hearing aids, not by cochlear implantation. In contrast, sensorineural hearing loss predominantly develops from disorders resulting in cochlear hair-cell injury (e.g., noise exposure or older age) and much less frequently from conditions involving the cochlear nerve or central nervous system.

common causes of sensorineural hearing loss that develops in adulthood (Fig. 1).^{2,12,22} Numerous conditions lead to the development of sensorineural hearing loss, and most share a common pathway whereby damaged or missing sensory hair cells in the cochlea no longer transduce acoustic energy into neural action potentials that are processed by the central nervous system and perceived as understandable speech.

In the search to identify a cure for sensorineural hearing loss, regeneration of cochlear hair cells has become a focus of gene therapy and stem-cell research. Fish, birds, and amphibians are able to regenerate damaged hair cells, but cochlear hair-cell loss in mammals is currently largely irreversible.²³ The two leading strategies under investigation for mammalian regeneration of inner-ear hair cells are the identification of existing progenitor-like cells that might be used to repopulate cochlear hair cells, a technique that has been approached in mouse models that are transgenic for *Nestin*²⁴ and *Lgr5*,²⁵ and the transplantation and engraftment of murine pluripotent stem cells in mouse,²⁶ rat,²⁷ and guinea pig²⁸ models.

Currently, there are no FDA-approved pharmacologic or surgical treatments that reverse sensorineural hearing loss and reinstate normal acoustic hearing. Thus, current clinical management revolves around prevention of hearing loss (e.g., workplace hearing-conservation programs) and rehabilitation through the use of hearing aids or cochlear implants.

The decision to recommend treatment with a cochlear implant is driven by the results of audiometric testing, as well as the demands of the

everyday listening environment. A person with limited unilateral hearing loss or mild bilateral high-frequency hearing loss may function reasonably well in a quiet environment. However, people with greater degrees of hearing loss frequently struggle with everyday communication, particularly in the presence of background noise. Fitted hearing aids provide sufficient benefit for most patients who have relatively good speech recognition (a score $\geq 60\%$ on word-recognition testing) and less than severe-to-profound hearing loss (i.e., ≤ 70 dB hearing level [HL] at low frequencies and ≤ 90 dB HL at high frequencies). However, for patients with poor speech recognition and more advanced bilateral sensorineural hearing loss, cochlear implantation provides the only effective means of auditory rehabilitation currently available. Whereas hearing aids function by amplifying sound, cochlear implants bypass nonfunctional or missing cochlear hair cells and directly stimulate surviving spiral ganglion cells of the distal cochlear nerve, enhancing both audibility and speech recognition (Fig. 2).

ASSESSMENT AND CURRENT CRITERIA FOR COCHLEAR IMPLANTATION

Evaluation of an adult for cochlear implantation entails a detailed hearing assessment by an audiologist and a medical assessment by a cochlear-implant surgeon. The chief objective of the audiologic assessment is to determine whether the patient will receive more benefit from a cochlear implant than from properly fitted hearing aids, whereas the primary goal of the medical assessment is to determine whether surgery is sufficiently safe and feasible for the patient.

During unilateral cochlear-implant surgery in an adult, the typical time under anesthesia is less than 2 hours and limited blood loss occurs; therefore, the perioperative anesthetic risks are relatively low. Most centers now routinely perform cochlear-implant surgery in patients ranging from infants to nonagenarians.^{29,30} In contrast to a subgroup of children with congenital sensorineural hearing loss, adults with postlingual hearing loss generally do not have severe cochlear dysplasia, cochlear-nerve hypoplasia, or a central process that precludes cochlear implantation.³¹ The one notable exception is patients with profound hearing loss after bacterial men-




Figure 2 (facing page). Components and Mechanism of a Cochlear Implant.

The external sound processor converts acoustic sound to a coded electrical signal that is transmitted to the internal device through transcutaneous radiofrequency coupling. The internal device, in turn, relays the signal to the electrode array of the cochlear implant, bypassing damaged sensory hair cells in the cochlea and directly stimulating spiral ganglion cells of the cochlear nerve. Stimulation is delivered in a rapid, sequential, interleaved pattern through which neighboring electrode signals are temporally separated by milliseconds to avoid current summation of adjacent electrodes. This strategy improves sound quality and speech recognition by maximizing channel independence and spectral resolution.

ingitis. Labyrinthitis ossificans develops concomitantly in up to 35% of such patients, causing irreversible scarring within the cochlea.³² If cochlear implantation is being considered in a patient with profound hearing loss due to bacterial meningitis, surgery is generally best performed soon after the onset of hearing loss so that the electrode can be inserted before prohibitive ossification ensues.

Currently, most insurance programs in the United States specify that to qualify for coverage, an adult must have bilateral sensorineural hearing loss that is moderate or more severe at low frequencies and profound at high frequencies, with no more than 50% of sentences understood correctly in the ear that will receive the implant or no more than 60% of sentences understood correctly when both ears are fitted with hearing aids (Fig. 3; and Table S1 in the Supplementary Appendix, available with the full text of this article at NEJM.org).^{14,16} In most cases, the ear with the greater degree of hearing loss is chosen for implantation so that the patient can continue to use a hearing aid in the contralateral ear while adjusting to the new implant. Since the cochlear implant is not typically activated until 2 to 4 weeks after implantation, this strategy also allows most people to resume work and other social activities within 5 to 10 days after surgery. Depending on how well the patient is functioning with the cochlear implant and how much residual hearing is present in the ear without the implant, implantation in the second ear may be performed at a later date if the patient wishes.

There is a common misconception that patients


must have bilateral profound sensorineural hearing loss to meet current criteria for cochlear implantation in adults. FDA device labeling has been broadened and currently encompasses device implantation in patients with varying degrees of preoperative residual acoustic hearing. In 2014, a hybrid cochlear-implant system with a short electrode was approved for use in adults with normal hearing to moderate hearing loss at low frequencies, severe to profound hearing loss at middle-to-high frequencies, and poor word recognition (www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMA/pma.cfm?id=P130016). The use of this specialized short electrode reduces apical cochlear trauma and allows for preservation of preoperative low-frequency acoustic hearing in 66% of cases (Fig. 3).³³ When functional acoustic hearing is preserved after surgery, patients can use a hearing aid in the ear with the cochlear implant, a strategy called hybrid stimulation, which may confer enhanced speech perception and music appreciation.³⁴ In July 2019, cochlear implantation was approved by the FDA for patients, 5 years of age or older, who have profound hearing loss in one ear and a range from normal hearing to moderate hearing loss in the other ear (www.fda.gov/medical-devices/recently-approved-devices/med-el-cochlear-implant-system-p000025s104).

There is no screening questionnaire or finding on clinical examination that can reliably predict whether a patient will benefit from cochlear implantation or will meet current FDA labeling criteria for implantation in adults. Furthermore, the degree of hearing loss is commonly underestimated during a typical face-to-face clinic visit. Outpatient encounters typically occur in a quiet room, where patients can compensate with lip reading and other visual cues or discreetly rely on an accompanying friend or family member to communicate. As a general rule, people who are struggling to understand speech during routine everyday activities, despite using hearings aids, may benefit from cochlear implantation (Table 1).^{15,35,36} Ultimately, a comprehensive hearing assessment can be considered when more advanced hearing loss is suspected.


COCHLEAR-IMPLANT TECHNOLOGY

Currently, three device companies manufacture FDA-approved cochlear implants. Notwithstand-


A




B FDA Eligibility for Conventional Cochlear Implant



C FDA Eligibility for Hybrid Cochlear Implant



Conventional cochlear implant



Hybrid cochlear implant




Figure 3 (facing page). Degree of Hearing Loss and Criteria for Cochlear Implantation in Adults.

Panel A shows the various degrees of hearing loss. Panel B shows that historically, only adults with bilateral, profound hearing loss underwent cochlear implantation (initial Food and Drug Administration [FDA] eligibility criteria), whereas currently, adults with moderate or more severe low-frequency hearing loss and profound high-frequency hearing loss meet the criteria for receipt of a conventional cochlear implant. Both the degree of hearing loss and word- or sentence-recognition scores are used to assess audiometric performance before cochlear implantation (see Table S1 in the Supplementary Appendix). Neurosensory cells that collect low frequencies aggregate in the apex of the cochlea, and those that collect high frequencies aggregate in the base. This serendipitous arrangement permits a cochlear-implant electrode to selectively activate different regions of the cochlea in order to precisely control pitch. Panel C shows that the tonotopic organization of the cochlea also allows specialized shorter-length electrodes, used in hybrid cochlear-implants, to treat mid- and high-frequency hearing loss through preferential electrical stimulation of the basal cochlea while simultaneously preserving preoperative low-frequency acoustic hearing by limiting trauma to the more apical regions of the cochlea.

ing small variations, virtually all commercially available devices share a similar design: the external component contains a microphone, battery, sound processor, and transmitting coil, and the internal component includes a radiofrequency receiver coil, microprocessor-based stimulator, and multichannel electrode (Fig. 2). The intracochlear electrode contains 12 to 22 active contacts, depending on the model, that independently stimulate different regions of the cochlea. Cochlear-implant technology takes advantage of the tonotopic anatomy of the cochlea, with neurons that collect low-frequency signals aggregating in the apical regions of the cochlea, and high-frequency signals collected in the basal regions. Precise pitch control is achieved through altering the stimulation rate and spatial location of stimulation along the tonotopic axis of the cochlea. Sound intensity is controlled by modifying the current amplitude and pulse duration at each individual electrode contact. The fact that many implant recipients can decipher complex speech with as few as 8 to 10 independent stimulating electrodes suggests that central nervous system processing and neuroplasticity play pivotal roles in speech perception.^{13,37,38}

Table 1. Clinical Screening Questions to Help Identify Adults Who May Benefit from Cochlear Implantation.

When using your hearing aid (or aids), are you able to carry on a conversation with another person when you cannot directly see his or her face?
Are you able to use the telephone without using visual aids such as caption or video?
Do you struggle when communicating with others at large gatherings such as dinner parties, even when you are using hearing aids?
Do you avoid social gatherings, meetings, and other events at which there is background noise, because you are unable to hear conversations well, even when you are using hearing aids?

External cochlear-implant components have become increasingly miniaturized, and most current designs can be concealed relatively well under medium-length or long hair (Fig. S1). Virtually all current sound processors integrate smartphone and wireless Bluetooth technology, offer specific programs for various listening environments, and incorporate water-resistant encasements. As of June 2019, all manufacturers of FDA-approved cochlear implants integrate internal magnets that are “MRI conditional” (i.e., safe for a patient to undergo magnetic resonance imaging [MRI] under very specific conditions provided in the labeling) in a field strength of up to 3 Tesla (www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?start_search=1&PMANumber=P970051&SupplementNumber=S183). This development is particularly important, since at least half of all cochlear-implant recipients will have a clinical indication for at least one MRI study during their lifetime.^{39,40} Currently, several commercial implant manufacturers are evaluating the safety and feasibility of a totally implantable cochlear-implant system.^{41,42} A totally implantable system not only would render the device virtually invisible but also would support full-time use. With current devices, the external component is typically removed while the patient is swimming, showering, and sleeping.

RISKS, BENEFITS, AND OUTCOMES OF COCHLEAR IMPLANTATION

SURGICAL AND DEVICE-RELATED COMPLICATIONS

As noted above, cochlear-implant surgery has evolved into a relatively low-risk outpatient surgical procedure performed while the patient is



A video showing
cochlear-implant
surgery is available
at NEJM.org

under general anesthesia. A narrated video of cochlear-implant surgery in adults can be viewed at NEJM.org. Data on the estimated prevalence of complications associated with cochlear implantation, compiled from 10 large studies, are presented in Table S2 and summarized here.⁴³⁻⁵² The overall prevalence of complications is 12.8% (reported in 964 of 7513 patients),^{43-49,51,52} and the prevalence of major complications is 2.7% (207 of 7542).^{43,44,46-52} The prevalence of specific complications includes wound infection in 1.9% of patients (104 of 5556),^{43,44,46-48,51,52} hematoma in 1.1% (85 of 7513),^{43-49,51,52} chronic surgical-site pain in 1.7% (49 of 2889),^{44,46,47,49,51,52} persistent vestibular symptoms in 2.2% (102 of 4664),^{44,46,47,49,51,52} and permanent facial-nerve paralysis in 0.1% (7 of 8779).⁴³⁻⁵² The risk of postoperative meningitis has been strongly associated with an age of less than 6 years (when recurrent otitis media is most prevalent), with the presence of inner-ear malformations, and with the use of a specific cochlear-implant electrode positioner that was removed from the market in July 2002.⁵³ With the use of recent device models, the prevalence of postoperative meningitis was less than 0.1% (5 of 7167).^{44-47,49-52} This risk might be further attenuated by adhering to Centers for Disease Control and Prevention guidelines for pneumococcal vaccine prophylaxis.⁵⁴ Total device failure requiring reimplantation has been reported in 1.9% of cases (125 of 6461).^{44-46,48,49,51,52}

Until relatively recently, the loss of preoperative acoustic hearing in the ear that received the implant was considered an unavoidable consequence of cochlear-implant surgery.⁵⁵ However, in the late 1990s, several reports documented the feasibility of hearing preservation after cochlear implantation.^{56,57} As FDA labeling criteria have expanded to include patients with better residual hearing, the ability to preserve functional preoperative hearing has become a priority. A meta-analysis by Santa Maria et al. showed that among adults who received cochlear implants with a conventional-length electrode (186 patients in 13 studies), preoperative low-frequency (≤ 1000 Hz) hearing was fully preserved (i.e., <10 dB loss) in 27.4% of patients and partially preserved (i.e., between 10 and 20 dB loss) in 25.3% but was not preserved (i.e., >20 dB loss) in 47.3%.⁵⁸ Thus, it appears that although most patients today benefit from the electrical stimu-

lation provided by the cochlear implant, approximately half of adults who have functional preoperative acoustic hearing lose this residual hearing as a complication of surgery. Beyond modifying surgical techniques and electrode designs, research efforts have focused on advances that include automated electrode insertion,⁵⁹ glucocorticoid-eluting electrodes,⁶⁰ and real-time intraoperative electrophysiological feedback⁶¹ to mitigate the risk of early or delayed acoustic hearing loss after cochlear-implant surgery.

SPEECH PERCEPTION AND SOUND QUALITY

Refinements in surgery, device design, programming strategy, and FDA labeling criteria have resulted in substantial improvements in multichannel cochlear-implant performance since the early devices were first approved for commercial use in 1985.⁴¹ After cochlear implantation, there is a variable period of acclimatization to the new electrical input. Many recipients initially report that voices sound high-pitched and mechanical, though sound quality generally improves during the following 3 to 6 months. Sound clips are available as examples of what a cochlear-implant recipient hears (https://asha.figshare.com/articles/single-sides_deaf_ci_voice_quality_dorman_et_al_2019_9341651).^{62,63} For the first year after surgery, cochlear-implant recipients need to attend several device-programming sessions at which the sound quality and loudness of the implant are adjusted to improve speech recognition. During this time, patients are also encouraged to use their implant consistently throughout the day while watching television, listening to familiar music or audiobooks, or reading aloud in order to train the central nervous system to reconcile what is heard with known words.⁶⁴ At most centers, there is a routine delay of 2 to 4 weeks after surgery before the device is turned on. Recently, device activation within 24 hours after surgery was successfully implemented at several centers to expedite rehabilitation and potentially reduce the number of early postoperative return visits.^{65,66}

Although the rate at which speech perception improves after cochlear implantation is variable, increases are usually steepest within the first 6 months of use. However, continued progress can be seen up to 3 years after surgery.^{67,68} Data from an unselected, consecutive series of 259 adults (277 ears) who recently received implants at the Mayo Clinic provide information regarding

the audiometric profile and anticipated outcomes with the use of current device models (unpublished data). The median preoperative scores on tests of monosyllabic-word and sentence recognition were 8% (interquartile range [IQR], 0 to 24) and 7% (IQR, 0 to 29), respectively. Within the first 12 months after cochlear implantation, the median postoperative scores for word and sentence recognition were 58% (IQR, 36 to 72) and 75% (IQR, 53 to 91), respectively (Table S3 and Fig. S2).

ENVIRONMENTAL SOUND PERCEPTION

In addition to improving speech perception, cochlear implants have the capacity to restore sound thresholds to normal or near-normal levels. Although this benefit of implants is commonly underappreciated in the medical literature, patients frequently reflect positively on their newfound ability to hear many quiet or high-pitched environmental sounds, such as the faucet dripping, the click of a car's turn signal, or birds chirping. Beyond such perceptual benefits, enhanced environmental sound awareness improves personal safety, increases autonomy, and alleviates possible fears of not hearing a fire alarm, oncoming traffic, a doorbell, or other "directing" sounds.^{10,11,69}

QUALITY OF LIFE AND COST-EFFECTIVENESS

Although scores for word and sentence recognition are the primary benchmarks used to assess outcome, these traditional metrics do not adequately capture the complex and multifaceted benefits that many implant recipients report, particularly with regard to social and emotional domains. A substantial body of literature elucidates consistent improvements in quality of life after implantation that appear to be independent of audiometric testing performance.^{70,71} In a recent pooled analysis, Crowson et al. reported that the average improvement in quality of life for adult recipients of unilateral implants was 44.3% with the use of generic measures (among 954 patients in 15 studies) and 89.1% with the use of disease-specific instruments (among 360 patients in 11 studies).⁷¹ Moreover, 6 studies (with a total of 674 patients) that assessed cost-effectiveness with the use of the Health Utilities Index showed a weighted mean cost (in U.S. dollars) per quality-adjusted life-year (QALY) gained of \$23,310 (range, \$9,426 to \$33,656).⁷²⁻⁷⁷

For context, a ratio of less than \$50,000 per QALY has long served as the accepted benchmark for value in the United States.⁷⁸

KEY ISSUES MOVING FORWARD

INDIVIDUAL VARIATION IN OUTCOMES

Although most implant recipients have significant gains in speech perception, a subgroup of recipients have poor outcomes, even after extended use of the implant and supplemental rehabilitation.^{38,67,79} A recent analysis showed that up to 16% of adult cochlear-implant recipients had poor long-term outcomes (defined as a word- or sentence-recognition score of less than 30% at 12 months) (unpublished data; Table S3 and Fig. S2). Rumeau et al. reported that 9 of 26 experienced adult implant recipients (35%) were unable to routinely use the telephone.⁸⁰ The duration of deafness before implantation and the preoperative speech-perception scores have the highest predictive value for the speech-perception outcome among adults who have received cochlear implants, whereas age at the time of implantation and the cause of sensorineural hearing loss do not appreciably influence the outcome in most studies.⁸¹⁻⁸³ Nevertheless, even the most robust predictive models cannot fully account for the observed range in speech-perception scores.^{67,84} Understanding the complex mechanisms underlying outcome variation and the development of new interventions to successfully remediate poor test performance remain two critical objectives in the field today.

ASSOCIATION BETWEEN COCHLEAR IMPLANTATION AND COGNITIVE FUNCTION

Growing evidence links hearing loss with cognitive impairment and dementia.^{8,85} In the 2017 report from the Lancet Commission on Dementia Prevention, Intervention, and Care, Livingston et al. estimated that up to 35% of dementia cases are potentially preventable, with hearing loss considered the largest modifiable risk factor.⁸ As a natural extension, whether rehabilitation with hearing aids or cochlear implants can successfully mitigate the risk of cognitive impairment or dementia has become an area of active research.⁸⁶ Five prospective longitudinal studies involving a total of 259 patients have identified improvements in attention, processing speed, and working memory after cochlear implantation.⁸⁶⁻⁹⁰

SUMMARY

Hearing loss is often viewed as an inconsequential process or an accepted part of aging rather than an important health risk that warrants treatment. Patients who struggle to understand speech in typical everyday listening environments despite using hearing aids may benefit from cochlear implantation. Age and coexisting medical conditions generally do not preclude cochlear-implant surgery and have a limited effect on performance outcomes. Although most adult cochlear-implant recipients gain clinically significant improvements in speech recognition and quality of life, a subgroup of patients do not have such benefits. The variation in outcomes, the ability to reliably preserve natural hearing

after surgery, and the potential role of hearing aids and cochlear implants in mitigating the risk of late-life dementia represent three important areas of active research.

Dr. Carlson reports receiving meal reimbursement from Advanced Bionics and Stryker, meal reimbursement and travel support from Cochlear Americas, MED-EL, and Oticon Medical, and holding pending patents 62/346,306, PCT/US2017/035617, and WO2017213978A1 on devices and methods for treating tinnitus with the use of electrical stimulation. No other potential conflict of interest relevant to this article was reported.

Disclosure forms provided by the author are available with the full text of this article at NEJM.org.

I thank my colleagues at Mayo Clinic, including John P. Marinelli, M.D., for editorial assistance with earlier versions of the manuscript, Andrew J. Goates, M.D., for editorial assistance with an earlier version of the narrated video, and Christine M. Lohse, M.S., for assistance with statistical analysis of the cochlear-implant outcome data presented in the Supplementary Appendix.

REFERENCES

1. GBD 2015 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016; 388:1545-602.
2. Wilson BS, Tucci DL, Merson MH, O'Donoghue GM. Global hearing health care: new findings and perspectives. *Lancet* 2017;390:2503-15.
3. Lin FR, Yaffe K, Xia J, et al. Hearing loss and cognitive decline in older adults. *JAMA Intern Med* 2013;173:293-9.
4. Maharani A, Pendleton N, Leroy I. Hearing impairment, loneliness, social isolation, and cognitive function: longitudinal analysis using English Longitudinal Study on Ageing. *Am J Geriatr Psychiatry* 2019;27:1348-56.
5. Manrique-Huarte R, Calavia D, Huarte Irujo A, Girón L, Manrique-Rodríguez M. Treatment for hearing loss among the elderly: auditory outcomes and impact on quality of life. *Audiol Neurotol* 2016;21: Suppl 1:29-35.
6. Sonnet MH, Montaut-Verient B, Niemier JY, Hoen M, Ribeyre L, Parietti-Winkler C. Cognitive abilities and quality of life after cochlear implantation in the elderly. *Otol Neurotol* 2017;38(8):e296-e301.
7. Clinkard D, Barbic S, Amoodi H, Shipp D, Lin V. The economic and societal benefits of adult cochlear implantation: a pilot exploratory study. *Cochlear Implants Int* 2015;16:181-5.
8. Livingston G, Sommerlad A, Orgeta V, et al. Dementia prevention, intervention, and care. *Lancet* 2017;390:2673-734.
9. Reed NS, Altan A, Deal JA, et al. Trends in health care costs and utilization associated with untreated hearing loss over 10 years. *JAMA Otolaryngol Head Neck Surg* 2019;145:27-34.
10. Harris MS, Boyce L, Pisoni DB, Shafiro V, Moberly AC. The relationship between environmental sound awareness and speech recognition skills in experienced cochlear implant users. *Otol Neurotol* 2017;38(9):e308-e314.
11. Ng ZY, Lamb B, Harrigan S, Archbold S, Athalye S, Allen S. Perspectives of adults with cochlear implants on current CI services and daily life. *Cochlear Implants Int* 2016;17:Suppl 1:89-93.
12. Cunningham LL, Tucci DL. Hearing loss in adults. *N Engl J Med* 2017;377: 2465-73.
13. Wilson BS. Toward better representations of sound with cochlear implants. *Nat Med* 2013;19:1245-8.
14. Sorkin DL, Buchman CA. Cochlear implant access in six developed countries. *Otol Neurotol* 2016;37(2):e161-e164.
15. Holder JT, Reynolds SM, Sunderhaus LW, Gifford RH. Current profile of adults presenting for preoperative cochlear implant evaluation. *Trends Hear* 2018;22: 1-16.
16. Sorkin DL. Cochlear implantation in the world's largest medical device market: utilization and awareness of cochlear implants in the United States. *Cochlear Implants Int* 2013;14:Suppl 1:S4-S12.
17. Goman AM, Lin FR. Prevalence of hearing loss by severity in the United States. *Am J Public Health* 2016;106:1820-2.
18. WHO global estimates on prevalence of hearing loss: mortality and burden of diseases and prevention of blindness and deafness. Geneva: World Health Organization, 2012 (https://www.who.int/pbd/deafness/WHO_GE_HL.pdf).
19. Wattamwar K, Qian ZJ, Otter J, et al. Increases in the rate of age-related hearing loss in the older old. *JAMA Otolaryngol Head Neck Surg* 2017;143:41-5.
20. Lin FR, Niparko JK, Ferrucci L. Hearing loss prevalence in the United States. *Arch Intern Med* 2011;171:1851-2.
21. Colby SL, Ortman JM. Projections of the size and composition of the U.S. population: 2014 to 2060. Current Population Reports P25-114 (March). Washington, DC: Census Bureau, 2015.
22. Agrawal Y, Platz EA, Niparko JK. Prevalence of hearing loss and differences by demographic characteristics among US adults: data from the National Health and Nutrition Examination Survey, 1999-2004. *Arch Intern Med* 2008;168:1522-30.
23. Ronaghi M, Nasr M, Heller S. Concise review: inner ear stem cells — an oxymoron, but why? *Stem Cells* 2012;30:69-74.
24. Watanabe R, Morell MH, Miller JM, et al. Nestin-expressing cells in the developing, mature and noise-exposed cochlear epithelium. *Mol Cell Neurosci* 2012;49: 104-9.
25. Shi F, Kempfle JS, Edge AS. Wnt-responsive Lgr5-expressing stem cells are hair cell progenitors in the cochlea. *J Neurosci* 2012;32:9639-48.
26. Takeda H, Minoda R, Miwa T, Yamada T, Ise M. Transplanting mouse induced pluripotent stem cells into mouse otocysts in vivo. *Neurosci Lett* 2017;647:153-8.
27. Zhao LD, Li L, Wu N, et al. Migration and differentiation of mouse embryonic stem cells transplanted into mature cochlea of rats with aminoglycoside-induced hearing loss. *Acta Otolaryngol* 2013;133: 136-43.
28. Nayagam BA, Backhouse SS, Cimenkaya C, Shepherd RK. Hydrogel limits

- stem cell dispersal in the deaf cochlea: implications for cochlear implants. *J Neural Eng* 2012;9:065001.
29. Carlson ML, Breen JT, Gifford RH, et al. Cochlear implantation in the octogenarian and nonagenarian. *Otol Neurotol* 2010;31:1343-9.
 30. Carlson ML, Sladen DP, Gurgel RK, Tombers NM, Lohse CM, Driscoll CL. Survey of the American Neurotology Society on Cochlear Implantation: part 1, candidacy assessment and expanding indications. *Otol Neurotol* 2018;39(1):e12-e19.
 31. Brown CS, Choi KJ, Kaylie DM. Preoperative imaging findings and cost in adults with postlingual deafness prior to cochlear implant. *Ann Otol Rhinol Laryngol* 2018;127:270-4.
 32. Caye-Thomasen P, Dam MS, Omland SH, Mantoni M. Cochlear ossification in patients with profound hearing loss following bacterial meningitis. *Acta Otolaryngol* 2012;132:720-5.
 33. Roland JT Jr, Gantz BJ, Waltzman SB, Parkinson AJ. United States multicenter clinical trial of the cochlear nucleus hybrid implant system. *Laryngoscope* 2016;126:175-81.
 34. Gantz BJ, Turner C, Gfeller KE, Lowder MW. Preservation of hearing in cochlear implant surgery: advantages of combined electrical and acoustical speech processing. *Laryngoscope* 2005;115:796-802.
 35. Sladen DP, Gifford RH, Haynes D, et al. Evaluation of a revised indication for determining adult cochlear implant candidacy. *Laryngoscope* 2017;127:2368-74.
 36. Luntz M, Khuri M, Khayr R, Gedaon Khuri W, Yehudai N, Shpak T. Cochlear implantation outcome in straightforward cases: can we do more for adults who cannot understand speech using hearing aids alone? *Acta Otolaryngol* 2018;138:1070-9.
 37. O'Donoghue G. Cochlear implants — science, serendipity, and success. *N Engl J Med* 2013;369:1190-3.
 38. Anderson CA, Lazard DS, Hartley DE. Plasticity in bilateral superior temporal cortex: effects of deafness and cochlear implantation on auditory and visual speech processing. *Hear Res* 2017;343:138-49.
 39. Edmonson HA, Carlson ML, Patton AC, Watson RE. MR imaging and cochlear implants with retained internal magnets: reducing artifacts near highly inhomogeneous magnetic fields. *Radiographics* 2018;38:94-106.
 40. Carlson ML, Neff BA, Link MJ, et al. Magnetic resonance imaging with cochlear implant magnet in place: safety and imaging quality. *Otol Neurotol* 2015;36:965-71.
 41. Carlson ML, Driscoll CL, Gifford RH, McMenomey SO. Cochlear implantation: current and future device options. *Otolaryngol Clin North Am* 2012;45:221-48.
 42. Australian New Zealand Clinical Trials Registry. Feasibility of the Cochlear™ Nucleus® cochlear implant in a newly implanted adult population. Sydney: NHMRC Clinical Trials Centre, University of Sydney (<https://www.anzctr.org.au/ACTRN12618001593224.aspx>).
 43. Black B. Keyhole cochlear implantation: current status. *Otol Neurotol* 2011;32:1459-63.
 44. Brito R, Monteiro TA, Leal AF, Tsuji RK, Pinna MH, Bento RF. Surgical complications in 550 consecutive cochlear implantation. *Braz J Otorhinolaryngol* 2012;78:80-5.
 45. Ding X, Tian H, Wang W, Zhang D. Cochlear implantation in China: review of 1,237 cases with an emphasis on complications. *ORL J Otorhinolaryngol Relat Spec* 2009;71:192-5.
 46. Halawani R, Aldhafeeri A, Alajlan S, Alzhrani F. Complications of post-cochlear implantation in 1027 adults and children. *Ann Saudi Med* 2019;39:77-81.
 47. Hansen S, Anthonsen K, Stangerup SE, Jensen JH, Thomsen J, Cayé-Thomasen P. Unexpected findings and surgical complications in 505 consecutive cochlear implantations: a proposal for reporting consensus. *Acta Otolaryngol* 2010;130:540-9.
 48. Jiang Y, Gu P, Li B, et al. Analysis and management of complications in a cohort of 1,065 minimally invasive cochlear implantations. *Otol Neurotol* 2017;38:347-51.
 49. Kim CS, Oh SH, Chang SO, Kim HM, Hur DG. Management of complications in cochlear implantation. *Acta Otolaryngol* 2008;128:408-14.
 50. Petersen H, Walshe P, Glynn F, et al. Occurrence of major complications after cochlear implant surgery in Ireland. *Cochlear Implants Int* 2018;19:297-306.
 51. Theunisse HJ, Pennings RJE, Kunst HPM, Mulder JJ, Mylanus EAM. Risk factors for complications in cochlear implant surgery. *Eur Arch Otorhinolaryngol* 2018;275:895-903.
 52. Venail F, Sicard M, Piron JP, et al. Reliability and complications of 500 consecutive cochlear implantations. *Arch Otolaryngol Head Neck Surg* 2008;134:1276-81.
 53. Reefhuis J, Honein MA, Whitney CG, et al. Risk of bacterial meningitis in children with cochlear implants. *N Engl J Med* 2003;349:435-45.
 54. Cochlear implants and vaccination recommendations. Atlanta: Centers for Disease Control and Prevention, August 8, 2019 (<https://www.cdc.gov/vaccines/vpd/mening/public/dis-cochlear-faq-gen.html>).
 55. Boggess WJ, Baker JE, Balkany TJ. Loss of residual hearing after cochlear implantation. *Laryngoscope* 1989;99:1002-5.
 56. Hodges AV, Schloffman J, Balkany T. Conservation of residual hearing with cochlear implantation. *Am J Otol* 1997;18:179-83.
 57. Fraysse B, Dillier N, Klenzner T, et al. Cochlear implants for adults obtaining marginal benefit from acoustic amplification: a European study. *Am J Otol* 1998;19:591-7.
 58. Santa Maria PL, Gluth MB, Yuan Y, Atlas MD, Blevins NH. Hearing preservation surgery for cochlear implantation: a meta-analysis. *Otol Neurotol* 2014;35(10):e256-e269.
 59. Torres R, Jia H, Drouillard M, et al. An optimized robot-based technique for cochlear implantation to reduce array insertion trauma. *Otolaryngol Head Neck Surg* 2018;159:900-7.
 60. Bas E, Bohorquez J, Goncalves S, et al. Electrode array-eluted dexamethasone protects against electrode insertion trauma induced hearing and hair cell losses, damage to neural elements, increases in impedance and fibrosis: a dose response study. *Hear Res* 2016;337:12-24.
 61. Giardina CK, Brown KD, Adunka OF, et al. Intracochlear electrocochleography: response patterns during cochlear implantation and hearing preservation. *Ear Hear* 2019;40:833-48.
 62. Dorman MF, Natale SC, Butts AM, Zeitler DM, Carlson ML. The sound quality of cochlear implants: studies with single-sided deaf patients. *Otol Neurotol* 2017;38(8):e268-e273.
 63. Dorman MF, Natale SC, Zeitler DM, Baxter L, Noble JH. Looking for Mickey Mouse™ but finding a munchkin: the perceptual effects of frequency upshifts for single-sided deaf, cochlear implant patients. *J Speech Lang Hear Res* 2019;62:3493-9.
 64. Harris MS, Capretta NR, Henning SC, Feeney L, Pitt MA, Moberly AC. Postoperative rehabilitation strategies used by adults with cochlear implants: a pilot study. *Laryngoscope Invest Otolaryngol* 2016;1:42-8.
 65. Batuk MO, Yarali M, Cinar BC, et al. Is early cochlear implant device activation safe for all on-the-ear and off-the-ear sound processors? *Audiol Neurotol* 2019;24:279-84.
 66. Roux-Vaillard S, Pineau A, Laccourreye L, Boucher S. Immediate activation after cochlear implantation: preliminary study. *Eur Ann Otorhinolaryngol Head Neck Dis* 2020;137:17-20.
 67. Holden LK, Finley CC, Firszt JB, et al. Factors affecting open-set word recognition in adults with cochlear implants. *Ear Hear* 2013;34:342-60.
 68. Cusumano C, Friedmann DR, Fang Y, Wang B, Roland JT Jr, Waltzman SB. Performance plateau in prelingually and postlingually deafened adult cochlear implant recipients. *Otol Neurotol* 2017;38:334-8.

69. Powell W, Jacobs JA, Noble W, Bush ML, Snell-Rood C. Rural adult perspectives on impact of hearing loss and barriers to care. *J Community Health* 2019;44:668-74.
70. McRackan TR, Bauschard M, Hatch JL, et al. Meta-analysis of quality-of-life improvement after cochlear implantation and associations with speech recognition abilities. *Laryngoscope* 2018;128:982-90.
71. Crowson MG, Semenov YR, Tucci DL, Niparko JK. Quality of life and cost-effectiveness of cochlear implants: a narrative review. *Audiol Neurotol* 2017;22:236-58.
72. Wyatt JR, Niparko JK, Rothman M, deLissovoy G. Cost utility of the multichannel cochlear implants in 258 profoundly deaf individuals. *Laryngoscope* 1996;106:816-21.
73. Palmer CS, Niparko JK, Wyatt JR, Rothman M, de Lissovoy G. A prospective study of the cost-utility of the multichannel cochlear implant. *Arch Otolaryngol Head Neck Surg* 1999;125:1221-8.
74. UK Cochlear Implant Study Group. Criteria of candidacy for unilateral cochlear implantation in postlingually deafened adults II: cost-effectiveness analysis. *Ear Hear* 2004;25:336-60.
75. Lee HY, Park EC, Kim HJ, Choi JY, Kim HN. Cost-utility analysis of cochlear implants in Korea using different measures of utility. *Acta Otolaryngol* 2006;126:817-23.
76. Kuthubutheen J, Mittmann N, Amoodi H, Qian W, Chen JM. The effect of different utility measures on the cost-effectiveness of bilateral cochlear implantation. *Laryngoscope* 2015;125:442-7.
77. Francis HW, Chee N, Yeagle J, Cheng A, Niparko JK. Impact of cochlear implants on the functional health status of older adults. *Laryngoscope* 2002;112:1482-8.
78. Neumann PJ, Cohen JT, Weinstein MC. Updating cost-effectiveness — the curious resilience of the \$50,000-per-QALY threshold. *N Engl J Med* 2014;371:796-7.
79. Pisoni DB, Kronenberger WG, Harris MS, Moberly AC. Three challenges for future research on cochlear implants. *World J Otorhinolaryngol Head Neck Surg* 2018;3:240-54.
80. Rumeau C, Frère J, Montaut-Verient B, Lion A, Gauchard G, Parietti-Winkler C. Quality of life and audiological performance through the ability to phone of cochlear implant users. *Eur Arch Otorhinolaryngol* 2015;272:3685-92.
81. Semenov YR, Martinez-Monedero R, Niparko JK. Cochlear implants: clinical and societal outcomes. *Otolaryngol Clin North Am* 2012;45:959-81.
82. Chen SY, Grisel JJ, Lam A, Golub JS. Assessing cochlear implant outcomes in older adults using HERMES: a national Web-based database. *Otol Neurotol* 2017;38(10):e405-e412.
83. van der Jagt MA, Briaire JJ, Verbist BM, Frijns JH. Comparison of the HiFocus Mid-Scala and HiFocus 1J electrode array: angular insertion depths and speech perception outcomes. *Audiol Neurotol* 2016;21:316-25.
84. Rubinstein JT, Parkinson WS, Tyler RS, Gantz BJ. Residual speech recognition and cochlear implant performance: effects of implantation criteria. *Am J Otol* 1999;20:445-52.
85. Gurgel RK, Ward PD, Schwartz S, Norton MC, Foster NL, Tschanz JT. Relationship of hearing loss and dementia: a prospective, population-based study. *Otol Neurotol* 2014;35:775-81.
86. Sarant J, Harris D, Busby P, et al. The effect of cochlear implants on cognitive function in older adults: initial baseline and 18-month follow up results for a prospective international longitudinal study. *Front Neurosci* 2019;13:789.
87. Völter C, Götze L, Dazert S, Falkenstein M, Thomas JP. Can cochlear implantation improve neurocognition in the aging population? *Clin Interv Aging* 2018;13:701-12.
88. Jayakody DMP, Friedland PL, Nel E, Martins RN, Atlas MD, Sohrabi HR. Impact of cochlear implantation on cognitive functions of older adults: pilot test results. *Otol Neurotol* 2017;38(8):e289-e295.
89. Cosetti MK, Pinkston JB, Flores JM, et al. Neurocognitive testing and cochlear implantation: insights into performance in older adults. *Clin Interv Aging* 2016;11:603-13.
90. Mosnier I, Bebear JP, Marx M, et al. Improvement of cognitive function after cochlear implantation in elderly patients. *JAMA Otolaryngol Head Neck Surg* 2015;141:442-50.

Copyright © 2020 Massachusetts Medical Society.

IMAGES IN CLINICAL MEDICINE

The *Journal* welcomes consideration of new submissions for Images in Clinical Medicine. Instructions for authors and procedures for submissions can be found on the *Journal*'s website at NEJM.org. At the discretion of the editor, images that are accepted for publication may appear in the print version of the *Journal*, the electronic version, or both.