

## CORRESPONDENCE

## Cochlear Implantation in Adults

**TO THE EDITOR:** In response to the review article by Carlson (April 16 issue)<sup>1</sup> describing cochlear implantation in adults, we totally agree that such surgery is an excellent method of rehabilitation for patients who cannot hear even with the use of hearing aids. However, we would like to emphasize the importance of postoperative mapping and speech therapy in such patients. Mapping refers to the customized programming of a cochlear implant according to the specific needs of its user, a process that allows the patient to hear sound most clearly from the electrodes of the cochlear implant array.<sup>2,3</sup> Mapping consists of various factors that include setting of threshold and comfort levels and adjustment of such measures as pulse width (control of stimulation rate), sensitivity (fine adjustment of the sound processor), and volume.<sup>4,5</sup> Speech therapy is an aural rehabilitation for patients who have received cochlear implants to improve their speech perception and to learn speaking and language skills. In addition, such therapy is essential for prelingually deaf adults and children.<sup>6</sup> In practice, patients may expect normal hearing, but it takes prolonged rehabilitation for the best speech recognition. Thus, postoperative rehabilitation, including mapping and speech therapy, is very important in order to obtain the best outcome after cochlear implant surgery.

Euyhyun Park, M.D.

Pyung Kon Thak, Ph.D.

Gi Jung Im, M.D., Ph.D.

Korea University College of Medicine  
Seoul, South Korea

logopas@korea.ac.kr

No potential conflict of interest relevant to this letter was reported.

1. Carlson ML. Cochlear implantation in adults. *N Engl J Med* 2020;382:1531-42.
2. Stakhovskaya O, Sridhar D, Bonham BH, Leake PA. Frequency map for the human cochlear spiral ganglion: implications for cochlear implants. *J Assoc Res Otolaryngol* 2007;8:220-33.
3. Eikelboom RH, Jayakody DM, Swanepoel DW, Chang S, Atlas MD. Validation of remote mapping of cochlear implants. *J Telemed Telecare* 2014;20:171-7.
4. Loizou PC, Poroy O, Dorman M. The effect of parametric variations of cochlear implant processors on speech understanding. *J Acoust Soc Am* 2000;108:790-802.
5. Bonnet RM, Boermans P-PBM, Avenarius OF, Briaire JJ, Frijns JHM. Effects of pulse width, pulse rate and paired electrode stimulation on psychophysical measures of dynamic range and speech recognition in cochlear implants. *Ear Hear* 2012;33:489-96.

6. Waters T. Speech therapy with cochlear implant wearers. *Br J Audiol* 1986;20:35-43.

DOI: 10.1056/NEJMc2017537

**THE AUTHOR REPLIES:** I am grateful to Park et al. for their commentary. With 15,000 mechanosensory cochlear hair cells and more than 35,000 spiral ganglion cells, the “normal” human auditory system can discriminate pitch variations of less than 5 Hz and differences in sound level as small as 1 dB. Given the inherent complexity of the inner ear, it is remarkable that many users of cochlear implants can decipher complex speech through broad stimulation of severely degraded auditory pathways using less than 10 functionally independent electrodes.<sup>1</sup> Thus, the brain has a remarkable capacity to fill in the gaps and adapt to a relatively crude afferent input. Nevertheless, this neuroplasticity requires time. Patients often do not have appreciable improvement in speech perception for several months after surgery, and auditory performance plateaus at approximately 1 to 3 years.<sup>2</sup> Furthermore, music appreciation and understanding speech in a noisy environment remain problematic for many experienced implant users. Ultimately, a person’s auditory outcome depends on a complex interplay between “bottom-up” auditory sensitivity to spectrotemporal cues (which is enhanced by device programming) and “top-down” linguistic and neurocognitive features that may be partially addressed through personalized intensive rehabilitation.<sup>3</sup>

Matthew L. Carlson, M.D.

Mayo Clinic  
Rochester, MN

carlson.matthew@mayo.edu

Since publication of his article, the author reports no further potential conflict of interest.

1. Riss D, Arnoldner C, Baumgartner W-D, Kaider A, Hamzavi J-S. A new fine structure speech coding strategy: speech perception at a reduced number of channels. *Otol Neurotol* 2008;29:784-8.
2. Cusumano C, Friedmann DR, Fang Y, Wang B, Roland JT Jr, Waltzman SB. Performance plateau in prelingually and postlingually deafened adult cochlear implant recipients. *Otol Neurotol* 2017;38:334-8.
3. Moberly AC, Bates C, Harris MS, Pisoni DB. The enigma of poor performance by adults with cochlear implants. *Otol Neurotol* 2016;37:1522-8.

DOI: 10.1056/NEJMc2017537

Correspondence Copyright © 2020 Massachusetts Medical Society.