

REVIEW ARTICLE

# Repair Schwann cell update: Adaptive reprogramming, EMT, and stemness in regenerating nerves

Kristjan R Jessen<sup>1</sup>  | Peter Arthur-Farraj<sup>2</sup> 

<sup>1</sup>Department of Cell and Developmental Biology, University College London, London, United Kingdom

<sup>2</sup>John Van Geest Centre for Brain Repair, Department of Clinical Neurosciences, University of Cambridge, Cambridge, United Kingdom

## Correspondence

Peter Arthur-Farraj, John Van Geest Centre for Brain Repair, Department of Clinical Neurosciences, University of Cambridge, Cambridge, United Kingdom.  
Email: pja47@cam.ac.uk

## Funding information

The work from the authors laboratory discussed in this article was supported by the Wellcome Trust (Programme Grant 074665 to K.R.J. and R.M. and a Clinical Research Career Development Fellowship 206634 to P.A-F), the Medical Research Council (Project Grant G0600967 to K.R.J. and R.M.), and the European Community (Grant HEALTH-F2-2008-201535 from FP7/2007-3013).

## Abstract

Schwann cells respond to nerve injury by cellular reprogramming that generates cells specialized for promoting regeneration and repair. These repair cells clear redundant myelin, attract macrophages, support survival of damaged neurons, encourage axonal growth, and guide axons back to their targets. There are interesting parallels between this response and that found in other tissues. At the cellular level, many other tissues also react to injury by cellular reprogramming, generating cells specialized to promote tissue homeostasis and repair. And at the molecular level, a common feature possessed by Schwann cells and many other cells is the injury-induced activation of genes associated with epithelial-mesenchymal transitions and stemness, differentiation states that are linked to cellular plasticity and that help injury-induced tissue remodeling. The number of signaling systems regulating Schwann cell plasticity is rapidly increasing. Importantly, this includes mechanisms that are crucial for the generation of functional repair Schwann cells and nerve regeneration, although they have no or a minor role elsewhere in the Schwann cell lineage. This encourages the view that selective tools can be developed to control these particular cells, amplify their repair supportive functions and prevent their deterioration. In this review, we discuss the emerging similarities between the injury response seen in nerves and in other tissues and survey the transcription factors, epigenetic mechanisms, and signaling cascades that control repair Schwann cells, with emphasis on systems that selectively regulate the Schwann cell injury response.

## KEYWORDS

adaptive reprogramming, c-Jun, epithelial-mesenchymal transition, nerve injury, nerve regeneration, repair cells, Schwann cells


## 1 | INTRODUCTION: THE CHANGING VIEW ON THE SCHWANN CELL INJURY RESPONSE

It has long been recognized that after injury to a peripheral nerve, Schwann cells undergo a radical change in identity by converting to denervated Schwann cells, which unlike the Schwann cells of uninjured nerves provide powerful support for regeneration (reviewed in Jessen & Mirsky, 2005, 2008, 2016; Chen, Yu, & Strickland, 2007; Zochodne, 2012; Glenn & Talbot, 2013; Scheib & Höke, 2013; Brosius Lutz & Barres, 2014; Faroni, Mobasseri, Kingham, & Reid, 2015; Jessen, Mirsky, & Lloyd, 2015; Cattin & Lloyd, 2016; Boerboom, Dion, Chariot, & Franzen, 2017). Although this Schwann cell injury response

is central to nerve repair, the nature of this process has been uncertain and disputed (Figure 1).

A recent contribution to this debate, namely the identification of partial epithelial-mesenchymal transition (EMT) in injured nerves (Arthur-Farraj et al., 2017; Clements et al., 2017), highlights how our understanding of the Schwann cell injury response has progressed by degrees, with each step providing new insights into this complex process.

Early studies on injured nerves focused on the obvious structural changes distal to injury, in particular the fragmentation for the myelin sheath following axonal death, rearrangement of Schwann cells into cellular columns, Bungner Bands, and invasion of macrophages, events



**FIGURE 1** Major cell components of regenerating nerves. The diagram shows the proximal and distal stumps and the bridge that connects them. (a) Regeneration unit, (b) blood vessel, (c) Schwann cells migrating from the distal stump, (d) macrophages and fibroblasts, and (e) Bunger bands. The basal lamina that covers the Schwann cells in the proximal stump and the Bunger bands in the distal stump is not shown. In crushed nerves, this basal lamina and surrounding connective tissue remain uninterrupted, allowing axons to regenerate within their original basal lamina tubes from the proximal nerve and along the distal stump. In cut nerves, axons accompanied by Schwann cells must traverse the bridge and successfully associate with a regeneration track for regeneration and target reinnervation to be possible. The length of the bridge depends on the extent of the injury and the type of surgical intervention

traditionally referred to as Wallerian degeneration (e.g., Lubinska, 1977; Stoll, Griffin, Li, & Trapp, 1989).

With the identification of molecular profile of myelin and Remak cells, and of their developmental ancestors in the Schwann cell lineage, it became clear that the generation of denervated cells after injury was accompanied by a striking downregulation of the molecules that characterized myelin cells. These denervated cells therefore shared one prominent feature, namely the absence of high levels of myelin gene expression, with immature Schwann cells of developing nerves prior to myelination. This reversal of myelin differentiation was instrumental in generating a view of the entire Schwann cell injury response essentially as a developmental regression to an earlier developmental stage, or dedifferentiation (e.g., Chen et al., 2007; Jessen & Mirsky, 2008).

The picture changed again when it was found that the generation of denervated cells fully capable of supporting repair depended not only on myelin dedifferentiation, but also on the activation of transcriptional mechanisms that were not significantly involved in Schwann cell development. Two transcriptional controls, c-Jun and chromatin modifications involving H3K27 demethylation, H3K27 deacetylation and H3K4 methylation, were found to be important for the normal execution of the Schwann cell injury response, although these mechanisms did not regulate neonatal Schwann cells or myelination (Arthur-Farraj et al., 2012; Hung, Sun, Keles, & Svaren, 2015; reviewed in Ma & Svaren, 2018). Additional control mechanisms that are selectively important in denervated cells have now been identified, including Merlin and STAT3 (Benito et al., 2017; Mindos et al., 2017). Denervated adult Schwann cells have also been shown to express *de novo* molecular markers that are low/absent during development, including Sonic Hedgehog (Shh), and Olig1 (Lu et al., 2000; Bosse et al., 2006; Arthur-Farraj et al., 2012; Fontana et al., 2012; Lin, Oksuz, Edward Hurley,

Wrabetz, & Awatramani, 2015; reviewed in Jessen & Mirsky, 2016). Again in contrast to developing cells, denervated cells express high levels of cytokines, recruit macrophages, and activate myelin autophagy to clear myelin (Gomez-Sanchez et al., 2015; reviewed in Martini, Fischer, López-Vales, & David, 2008; Rotshenker, 2011; Stratton & Shah, 2016). Denervated adult cells also adopt a striking elongated and often branched morphology, which is very different from that of Schwann cells in developing nerves, and which enables them to form regeneration tracks, which guide regrowing axons (Gomez-Sanchez et al., 2017).

In view of these substantive differences between developing and denervated Schwann cells, it is no longer useful to view the Schwann cell injury response simply as a developmental regression or dedifferentiation. Notably, most of the properties of denervated cells that are distinctive and differentiate them from developing cells, also make these cells exceptionally well suited for supporting nerve regeneration, leading to the suggestion that these cells are best understood as cells specialized for supporting this process, repair Schwann cells (reviewed in Jessen & Mirsky, 2016). This concept recognizes that the Schwann cell injury response represents a dual change: loss of myelin differentiation coupled with the activation of a set of features that support regeneration, a repair program. As outlined in the subsequent section, this adaptive response shows similarities to injury-induced cell conversions in other tissues, which are typically referred to as direct (or lineage) reprogramming. We have therefore referred to it as adaptive cellular reprogramming (Jessen, Mirsky, & Arthur-Farraj, 2015).

The most recent addition to the changing picture of the Schwann cell injury response, namely the finding that it includes upregulation of the genes associated with EMT and stemness, brings Schwann cells into line with several other systems in which injury also activates


EMT/stemness genes (Arthur-Farraj et al., 2017; Clements et al., 2017). This adds important components to our understanding of the repair program, and makes biological sense because EMT-like changes typically provide cells with increased motility and morphological flexibility. In Schwann cells, this would facilitate the extensive tissue remodeling that takes place when the nerve trunk distal to the injury converts into a collection of regeneration tracks, and when a tissue bridge forms across the injury site in severed nerves (Figures 1 and 2). Further increase in stemness generally shifts cells to a state associated with a change of identity. This is consistent with the view that injury reprograms Remak and myelin cells to adopt the distinct alternative identity of a repair cell.

We will now discuss the general concept of adaptive reprogramming and how this notion relates to the Schwann cell response to injury.

## 2 | ADAPTIVE CELLULAR REPROGRAMMING

Peripheral nerves provide a striking example of how mammalian tissues can regenerate and recover function after injury. Thus, some 3–4 weeks after crush of the sciatic nerve in rodents, which results in the destruction of axons and myelin and loss of both sensory and motor function in the hind limb, the nerve has regenerated, myelin sheaths have reformed and the limb is again fully functional.

The radical change in the differentiation state of myelin and Remak Schwann cells to a repair-supportive phenotype, which is a key mechanism behind this impressive regenerative capacity, shows



**FIGURE 2** Regeneration tracks (Bungner bands). Electron micrograph showing a transverse section through the distal stump of the mouse tibial nerve 4 weeks after transection (regeneration prevented). Several Bungner bands (example indicated at B) are visible, one of which shows a repair cell nucleus (N). The number of cell profiles differs considerably between the bands, each of which is surrounded by a basal lamina (examples arrowed) and collagen-rich endoneurial connective tissue. This represents the terrain along which axons grow during the often extended period of nerve repair in larger animals

interesting similarities with the cellular response to damage in a number of other tissues, including the pancreas, ear, liver eye, and skin. In common, these systems respond to injury, not by developmental regression, but by cellular reprogramming to new differentiation states, which restore tissue homeostasis.

Several tissues show this response, which works in one of two ways (reviewed in Jessen, Mirsky, & Arthur-Farraj, 2015). In some systems, the surviving cells convert to cells similar to those that were lost during injury. Such generation of replacement cells takes place for instance in the murine pancreatic isles. Following destruction of the insulin secreting  $\beta$ -cells, the remaining glucagon-secreting  $\alpha$ -cells replace the loss by converting to  $\beta$  cells in a process that combines the downregulation of glucagon expression with activation of insulin secretion. The new replacement cells are functional and capable of partially restoring glycemic control to the animal (reviewed in Chera & Herrera, 2016; Thorel et al., 2010). Formation of replacement cells is also seen the vestibular and cochlear epithelium of the ear. Here, damage to hair cells, involved in balance and hearing, triggers the direct conversion of neighboring glial-like supportive cells to new hair cells. (Bramhall, Shi, Arnold, Hochedlinger, & Edge, 2014; Cox et al., 2014; Forge, Li, & Nevill, 1998; Mizutani et al., 2013; Richardson & Atkinson, 2015). In the liver also, damage to biliary epithelial cells sparks the reprogramming of surrounding hepatocytes to form new biliary epithelial cells (reviewed in Eberhard & Tosh, 2008; Yanger et al., 2013). The classical non-mammalian example of this type of cell conversion is the transformation of pigmented epithelial cells of the dorsal iris to transparent lens cells, illustrating the combination of partial dedifferentiation (loss of the pigment phenotype), with activation of alternative differentiation (upregulation of lens cell features, such as crystalline expression) (Henry & Tsonis, 2010).

Illustration of the other form of adaptive reprogramming, the generation of repair cells, is provided by myofibroblasts in skin and heart injuries (Davis, Burr, Davis, Birnbaumer, & Molkenin, 2012; reviewed in Hinz et al., 2012). Myofibroblasts are absent from normal uninjured tissue. But after injury, myofibroblasts are generated from fibroblast and function to accelerate wound healing by secretion of extracellular matrix and restoring tension to the damaged tissue. Blocking the fibroblast–myofibroblast conversion results in defective skin and heart repair, demonstrating the adaptive nature of this reprogramming event.

In peripheral nerves, neurons respond to injury by activating gene programs that temporarily shift their function from that of cell–cell communication to that of regenerating axons, a response classically referred to as the signaling to growth mode switch or the cell body response (Allodi et al., 2012; Blesch et al., 2012; Doron-Mandel et al., 2015; reviewed in Fu & Gordon, 1997). In many ways, this response is comparable to the events discussed here as adaptive reprogramming.

As detailed in the following section, Schwann cells provide a further example of adaptive cellular reprogramming. In this case also, cells with a repair-supportive phenotype, which are normally absent, are generated directly from resident differentiated cells of the injured tissue (Gomez-Sanchez et al., 2017) and function to restore tissue homeostasis.



## 2.1 | The generation of repair Schwann cells

The Schwann cell injury response has been reviewed elsewhere (Boerboom et al., 2017; Chen et al., 2007; Jessen & Mirsky, 2008, 2016; Monk, Feltri, & Taveggia, 2015). Briefly, it has two principal components. One is the reversal of myelin differentiation. This involves transcriptional downregulation of the pro-myelin transcription factor *Egr2* (*Krox20*), the enzymes of cholesterol synthesis, and a number of structural and membrane associated myelin related proteins. At the same time, the cells reexpress molecules that were suppressed when immature Schwann cells started to myelinate during development, including *L1*, *NCAM*, *p75NTR*, and glial fibrillary acidic protein and many others.

The other component of the injury response is the sequential activation of a diverse features all of which support repair: (a) Upregulation of proteins that support neuronal survival and promote axonal regeneration, such as *GDNF*, *artemin*, *BDNF*, *NT3*, *NGF*, *VEGF*, *erythropoietin*, *pleiotrophin*, *p75NTR*, and *N-cadherin* (Brushart et al., 2013; Chen et al., 2007; Fontana et al., 2012; reviewed in Boyd & Gordon, 2003; Scheib & Höke, 2013; Wood & Mackinnon, 2015). (b) Activation of an innate immune response, comprising the upregulation of cytokines including *TNF $\alpha$* , *LIF* *Il-1 $\alpha$* , *Il-1 $\beta$* , *LIF*, and *MCP-1* (reviewed in Martini et al., 2008; Rotshenker, 2011). This recruits macrophages, which help Schwann cells to clear redundant myelin, stimulate the formation of blood vessels and axon growth (Barrette et al., 2008; Bauer, Kerr, & Patterson, 2007; Cafferty et al., 2001; Cattin et al., 2015; Hirota, Kiyama, Kishimoto, & Taga, 1996; Niemi et al., 2013; reviewed in Hirata & Kawabuchi, 2002). (c) Structural reorganization, generating extremely elongated, cells, which are about threefold longer than myelin and Remak cells. Lineage tracing has shown that these cells are generated directly from myelin and Remak cells, and that they can convert directly to myelin cells following regeneration (Gomez-Sanchez et al., 2017). These cells often carry long, parallel processes, which together with their length, allows repair cells to partly overlap and promotes the formation of the compact cellular columns (Bungner bands—Figures 1 and 2), which are obligatory regeneration tracks for regenerating axons (Gomez-Sanchez et al., 2017). (d) Activation of myelinophagy by which myelin cells initiate the breakdown of their own myelin sheath (Gomez-Sanchez et al., 2015; Suzuki, Lovera, Schmachtenberg, & Couve, 2015; Jang et al., 2016; Brosius Lutz et al., 2017).

The phenotype of repair Schwann cells, like other cells in the lineage, depends on environmental signals. Thus, as regeneration proceeds and repair cells become redundant, they obey axonal signals to return to the myelin and Remak phenotypes. The repair cells are therefore transient, produced on demand, and present as long as needed (reviewed in Jessen & Mirsky, 2016).

## 2.2 | Timing and activation of the repair program

The changes that constitute the Schwann cell injury response, and outlined above, do not all take place simultaneously. Rather, nerve injury starts the execution of a repair program, which is a temporal sequence of changes in cellular gene expression, structure, and function. Thus, expression of cytokines such as *Il-1 $\beta$*  and *TNF $\alpha$*  reaches

peak levels within 1 day after injury (reviewed in Rotshenker, 2011), myelin autophagy is maximally activated around Day 5 (Gomez-Sanchez et al., 2017), *GDNF* protein levels peak around Day 7, while *BDNF* protein peaks at 2–3 weeks. *c-Jun* is activated rapidly, but *c-Jun* protein levels continue to increase at least until 7–10 days after injury (J. Gomez-Sanchez, K. R. Jessen, and R. Mirsky unpublished observations), and full cellular elongation is not achieved until about four weeks after nerve injury or later (Gomez-Sanchez et al., 2017). The phenotype of the repair cell therefore undergoes an orderly series of changes, which collectively promote nerve regeneration.

The identification of the cell-extrinsic signals that initiate these changes after injury is an important area of future work. The first signs of the injury response can be detected surprisingly early (5–24 hr) after nerve cut. This includes the elevation of phospholipase A2 immunoreactivity, expression of mRNA for *LIF*, *c-Jun* and several other *AP1*-family members, and of *c-Jun*, *IL1* *Il-1 $\beta$*  and *TNF $\alpha$*  protein, phosphorylation of *ErbB2* neuregulin receptor, *p38* MAPK activation, and actin polymerization (Banner & Patterson, 1994; Kurek, Austin, Cheema, Bartlett, & Murphy, 1996; De et al., 2003; Parkinson et al., 2008; Jung et al., 2011; Yang et al., 2012; Guertin et al., 2005; Arthur-Farraj et al., 2017; reviewed in Martini et al., 2008; Rotshenker, 2011; Wong, Babetto, & Beirowski, 2017). These Schwann cell responses are seen before appreciable structural axonal degeneration and substantial macrophage influx. They are therefore likely to be triggered mainly by signals from transected/crushed axons. These signals remain largely unidentified, although the early *ErbB2* phosphorylation raises the possibility of neuregulin involvement. There is evidence that purines, which have a role in many other injury responses, play a part through activation of Schwann cell protein kinase C or *ERK1/2* (Negro et al., 2016; Xu et al., 2013). Neurone-derived hydrogen peroxide has also been implicated in early Schwann cell activation involving Annexin proteins (Duregotti et al., 2015; Negro et al., 2018), and induction of injury-related cytoskeletal changes (Gomez-Sanchez et al., 2017; Reynolds & Wolf, 1992; Son, Trachtenberg, & Thompson, 1996).


## 2.3 | Signals that control repair cells

Below is an outline of the molecular signaling that controls the generation and function of repair cells (Figure 3). In addition, we summarize studies that describe a nerve injury phenotype *in vivo* following conditional gene inactivation in Schwann cells (Table 1). Many of the factors which regulate the Schwann cell injury response also control other aspects of Schwann cell development and myelination. Except for re-myelination after injury, these other functions will not be discussed in detail, nor will signals that control Schwann cell proliferation.

### 2.3.1 | Transcription factors

The transcription factor *c-Jun* is a key regulator of the Schwann cell injury response. After injury, it is rapidly upregulated in Schwann cells where it negatively regulates the myelin program, and promotes expression of the repair program. This includes increase in trophic support for neurons, acceleration of myelin breakdown by autophagy, promotion of Schwann cell elongation, and formation of the bands of Bungner. As a result, regeneration of motor and sensory axons, and





**FIGURE 3** Signals that regulate the Schwann cell injury response and re-myelination. The diagram shows the repair Schwann cell in the context of other cells in the Schwann cell lineage, and indicates the major molecular systems implicated in any aspect of repair cell generation (left of double vertical arrows) and myelination/re-myelination (right of double vertical arrows).  $\Delta$ H3K27/H3K4: Loss of H3K27 trimethylation and gain of H3K4 methylation; TGF $\beta$ R: TGF $\beta$  receptors; NRG: Neuregulin-1; for other abbreviations and references see text. The developmental options of Schwann cell precursors are also indicated (for references see Jessen, Mirsky, & Lloyd, 2015)

the survival of sensory neurons and facial motor neurons are severely compromised if Schwann cell c-Jun is inactivated. Conversely, in situations where normal regeneration is impaired, such as after chronic denervation and in aging animals, enhanced expression of Schwann cell c-Jun promotes axonal regeneration (Parkinson et al., 2008; Arthur-Farraj et al., 2012; Fontana et al., 2012; Fazal et al., 2017; Wagstaff et al., 2017; Figlia, Gerber, & Suter, 2018). Interestingly, macrophage recruitment and Schwann cell numbers, at least at early time points after injury, are less affected in the absence of c-Jun than other aspects of the injury response (Arthur-Farraj et al., 2012). A microarray screening shows that c-Jun controls 172 genes, including directly regulating several neurotrophic factors such as GDNF, BDNF, Shh, and Artemin (Arthur-Farraj et al., 2012; Fontana et al., 2012). However, after nerve injury microarray and RNA-Seq analysis demonstrate at least 4,000–5,000 differentially expressed RNAs, showing that c-Jun only controls a subset of injury induced genes (Arthur-Farraj et al., 2012, 2017; Barrette, Calvo, Vallières, & Lacroix, 2010; Bosse, Hasenpusch-Theil, Küry, & Müller, 2006; Chang et al., 2013; Nagarajan, Le, Mahoney, Araki, & Milbrandt, 2002).

The extracellular signals that activate Schwann cell c-Jun after injury are obscure. Nevertheless, a number of intracellular pathways that potentially regulate c-Jun levels have been identified. This includes the Rac1-MKK7-JNK pathway and mTORC, which elevate c-Jun and have been implicated as upstream activators of c-Jun after nerve injury, and the cAMP pathway that suppresses c-Jun (Arthur-Farraj et al., 2011; Monje, Soto, Bacallao, & Wood, 2010; Normén et al., 2018; Shin et al., 2013). Histone deacetylase 2 (HDAC2), which is activated in injured nerves, also downregulates c-Jun (Brügger et al., 2017; reviewed in Jacob, 2017). The transcription factors Krox20 and Oct6 also antagonize c-Jun expression (Brügger et al., 2017; Parkinson et al., 2008). Since Krox20 is suppressed but Oct6 elevated after injury, these factors modulate c-Jun levels in opposing directions in damaged nerves. It has recently been shown that c-Jun can be

O-GlcNAcylated. Loss of this posttranslational modification results in inappropriate elevation of the c-Jun-dependent repair program and impaired remyelination after regeneration (Kim et al., 2018).

Sox2 as well as other negative regulators of myelination, including Id2 and Pax-3 are also induced in repair cells, and are likely to take part in suppressing myelin genes after injury although this has not yet been shown directly in vivo. (Doddrell et al., 2012; Florio et al., 2018; Kioussi, Gross, & Gruss, 1995; Le et al., 2005; revived in Jessen & Mirsky, 2008; Stolt & Wegner, 2016). Sox2 activation in Schwann cells has a role in the tissue bridge between the proximal and distal stumps where it mediates ephrin-B/EphB2 signaling between fibroblasts and Schwann cells, which promotes the clustering of migrating Schwann cells and ordered axon growth in the bridge (Parrinello et al., 2010). Enforced expression of Sox2 in vivo suppresses myelination both after injury and during development, but does not control the number of regenerated axons in injured nerves (Roberts et al., 2017).

The Notch pathway is activated in Schwann cells upon nerve injury and inactivation of Notch signaling or constitutive Notch activation specifically in Schwann cells, retards or accelerates demyelination, respectively (Woodhoo et al., 2009). Notch suppresses myelin genes, and other signals including the transcription factor Zeb2 and let-7miRNA, which promotes myelination and myelin gene expression, are thought to act in part by inhibiting Notch (Gökbuget et al., 2015; Pereira et al., 2010; Quintes et al., 2016; Wu et al., 2016). Enforced activation of Notch alone, even in uninjured nerves, is sufficient to induce myelin breakdown (Woodhoo et al., 2009). Whether the Notch-mediated promotion of myelin clearance after injury affects regeneration or whether Notch has broader effect on repair cells is yet to be studied.

Zeb2 levels are elevated in Schwann cells of injured nerves. This does not appear to regulate the expression of c-Jun and Sox2 at early time points after injury suggesting that Zeb2 is relatively unimportant for the initial generation of repair cells. However, during re-

**TABLE 1** Summary of in vivo phenotypes caused by genetic modifications in Schwann cells

Gene	Phenotype						Reference
	Mouse	Demyelination	SC Proliferation	Macrophage	Axon regeneration	Neuronal survival	
<i>c-Jun</i>	<i>POCre c-Jun f/f</i>	Delayed	Normal	Normal influx but reduced ability to degrade myelin after chronic cut	Delayed small and large fibre motoneuron target reinnervation	Reduced motor (facial) and sensory (sciatic)	Uninjured nerve develops normally in these mice Arthur-Farraj et al., 2012 Fontana et al., 2012
<i>Oct6</i>	<i>POCre c-Jun stop f/f</i>	Accelerated	Not assessed	Not assessed	Normal	Not assessed	Fazal et al., 2017
<i>Sox2</i>	<i>Oct6ΔSCE</i>	Mildly delayed	Not assessed	Not assessed	Slightly accelerated	Not assessed	Brügger et al., 2017
	<i>POCre Sox2IRESGFP Stop f/f</i>	Normal	Not assessed	Increased numbers in uninjured neve	Normal	Not assessed	Roberts, Dun, Dee, et al., 2017
<i>Stat3</i>	<i>POCre Stat3 f/f</i>	Normal	Normal	Normal	Normal	Normal	Benito et al., 2017
<i>Zeb2</i>	<i>PLPCreERT2 Zebf/f</i>	Normal	Normal	Normal	Normal	Not assessed	Increased Schwann cell death in chronic denervation and downregulation of repair program genes Quintes et al., 2016
<i>erbB2</i>	<i>PLPCreERT2 erbB2 f/f</i>	Not assessed	Normal	Not assessed	Not assessed	Not assessed	Not assessed
	<i>S100 rtTA: tetO-caErbB2</i>	Not assessed	Increased	Not assessed	Increased	Not assessed	Enhanced c-Jun expression in SC Atanasoski et al., 2006 Han et al., 2017
<i>GPR126</i>	<i>PLPCreERT2 Gpr126 f/f</i>	Not initially delayed but increased myelin debris in nerve at late timepoints	Not assessed	Reduced	Delayed	Not assessed	Reduced cytokine expression in injured nerves Mogha, Harty, et al., 2016
<i>Notch1</i>	<i>POCre CALSD-NICD stop f/f</i>	Accelerated	Not assessed	Not assessed	Not assessed	Not assessed	Not assessed
	<i>POCre RBPJ f/f</i>	Delayed	Not assessed	Not assessed	Not assessed	Not assessed	Not assessed
<i>NRG1</i>	<i>SLICK A Cre Nrg1 f/f</i>	Not assessed	Not assessed	Not assessed	Delayed	Not assessed	Removal of NRG1 in neuronal subset Fricke et al., 2011
	<i>CAG CreER Nrg1 f/f</i>	Normal	Normal	Normal	Delayed	Not assessed	Removal of Nrg1 from SC and neurons with tamoxifen Fricke et al., 2013

Woodhoo et al., 2009



TABLE 1 (Continued)

Gene	Mouse	Phenotype							Comments	Reference
		Demyelination	SC Proliferation	Macrophage	Axon regeneration	Neuronal survival	Remyelination			
	<i>DhhCre Nrg1 f/f</i>	Not assessed	Not assessed	Not assessed	Not assessed	Not assessed	Impaired		Stassart et al., 2013	
<i>TGFβRII</i>	<i>POCre TGFβRII f/f</i>	Normal	Normal	Normal	Not assessed	Not assessed	Not assessed	Reduced SC migration into bridge	D'Antonio et al., 2006	
	<i>POCre TGFβRII f/f</i>	Normal	Normal	Not assessed	Delayed at bridge region But normal longterm	Not assessed	Normal		Clements et al., 2017	
<i>ATG7</i>	<i>POCre ATG7 f/f</i>	Delayed	Not assessed	Not assessed	Not assessed	Not assessed	Not assessed		Gomez-Sanchez et al., 2015	
<i>CEMIP</i>	<i>PLPCreERT CEMIP f/f</i>	Delayed	Not assessed	Not assessed	Normal functional recovery	Not assessed	Increased thickness		Boerboom et al., 2017	
<i>MEK1</i>	<i>PLPCreER MEK1DD stop f/f</i>	Not induced in uninjured nerve after 6 weeks Accelerated after cut	Not assessed	Not induced in uninjured nerve Increased influx after cut	Reduced small fibre regeneration and unmyelinated axon number	Not assessed	Normal thickness but reduced intermodal length and cajal bands	Constitutive MEK1 expression in SC with tamoxifen	Cervellini et al., 2018	
	<i>CNPCre MEK1DD stop f/f</i>	Induced in uninjured nerve at 8 months	Not assessed	BNB breakdown and influx in uninjured nerve	Not assessed	Not assessed	Not assessed	Increased ECM deposition after injury	Ishii et al., 2013	
<i>Merlin/NF2</i>	<i>POCre NF2 f/f</i>	Normal	Increased	Increased influx	Very reduced myelinated and unmyelinated axons	Not assessed	Impaired	Phenotype rescued by YAP SC knockout	Mindos et al., 2017	
	<i>PO-NF2Δ39-121</i>	Not assessed	Not assessed	Not assessed	Delayed with reduced myelinated axons	Not assessed	Impaired	Dominant negative merlin Greater ECM production	Truong et al., 2018	
<i>P38MAPKα</i>	<i>POCre P38α f/f</i>	Delayed	Normal	Normal	Normal	Not assessed	Normal		Roberts, Dun, Doddrell, et al., 2017	
<i>Raf</i>	<i>POCre RafTR</i>	Induced in uninjured nerve with tamoxifen	Not assessed	BNB breakdown and influx in uninjured nerve with tamoxifen	Not assessed	Not assessed	Not assessed	Constitutive Raf expression in SC with tamoxifen	Napoli et al., 2012	
<i>Dicer</i>	<i>PLPCreERT Dicer f/f</i>	Normal	Normal	Not assessed	Not assessed	Not assessed	Impaired		Viader et al., 2011	
<i>Hdac1/2</i>	<i>POCreERT2 Hdac1 f/f Hdac2 f/f</i>	Mildly accelerated	Increased	Not assessed	Increased	Not assessed	Impaired		Brügger et al., 2017	
<i>Hdac 3</i>	<i>PLPCreERT Hdac3 f/f</i>	Not assessed	Not assessed	Not assessed	Not assessed	Not assessed	Increased thickness		He et al., 2018	
<i>O-GlcNAc transferase</i>	<i>POCre OGT f/f</i>	Normal	Not assessed	Not assessed	Normal	Not assessed	Severely Impaired	OGT inhibits <i>c-Jun</i> activity	Kim et al., 2018	

SC = Schwann cell; BNB = blood-nerve barrier.



myelination, Zeb2 contributes to the suppression of negative regulators such as Sox2 and Id2 and the Notch pathway effector Hey2, and is therefore needed for normal generation of myelin in regenerating nerves, a function that mimics the role of this transcription factor during developmental myelination (Quintes et al., 2016; Wu et al., 2016).

Importantly, genome wide mapping of active enhancers in injured rat nerves with acetylated histone H3K27 CHIP-Seq found that the most abundant enriched motifs, in addition to c-Jun/AP-1, were for Runx2 and the Ets family of transcription factors (Hung et al., 2015). Ets transcription factors have previously been implicated in the support of Schwann cell survival after injury (Parkinson, Langner, Sharghi Namini, Jessen, & Mirsky, 2002). It will be important to define the role of these proteins in the Schwann cell injury response.

Axons take a long time, up to years in humans, to reach their targets after injury. Even for very distal injuries such as in the wrist, axons can take months to reinnervate the finger tips. For successful regeneration therefore, repair cell generation needs to be followed by long-term maintenance of the repair cell population. Because failure of repair cell maintenance is thought to be a significant reason for poor regeneration outcomes in humans, it is important to identify transcription factors and other signals that sustain repair cells. The first known transcription factor with this function is STAT3. This protein is not required for the initial generation of repair Schwann cells, but during chronic denervation, STAT3 is important both for the long-term survival of these cells and for promoting the expression of c-Jun and other repair program genes (Benito et al., 2017).

### 2.3.2 | Intracellular signaling pathways

Although myelin breakdown is an important component in the conversion of myelin cells to repair cells, the mechanism by which Schwann cells get rid of their redundant myelin sheaths has long been debated. However, a Schwann cell specific knockout of ATG7, a key protein in the autophagy process has now demonstrated that Schwann cells digest their own myelin through a process termed, myelinophagy, a type of macro-autophagy (Brosius Lutz et al., 2017; Gomez-Sanchez et al., 2015; Jang et al., 2016). In addition to myelinophagy, Schwann cells also phagocytose smaller amounts of myelin debris via TAM receptor phagocytosis (Brosius Lutz et al., 2017). Interestingly, ATG7 null Schwann cells demonstrated impaired upregulation of other repair program genes such as *GDNF*, *Olig1* and *Shh*, suggesting that disrupting myelinophagy impairs generation of the repair cell phenotype (Gomez-Sanchez et al., 2015). It remains to be tested, whether delaying myelinophagy ultimately affects axonal regeneration and functional nerve repair.

Another early function of repair Schwann cells, in addition to myelinophagy, is to recruit macrophages into injured nerve. These cells make an important contribution to myelin clearance by phagocytosing myelin debris, in addition to stimulating vascularization and likely promoting axonal regrowth by direct signaling to neurons (Barrette et al., 2008; Cafferty et al., 2001; Cattin et al., 2015; Hirota et al., 1996; Martini et al., 2008; reviewed in Bauer et al., 2007). Schwann cell recruitment of macrophages is under strong regulation by the Raf-MEK-ERK mitogen-activated protein kinase (MAPK) pathway. ERK1/2 phosphorylation is rapidly induced in Schwann cells after

injury, and implicated in the upregulation of the major macrophage recruitment signal MCP-1 (Fischer, Weishaupt, Troppmair, & Martin, 2008; Sheu, Kulhanek, & Eckenstein, 2000). Pharmacological inhibition of ERK1/2 activation, or genetic inactivation of CEMIP, a protein that stimulates ERK1/2 activity, delays myelin clearance after injury (Harrisingh et al., 2004; Napoli et al., 2012; reviewed in Boerboom et al., 2017). Conversely, strong activation of ERK1/2, whether by constitutively active Raf or MEK1, promotes macrophage recruitment, myelin breakdown and suppression of myelin genes. This is seen even in uninjured nerves in experiments using Raf (Napoli et al., 2012), but only after injury in studies using a constitutively active MEK1 variant to stimulate ERK1/2 (Cervellini et al., 2018). It still remains controversial whether Raf-MEK-ERK1/2 signaling has a role in activating c-Jun in Schwann cells (Monje et al., 2010; Shin et al., 2013; Syed et al., 2010).

The function of ERK1/2 in Schwann cells is, however complex, and not restricted to promoting the injury response, because ERK1/2 is also essential for developmental myelination (Newbern et al., 2011). In agreement with that, enhanced ERK1/2 activation in Schwann cells during early development results in increased myelin thickness (Ishii et al., 2013; Sheean et al., 2014). Notably, during re-myelination of regenerating nerves, enhanced Schwann cell ERK1/2 activation has a different outcome. In this case, there is no effect on myelin thickness, but myelin compaction, Cajal bands, internodal length and axonal regeneration are compromised (Cervellini et al., 2018). This indicates that repair cells differ from immature Schwann cells with respect to the intracellular signaling pathways that control myelination. The role of neuregulin in myelination also differs between these two Schwann cell types (see below).

The c-Jun N terminal kinase (JNK) pathway is rapidly activated within minutes to hours after nerve injury, although c-Jun protein levels continue to rise for several days after injury (De Felipe & Hunt, 1994; Parkinson et al., 2008; Yang et al., 2012; J. Gomez-Sanchez, K. R. Jessen, and R. Mirsky unpublished). While pharmacological experiments indicate that the Rac1-MKK7-JNK pathway is important for c-Jun upregulation (Monje et al., 2010; Shin et al., 2013), the requirement for JNK and c-Jun phosphorylation after nerve injury has not yet been tested through genetic inactivation. Although c-Jun is a major phosphorylation target of JNK, JNK phosphorylates a number of other proteins, and at least in developing Schwann cells, c-Jun suppresses myelin genes without phosphorylation (Parkinson et al., 2008). The relationship between JNK and the functions of Schwann cell c-Jun in injured nerves is therefore unclear.

The above findings suggest that the Raf-MEK-ERK1/2 and the Rac1-MKK7-JNK signaling pathways carry out distinct functions in repair cells (reviewed in Lee, Shin, & Park, 2014). The Raf-MEK-ERK1/2 pathway, is particularly implicated in cytokine expression, macrophage recruitment and myelin breakdown, while its role in c-Jun activation is not clear. The Rac1-MKK7-JNK pathway activates c-Jun and regeneration-associated genes, including *GDNF* and *P75NTR*, downregulates the pro-myelin transcription factor *Krox20*, and promotes myelinophagy. In line with this, there does not appear to be significant overlap in the genes, particularly cytokines, dysregulated by Raf activation and those by c-Jun inactivation in Schwann cells (Arthur-Farraj et al., 2012; Napoli et al., 2012).



Although the p38 mitogen-activated protein kinase (p38MAPK) pathway is strongly and rapidly activated in Schwann cells after injury, a Schwann cell specific knockout of the major p38 isoform, p38 $\alpha$ , which abolished p38MAPK activity after nerve injury, demonstrated only slightly slower clearance of myelin proteins, and no effect on macrophage influx, axonal regeneration or functional recovery (Roberts, Dun, Dee, et al., 2017).

Merlin is a tumor suppressor protein which has a significant role in repair Schwann cells (Hilton & Hanemann, 2014; Mindos et al., 2017). In mice with Schwann cell-specific knockout of Merlin, axonal regeneration is severely impaired and there is noticeable loss of regenerated large myelinated fibers. Re-myelination is suppressed and there is a large expansion in endoneurial size, likely due to increased extracellular matrix production. The hippo pathway effector YAP is overexpressed in Merlin null nerves, and genetic inactivation of YAP almost completely restores the regenerated nerve to normal (Mindos et al., 2017). A comparable, albeit much milder, regeneration phenotype is seen following overexpression of a dominant negative Merlin construct in Schwann cells, possibly due to incomplete inhibition (Truong et al., 2018). The molecular mechanism of how Merlin regulates the phenotype of repair Schwann cells and axonal regeneration is unclear, although both c-Jun and ERK1/2 activation are dysregulated in Merlin null Schwann cells (Mindos et al., 2017).

The mTORC1 pathway is activated in Schwann cells by injury. This is required for normal c-Jun activation likely because TORC1 promotes c-Jun translation. Genetic inactivation of the TORC1 pathway results in subdued activation of c-Jun and other repair cell genes, including *Olig1*, *Shh*, and *Runx2*. While macrophage recruitment is unaffected, the capacity of repair cells to clear myelin is impaired and re-myelination is delayed (Normén et al., 2018).

### 2.3.3 | Extracellular ligands, receptors, and cytokines

Among the growth factors and extracellular signaling molecules implicated in regulating the phenotype of repair Schwann cells, the neuregulin-1 signaling pathway is probably the most studied. Neuregulin-1 is dispensable for myelin maintenance in adult uninjured nerves (Atanasoski et al., 2006; Birchmeier & Nave, 2008). Although Schwann cells elevate expression of both ErbB2/3 receptors and neuregulin-1 I/II isoforms after injury, neuregulin-1 appears not to be involved in injury-induced Schwann cell proliferation, macrophage recruitment, and myelin breakdown (Atanasoski et al., 2006; Carroll, Miller, Frohnert, Kim, & Corbett, 1997; Fricker et al., 2013; Kwon et al., 1997; Ronchi et al., 2013; Stassart et al., 2013). Nerves with selective inactivation of neuregulin-1 in Schwann cells also re-myelinate after injury, although the remyelination rate is significantly impaired (Stassart et al., 2013). Even after removal of neuregulin-1 from both Schwann cells and axons, re-myelination is eventually normal after a substantial delay (Fricker et al., 2013; Stassart et al., 2013). Axonal regeneration is also slower in mice without neuregulin-1 in both axons and Schwann cells, suggesting that endogenous neuregulin signaling through ErbB2/3 receptors in repair cells promotes the repair phenotype and the capacity of these cells to support axon growth (Fricker et al., 2011, 2013). Direct evidence for this mechanism of action is missing, however. Nevertheless, artificially enhancing

neuregulin signaling might serve as a tool for promoting nerve repair, because enforced Schwann cell ErbB2 expression, and exogenously applied neuregulin increase axonal regeneration in vivo (Han et al., 2017; Gambarotta, Fregnan, Gnani, & Perroteau, 2013). Another factor that also potentially binds to ErbB receptors, betacellulin, is upregulated in Schwann cells after injury, but the involvement of betacellulin in injured nerves is unclear (Valli  res et al., 2017).

The G-protein coupled receptor GPR126 in Schwann cells is essential for both developmental myelination and re-myelination (Mogha, D'Rozario, & Monk, 2016; Mogha, Harty, et al., 2016). This protein is not required for the maintenance of myelin in uninjured nerves, but inactivation of Gpr126 in repair Schwann cells results in reduced axonal regeneration, reduced expression of TNF and a restricted group of chemokines, and impaired recruitment of macrophages, cells which potentially promote axonal regeneration (Barrette et al., 2008). In contrast, Gpr126 is dispensable for injury-induced c-Jun elevation in Schwann cells, Schwann cell proliferation, and myelin clearance (Mogha, Harty, et al., 2016). In developing nerves, GPR126 binds to laminin 211 to elevate cyclic AMP levels and promote myelination (Monk et al., 2009; Petersen et al., 2015). After nerve injury, the identity of the GPR126 ligand is unclear, though it may be laminin since removal of *laminin*  $\gamma$ 1 in Schwann cells impairs axon regeneration (reviewed in Chen & Strickland, 2003).

Disruption of collagen post-translational modifications through inactivation of *lysyl-hydroxylase 3 (lh3)* in zebrafish leads to selective mis-targeting of dorsal branch motoneurons into the ventral branch after laser axotomy. This is completely rescued through re-expression of *lh3* solely in Schwann cells. Interestingly ventral branch motoneurons regenerate normally suggesting that subsets of Schwann cells are responsible for target selective re-innervation after nerve injury (Isaacman-Beck et al., 2015).

Full knockouts of Toll-like receptor (TLR) 2 and 4 or their downstream effector MyD88 significantly impair early induction of cytokines, including MCP-1, IL-1 $\beta$  in injured nerves, and retard macrophage influx and myelin clearance. Conditional inactivation of TLRs is, however, required to investigate their specific role in repair Schwann cells (Boivin et al., 2007).

TGF- $\beta$ 1 is upregulated after nerve injury in Schwann cells and endoneurial fibroblasts and is also expressed by macrophages (Scherer, Kamholz, & Jakowlew, 1993; Sulaiman & Nguyen, 2016). In vitro, TGF- $\beta$ 1 shifts Schwann cells from a myelin toward a repair phenotype, since it suppresses myelin proteins and upregulates the adhesion molecules L1 and NCAM, both of which are elevated after nerve injury (Morgan, Jessen, & Mirsky, 1994; Stewart et al., 1995). TGF- $\beta$ 1 treatment of Schwann cells is also reported to improve axonal regeneration in vivo (Sulaiman & Nguyen, 2016). Removal of the TGF- $\beta$  receptor II in Schwann cells shows that TGF- $\beta$  has a role specifically at the site of nerve injury for promoting Schwann cell migration from the proximal stump, across a bridge region to reconnect with the distal stump. Disruption of this Schwann cell migration leads to mis-targeting of axons at the injury site, although long-term axonal regeneration and functional recovery were normal in the absence of Schwann cell TGF- $\beta$  signaling (Cattin et al., 2015; Clements et al., 2017; Parrinello et al., 2010).



### 2.3.4 | Epigenetic processes

Several epigenetic processes regulate the repair Schwann cell phenotype (Ma & Svaren, 2018; reviewed in Jacob, 2017). Injury sets in train chromatin remodeling involving, first, the loss of the active histone mark H3K27 acetylation on enhancers of a number of myelin genes. Second, a proportion of injury induced genes (around 20%), which are normally marked by the polycomb PRC2 complex with the repressive histone mark H3K27 trimethylation at enhancer regions, are demethylated. Simultaneously, they gain the active H3K4 methylation promoter mark and their expression is derepressed. It is likely that the histone demethylase JMJD3, which is upregulated by injury, mediates the demethylation of histone H3K27 in repair Schwann cells, because inhibition of this enzyme in nerve explants blocks upregulation of a number of repair program genes such as, *Fgf5*, *Shh*, and *Runx2* (Gomez-Sanchez et al., 2013; Hung et al., 2015; Ma et al., 2015; reviewed in Ma & Svaren 2016). JMJD3 also mediates promoter H3K27 demethylation of *p19Arf* and *p16Ink4a* to prevent Schwann cell over-proliferation (Gomez-Sanchez et al., 2013).

Histone deacetylase 1 and 2 (HDAC1/2) are upregulated in Schwann cells after injury. They suppress c-Jun and delay the conversion to repair cells. Accordingly, axonal regeneration is faster if Schwann cell HDAC1/2 is inactivated, but re-myelination is impaired (Brügger et al., 2017). HDAC4 also suppresses c-Jun and elevates myelin gene expression (Gomis-Coloma et al., 2018). In contrast, HDAC3 upregulates negative regulators of myelination, including *Id2* and *Sox2*, and the Notch and Hippo signaling pathways. Myelination is therefore enhanced if Schwann cell HDAC3 is inactivated (He et al., 2018).

MicroRNAs (miRNAs) are 21–24 nucleotide regulatory RNAs. After nerve injury, there is a significant change in expression of hundreds of miRNAs (Adilakshmi, Sudol, & Tapinos, 2012; Arthur-Farraj et al., 2017; Viader et al., 2011). Disruption of miRNA processing through deletion of the enzyme DICER impairs myelination during development and after injury (Pereira et al., 2010; Viader et al., 2011). Deletion of another component of miRNA processing, *Dgcr8*, similarly causes defective myelination, but results also in upregulation of a number of genes that are upregulated after injury, including *Sox2*, and repair program genes, such as *Shh*, *Gdnf*, *c-Jun*, and *Olig1* (Lin et al., 2015). MiR-9 in particular has been implicated in negative regulation of the injury response (Zhou et al., 2014). Taken together, this suggests that miRNAs repress the repair program and promote myelin gene expression. Interestingly, c-Jun controls expression of a number of miRNAs, including miR21 and miR34 (Arthur-Farraj et al., 2017).

Two other epigenetic mechanisms that may have a role in repair Schwann cells are long non-coding RNAs (lncRNAs) and DNA methylation. lncRNAs, are RNA molecules greater than 200 base pairs in length with no coding potential. There is a significant change in lncRNA expression after nerve injury, but the function of these molecules in repair Schwann cells is currently unknown (Arthur-Farraj et al., 2017). In contrast, there are only limited changes in genome-wide CpG DNA methylation after nerve injury, which argues against global regulation of gene expression through this mechanism in nerve repair (Arthur-Farraj et al., 2017). DNA methylation has been shown

to regulate developmental myelination but it is not known whether it also regulates re-myelination after injury (Varela-Rey et al., 2014).

### 2.4 | Distinctive control mechanisms in repair cells

In a number of cases, the molecular signals that control the Schwann cell injury response are not important for Schwann cell development, which is consistent with the view that injury generates a distinct repair-supportive Schwann cell. This includes c-Jun and Merlin. While these proteins are essential for the generation of functional repair cells as discussed above, c-Jun does not have a significant role in early Schwann cells, and Schwann cell development shows only minor impairments in mice without Merlin (Arthur-Farraj et al., 2012; Mindos et al., 2017). Similarly, developmental myelination is not controlled by the chromatin cross talk that in repair cells promotes silencing of myelin genes and activation of injury genes, and involves H3K27 demethylation coupled with H3K27 acetylation and H3K4 methylation (Hung et al., 2015; reviewed Ma & Svaren, 2018). STAT3 likewise is not involved in Schwann cell development although this factor is important for long-term repair cell maintenance (Benito et al., 2017). Furthermore, the proliferation of adult Schwann cells that is triggered by injury and associated with the generation of repair cells requires cyclin D1, although this protein is not involved in the proliferation of developing Schwann cells (Kim et al., 2000). Other differences between repair and developing cells concern neuregulin-1 and ERK1/2 as outlined earlier. Thus repair cells, but not developing Schwann cells, activate an autocrine neuregulin signaling loop that supports re-myelination, while axon-associated neuregulin-1 acts only as a timer for repair cell re-myelination, although it is essential for developmental myelination (Fricker et al., 2013; Stassart et al., 2013). Furthermore, enhanced ERK1/2 increases myelin thickness during development, but not during re-myelination (Cervellini et al., 2018; Ishii et al., 2013).

These data show that the reprogramming of myelin and Remak cells to repair cells involves dedicated injury-associated control systems that are not important in developing or undamaged, normal nerves. Interestingly, Schwann cell reprogramming has now been found to involve another set of gene changes that are not connected to Schwann cell development, namely that associated with EMT. EMT is also associated with tissue damage in many other systems. We will now outline the general association between EMT and injury, and the role of EMT in the Schwann cell injury response.

### 2.5 | Activation of EMT and stemness genes after tissue injury

EMT represents a process of cellular reprogramming that was initially conceived as a transition between two states, involving the wholesale conversion of epithelial cells to a motile mesenchymal phenotype (Forte et al., 2017; Kim et al., 2017; reviewed in Yang & Weinberg, 2008; Skrypek, Goossens, De Smedt, Vandamme, & Berx, 2017; Thiery, Acloque, Huang, & Nieto, 2009). EMT has been intensively studied in development and cancer, and is now known to be controlled by multiple signals, including secreted signals such as the TGF $\beta$  family, transcription factors, including the Snail, Slug, Zeb, and Twist families and others, in addition to miRNAs, for example, the miR-200

family. EMT is characteristically associated with downregulation of molecules that promote cell-cell adhesion, increased capacity for migration and matrix invasion, increased morphological plasticity, including loss of cell polarity, and proliferation. Recently, three important developments have modified the classical view of EMT.

First, it has become clear that rather than an all-or-none response, EMT-like changes are often graded, or partial. Evidence for partial EMT comes from developing systems, cancer, fibrosis and injury (Nieto, Huang, Jackson, & Thiery, 2016; reviewed in Grigore, Jolly, Jia, Farach-Carson, & Levine, 2016). This work has demonstrated the existence of a heterogeneous group of intermediate cellular states, which share both epithelial and mesenchymal features, showing a graded decrease in cell polarity and cell-cell adhesion often without resulting in frank cellular dispersion or loss of lineage identity.

Second, partial EMT has now been established as a normal physiological response to tissue injury, serving as a component of the healing mechanism. This is unsurprising because many of the features originally identified as characteristic of EMT in developing systems and listed above are also important for an effective response of adult tissues to injury. Injury-induced EMT occurs in many different systems (Nieto et al., 2016; reviewed in Weber, Li, Wai, & Kuo, 2012; Shaw & Martin 2016), in addition to peripheral nerve, as discussed further below. This includes, zebrafish heart, basal cells in the airway, ovarian epithelium, CNS meninges, skin wounds, and kidney tubules. In skin wounds, for instance, partial EMT is triggered in the cells at the wound boundary. This involves the downregulation of adherens and tight junctions, which promotes healing by allowing previously immobile epithelial cells to move and re-epithelialize the damaged area, in a process regulated by the classical EMT driver Slug (Hudson et al., 2009; Nunan et al., 2015; reviewed in Shaw & Martin 2016). In the kidney, obstruction or chemical injury induces partial EMT in the epithelial cells of the renal tubules, which also depends on Slug activation. The cells lose morphological differentiation and polarity markers, and express cytokines to attract macrophages, yet remain in the wall of the renal tubule and retain contact with the basal lamina they contacted before the injury (Grande et al., 2015; Lovisa et al., 2015). This response is reminiscent of the reprogramming events that generate repair Schwann cell in the distal nerve stump after injury, as discussed in a previous section.

Third, activation of EMT in injury and many other situations is frequently associated with activation of genes typical of stem cell states, and increased stemness (Liao & Yang, 2017; reviewed in Fabregat, Malfettone, & Soukupova, 2016). Stemness is associated with loosened phenotypic restrictions and enhanced plasticity. Therefore, the strong correlations between stemness and EMT, in particular intermediate or partial EMT (Forte et al., 2017; Nieto et al., 2016), is in line with the notion that EMT represents cellular reprogramming and a change of differentiation state. The EMT/stemness association is shown for instance by the findings that typical inducers of EMT, such as Snail, Slug, or TGF $\beta$ , also activate stem cell properties (Mani et al., 2008; Morel et al., 2008). A stem cell-like phenotype is also induced by other EMT inducers, including Zeb and Twist (reviewed in Fabregat et al., 2016). In line with this, invasive tumor cells express both EMT and stemness characteristics (reviewed in Jordan, Johnson, & Abell, 2011; Rhim et al., 2012).

Although the implications of the intriguing association between EMT and stemness have been the subject of much speculation, particularly in cancer, the significance may be clearer in the context of tissue injury. This is because, in many cases, the injury response requires not only the classical EMT attributes of motility, morphological flexibility, and proliferation, but also enhanced plasticity as surviving cells change differentiation states to acquire phenotypes adapted to promote healing, as discussed in the previous section on adaptive reprogramming. The combined activation of EMT/stemness programs in injured tissues therefore appears strongly adaptive and biologically useful. As discussed below, this mechanism has now been established in peripheral nerves.

## 2.6 | Repair Schwann cells undergo partial EMT

Two recent papers show the involvement of an EMT-like process in Schwann cells of injured nerves (Arthur-Farraj et al., 2017; Clements et al., 2017). RNA-Seq performed on 7-day injured whole nerves demonstrates enrichment for EMT mRNAs and miRNAs. This involves downregulation of RNAs associated with mesenchymal-epithelial transition, including *E-cadherin*, *Wt1*, *Fgf1*, *Ndr1*, *mir30*, *mir33*, and *mir137*, and upregulation of EMT associated RNAs, including *Tgf- $\beta$ 1*, *Met*, *Hmga2*, *mir21*, *mir221*, and *mir222*. *c-Jun* target genes are also enriched for an EMT signature (Arthur-Farraj et al., 2017). Similarly, a study involving FACs sorting of td-tomato labeled Schwann cells from the tissue bridge and from the distal stump of injured nerves shows enrichment for EMT genes (Clements et al., 2017). In line with the close relationship between EMT and stemness (Mani et al., 2008; Morel et al., 2008), this study also showed activation of the Myc stemness and Core pluripotency modules and suppression of polycomb-related factors.

In the distal stump, tissue remodeling involves converting myelin and Remak cells into repair cells in Bungner bands (reviewed in Jessen & Mirsky, 2016), while in the bridge it involves the formation of new tissue by migrating Schwann cells, together with fibroblasts and immune cells (reviewed in Cattin & Lloyd, 2016). Interestingly, EMT/stemness changes in Schwann cells were even more prominent in the bridge, where they were associated with localized signaling by TGF $\beta$ , a typical inducer of EMT and stem cell properties (Clements et al., 2017).

## 3 | CONCLUSIONS

Recent work has underlined the complexity of the Schwann cell injury response, and established the involvement of novel epigenetic mechanisms and morphogenetic transformations in this process. Significant issues have also been brought into focus. One of these is the close similarity between the way nerves and other tissues respond to injury. For instance, Schwann cells activate EMT/stemness-like programs during injury-induced tissue remodeling, as seen after damage in many other systems. Furthermore, similar to cells in a number of other tissues, Schwann cells adopt an alternative differentiation state after injury, which is adapted to meet the particular needs that arise after damage. On the other hand, the distinctive nature of the Schwann cell

injury response has also emerged, since it is now clear that it is controlled by molecular mechanisms a number of which have no or little role elsewhere in the Schwann cell lineage. The observation that repair Schwann cells are regulated by dedicated mechanisms, and the large number of novel signals that have recently been implicated in this process suggest that pharmacological or genetic tools can be developed to manipulate these particular cells. This is a useful development from the translation point of view, because there is a strong clinical need to learn how to amplify the repair-supportive functions of denervated Schwann cells, and how to prevent their deterioration in older age and during the prolonged periods required for axonal regeneration in human nerves.

## ACKNOWLEDGMENTS

The authors are grateful to R. Mirsky for constructive criticism of the manuscript and L. Wagstaff, J. Gomez-Sanchez, and R. Mirsky for Figure 2.

## ORCID

Kristjan R Jessen  <https://orcid.org/0000-0002-7883-1221>

Peter Arthur-Farraj  <https://orcid.org/0000-0002-1239-9392>

## REFERENCES

- Adilakshmi, T., Sudol, I., & Tapinos, N. (2012). Combinatorial action of miRNAs regulates transcriptional and post-transcriptional gene silencing following in vivo PNS injury. *PLoS One*, 7, e39674. <https://doi.org/10.1371/journal.pone.0039674>
- Allodi, I., Udina, E., & Navarro, X. (2012). Specificity of peripheral nerve regeneration: Interactions at the axon level. *Progress in Neurobiology*, 98, 16–37. <https://doi.org/10.1016/j.pneurobio.2012.05.005>
- Arthur-Farraj, P. J., Latouche, M., Wilton, D. K., Quintes, S., Chabrol, E., Banerjee, A., ... Jessen, K. R. (2012). c-Jun reprograms Schwann cells of injured nerves to generate a repair cell essential for regeneration. *Neuron*, 75, 633–647. <https://doi.org/10.1016/j.neuron.2012.06.021>
- Arthur-Farraj, P. J., Morgan, C. C., Adamowicz, M., Gomez-Sanchez, J. A., Fazal, S. V., Beucher, A., ... Aitman, T. J. (2017). Changes in the coding and non-coding transcriptome and DNA methylome that define the Schwann cell repair phenotype after nerve injury. *Cell Reports*, 20, 2719–2734. <https://doi.org/10.1016/j.celrep.2017.08.064>
- Arthur-Farraj, P., Wanek, K., Hantke, J., Davis, C. M., Jayakar, A., Parkinson, D. B., ... Jessen, K. R. (2011). Mouse Schwann cells need both NRG1 and cyclic AMP to myelinate. *Glia*, 59, 720–733. <https://doi.org/10.1002/glia.21144>
- Atanasoski, S., Scherer, S. S., Sirkowski, E., Leone, D., Garratt, A. N., Birchmeier, C., & Suter, U. (2006). ErbB2 signaling in Schwann cells is mostly dispensable for maintenance of myelinated peripheral nerves and proliferation of adult Schwann cells after injury. *Journal of Neuroscience*, 26, 2124–2131.
- Banner, L. R., & Patterson, P. H. (1994). Major changes in the expression of the mRNAs for CDF/LIF and its receptor after injury to adult peripheral nerves and ganglia. *Proceedings of the National Academy of Sciences USA*, 91, 7109–7113.
- Barrette, B., Calvo, E., Vallières, N., & Lacroix, S. (2010). Transcriptional profiling of the injured sciatic nerve of mice carrying the Wld(S) mutant gene: Identification of genes involved in neuroprotection, neuroinflammation, and nerve regeneration. *Brain and Behavioral Immunology*, 24, 1254–1267. <https://doi.org/10.1016/j.bbi.2010.07.249>
- Barrette, B., Hébert, M. A., Filali, M., Lafortune, K., Vallières, N., Gowing, G., ... Lacroix, S. (2008). Requirement of myeloid cells for axon regeneration. *Journal of Neuroscience*, 28, 9363–9376. <https://doi.org/10.1523/JNEUROSCI.1447-08.2008>
- Bauer, S., Kerr, B. J., & Patterson, P. H. (2007). The neuropoietic cytokine family in development, plasticity, disease and injury. *Nature Reviews of Neuroscience*, 8, 221–232. <https://doi.org/10.1038/nrn2054>
- Benito, C., Davis, C. M., Gomez-Sanchez, J. A., Turmaine, M., Meijer, D., Poli, V., ... Jessen, K. R. (2017). STAT3 controls the long-term survival and phenotype of repair Schwann cells during nerve regeneration. *Journal of Neuroscience*, 37, 4255–4269. <https://doi.org/10.1523/JNEUROSCI.3481-16.2017>
- Birchmeier, C., & Nave, K. A. (2008). Neuregulin-1, a key axonal signal that drives Schwann cell growth and differentiation. *Glia*, 56, 1491–1497. <https://doi.org/10.1002/glia.20753>
- Blesch, A., Lu, P., Tsukada, S., Alto, L. T., Roet, K., Coppola, G., ... Tuszynski, M. H. (2012). Conditioning lesions before or after spinal cord injury recruit broad genetic mechanisms that sustain axonal regeneration: superiority to camp-mediated effects. *Experimental Neurology*, 235, 162–173. <https://doi.org/10.1016/j.expneurol.2011.12.037>
- Boerboom, A., Dion, V., Chariot, A., & Franzen, R. (2017). Molecular mechanisms involved in Schwann cell plasticity. *Frontiers in Molecular Neuroscience*, 10, 38. <https://doi.org/10.3389/fnmol.2017.00038>
- Boivin, A., Pineau, I., Barrette, B., Filali, M., Vallières, N., Rivest, S., & Lacroix, S. (2007). Toll-like receptor signaling is critical for Wallerian degeneration and functional recovery after peripheral nerve injury. *Journal of Neuroscience*, 27, 12565–12576. <https://doi.org/10.1523/JNEUROSCI.3027-07.2007>
- Bosse, F., Hasenpusch-Theil, K., Küry, P., & Müller, H. W. (2006). Gene expression profiling reveals that peripheral nerve regeneration is a consequence of both novel injury-dependent and reactivNeurotrophic factors and their receptors in axonal regeneration and functional recovery after peripheral nerve injury. *Journal of Neurochemistry*, 96, 1441–1457. <https://doi.org/10.1111/j.1471-4159.2005.03635.x>
- Boyd, J. G., & Gordon, T. (2003). Neurotrophic factors and their receptors in axonal regeneration and functional recovery after peripheral nerve injury. *Molecular Neurobiology*, 27, 277–324. <https://doi.org/10.1385/MN:27:3:277>
- Bramhall, N. F., Shi, F., Arnold, K., Hochedlinger, K., & Edge, A. S. (2014). Lgr5-positive supporting cells generate new hair cells in the postnatal cochlea. *Stem Cell Reports*, 2, 311–322. <https://doi.org/10.1016/j.stemcr.2014.01.008>
- Brosius Lutz, A., & Barres, B. A. (2014). Contrasting the glial response to axon injury in the central and peripheral nervous systems. *Developmental Cell*, 28, 7–17. <https://doi.org/10.1016/j.devcel.2013.12.002>
- Brosius Lutz, A., Chung, W. S., Sloan, S. A., Carson, G. A., Zhou, L., Lovelett, E., ... Barres, B. A. (2017). Schwann cells use TAM receptor-mediated phagocytosis in addition to autophagy to clear myelin in a mouse model of nerve injury. *Proceedings of the National Academy of Sciences USA*, 114, 8072–8080. <https://doi.org/10.1073/pnas.1710566114>
- Brügger, V., Duman, M., Bochud, M., Münger, E., Heller, M., Ruff, S., & Jacob, C. (2017). Delaying histone deacetylase response to injury accelerates conversion into repair Schwann cells and nerve regeneration. *Nature Communications*, 8, 14272. <https://doi.org/10.1038/ncomms14272>
- Brushart, T. M., Aspalter, M., Griffin, J. W., Redett, R., Hameed, H., Zhou, C., ... Höke, A. (2013). Schwann cell phenotype is regulated by axon modality and central-peripheral location, and persists in vitro. *Experimental Neurology*, 247, 272–281. <https://doi.org/10.1016/j.expneurol.2013.05.007>
- Cafferty, W. B., Gardiner, N. J., Gavazzi, I., Powell, J., McMahon, S. B., Heath, J. K., ... Thompson, S. W. (2001). Leukemia inhibitory factor determines the growth status of injured adult sensory neurons. *Journal of Neuroscience*, 21, 7161–7170.
- Carroll, S. L., Miller, M. L., Frohner, P. W., Kim, S. S., & Corbett, J. A. (1997). Expression of neuregulins and their putative receptors, ErbB2 and ErbB3, is induced during Wallerian degeneration. *Journal of Neuroscience*, 17, 1642–1659.
- Cattin, A. L., Burden, J. J., Van Emmenis, L., Mackenzie, F. E., Hoving, J. J., Garcia Calavia, N., ... Lloyd, A. C. (2015). Macrophage-induced blood vessels guide Schwann cell-mediated regeneration of peripheral nerves. *Cell*, 162, 1127–1139. <https://doi.org/10.1016/j.cell.2015.07.021>



- Cattin, A. L., & Lloyd, A. C. (2016). The multicellular complexity of peripheral nerve regeneration. *Current Opinion in Neurobiology*, 39, 38–46. <https://doi.org/10.1016/j.conb.2016.04.005>
- Cervellini, I., Galino, J., Zhu, N., Allen, S., Birchmeier, C., & Bennett, D. L. (2018). Sustained MAPK/ERK activation in adult Schwann cells impairs nerve repair. *The Journal of Neuroscience*, 38(3), 679–690. <http://doi.org/10.1523/JNEUROSCI.2255-17.2017>
- Chang, L.-W., Viader, A., Varghese, N., Payton, J. E., Milbrandt, J., & Nagarajan, R. (2013). An integrated approach to characterize transcription factor and microRNA regulatory networks involved in Schwann cell response to peripheral nerve injury. *BMC Genomics*, 14, 84. <https://doi.org/10.1186/1471-2164-14-84>
- Chen, Z.-L., & Strickland, S. (2003). Laminin  $\gamma 1$  is critical for Schwann cell differentiation, axon myelination, and regeneration in the peripheral nerve. *The Journal of Cell Biology*, 163(4), 889–899. <http://doi.org/10.1083/jcb.200307068>
- Chen, Z. L., Yu, W. M., & Strickland, S. (2007). Peripheral regeneration. *Annual Reviews of Neuroscience*, 30, 209–233. <https://doi.org/10.1146/annurev.neuro.30.051606.094337>
- Chera, S., & Herrera, P. L. (2016). Regeneration of pancreatic insulin-producing cells by in situ adaptive cell conversion. *Current Opinion in Genetics and Development*, 40, 1–10. <https://doi.org/10.1016/j.gde.2016.05.010>
- Clements, M. P., Byrne, E., Camarillo Guerrero, L. F., Cattin, A. L., Zakka, L., Ashraf, A., ... Parrinello, S. (2017). The wound microenvironment reprograms Schwann cells to invasive mesenchymal-like cells to drive peripheral nerve regeneration. *Neuron*, 96, 98–114. <https://doi.org/10.1016/j.neuron.2017.09.008>
- Cox, B. C., Chai, R., Lenoir, A., Liu, Z., Zhang, L., Nguyen, D. H., ... Zuo, J. (2014). Spontaneous hair cell regeneration in the neonatal mouse cochlea in vivo. *Development*, 141, 816–829. <https://doi.org/10.1242/dev.103036>
- D'Antonio, M., Droggiti, A., Feltri, M. L., Roes, J., Wrabetz, L., Mirsky, R., & Jessen, K. R. (2006). TGF type II receptor signaling controls Schwann cell death and proliferation in developing nerves. *The Journal of Neuroscience*, 26(33), 8417–8427. <http://doi.org/10.1523/JNEUROSCI.1578-06.2006>
- Davis, J., Burr, A. R., Davis, G. F., Birnbaumer, L., & Molkentin, J. D. (2012). A TRPC6-dependent pathway for myofibroblast transdifferentiation and wound healing in vivo. *Developmental Cell*, 23, 705–715. <https://doi.org/10.1016/j.devcel.2012.08.017>
- De Felipe, C., & Hunt, S. P. (1994). The differential control of c-jun expression in regenerating sensory neurons and their associated glial cells. *Journal of Neuroscience*, 14, 2911–2923.
- De, S., Trigueros, M. A., Kalyvas, A., & David, S. (2003). Phospholipase A2 plays an important role in myelin breakdown and phagocytosis during Wallerian degeneration. *Molecular and Cellular Neuroscience*, 24, 753–765.
- Doddrell, R. D., Dun, X. P., Moate, R. M., Jessen, K. R., Mirsky, R., & Parkinson, D. B. (2012). Regulation of Schwann cell differentiation and proliferation by the Pax-3 transcription factor. *Glia*, 60, 1269–1212. <https://doi.org/10.1002/glia.22346>
- Doron-Mandel, E., Fainzilber, M., & Terenzio, M. (2015). Growth control mechanisms in neuronal regeneration. *FEBS Letters*, 589, 1669–1677. <https://doi.org/10.1016/j.febslet.2015.04.046>
- Duregotti, E., Negro, S., Scorsetto, M., Zornetta, I., Dickinson, B. C., Chang, C. J., ... Rigoni, M. (2015). Mitochondrial alarmins released by degenerating motor axon terminals activate perisynaptic Schwann cells. *Proceedings of the National Academy of Sciences U S A*, 112, E497–E505. <https://doi.org/10.1073/pnas.1417108112>
- Eberhard, D., & Tosh, D. (2008). Transdifferentiation and metaplasia as a paradigm for understanding development and disease. *Cellular and Molecular Life Sciences*, 65, 33–40.
- Fabregat, I., Malfettone, A., & Soukupova, J. (2016). New insights into the crossroads between EMT and stemness in the context of cancer. *Journal of Clinical Medicine*, 5, 37. <https://doi.org/10.3390/jcm5030037>
- Faroni, A., Mobasser, S. A., Kingham, P. J., & Reid, A. J. (2015). Peripheral nerve regeneration: Experimental strategies and future perspectives. *Advances in Drug Delivery Reviews*, 82–83, 160–167. <https://doi.org/10.1016/j.addr.2014.11.010>
- Fazal, S. V., Gomez-Sanchez, J. A., Wagstaff, L. J., Musner, N., Otto, G., Janz, M., ... Jessen, K. R. (2017). Graded elevation of c-jun in Schwann cells in vivo: gene dosage determines effects on development, remyelination, tumorigenesis, and hypomyelination. *The Journal of Neuroscience*, 37(50), 12297–12313. <http://doi.org/10.1523/JNEUROSCI.0986-17.2017>
- Figlia, G., Gerber, D., & Suter, U. (2018). Myelination and mTOR. *Glia*, 66, 693–707.
- Fischer, S., Weishaupt, A., Troppmair, J., & Martin, i. R. (2008). Increase of MCP-1 (CCL2) in myelin mutant Schwann cells is mediated by MEK-ERK signaling pathway. *Glia*, 56, 836–843. <https://doi.org/10.1002/glia.20657>
- Florio, F., Ferri, C., Scapin, C., Feltri, M. L., Wrabetz, L., & D'Antonio, M. (2018). Sustained expression of negative regulators of myelination protects Schwann cells from dysmyelination in a Charcot-Marie-Tooth 1B mouse model. *Journal of Neuroscience*, 38, 4275–4287. <https://doi.org/10.1523/JNEUROSCI.0201-18.2018>
- Fontana, X., Hristova, M., Da Costa, C., Patodia, S., Thei, L., Makwana, M., ... Behrens, A. (2012). c-Jun in Schwann cells promotes axonal regeneration and motoneuron survival via paracrine signaling. *Journal of Cell Biology*, 198, 127–141. <https://doi.org/10.1083/jcb.201205025>
- Forge, A., Li, L., & Nevill, G. (1998). Hair cell recovery in the vestibular sensory epithelia of mature guinea pigs. *Journal of Comparative Neurology*, 397, 69–88.
- Forte, E., Chimentelli, I., Rosa, P., Angelini, F., Pagano, F., Calogero, A., ... Messina, E. (2017). EMT/MET at the crossroad of stemness, regeneration and oncogenesis: The Ying-Yang equilibrium recapitulated in cell spheroids. *Cancers (Basel)*, 9, e98. <https://doi.org/10.3390/cancers9080098>
- Fricker, F. R., Antunes-Martins, A., Galino, J., Paramsothy, R., La Russa, F., Perkins, J., ... Bennett, D. L. (2013). Axonal neuregulin 1 is a rate limiting but not essential factor for nerve remyelination. *Brain*, 136, 2279–2297. <https://doi.org/10.1093/brain/awt148>
- Fricker, F. R., Lago, N., Balarajah, S., Tsantoulas, C., Tanna, S., Zhu, N., ... Bennett, D. L. (2011). Axonally derived neuregulin-1 is required for remyelination and regeneration after nerve injury in adulthood. *Journal of Neuroscience*, 31, 3225–3233. <https://doi.org/10.1523/JNEUROSCI.2568-10.2011>
- Fu, S. Y., & Gordon, T. (1997). The cellular and molecular basis of peripheral nerve regeneration. *Molecular Neurobiology*, 14(1-2), 67–116.
- Gambarotta, G., Fregnan, F., Gnani, S., & Perroteau, I. (2013). Neuregulin 1 role in Schwann cell regulation and potential applications to promote peripheral nerve regeneration. *International Review of Neurobiology*, 108, 223–256. <https://doi.org/10.1016/B978-0-12-410499-0.00009-5>
- Glenn, T. D., & Talbot, W. S. (2013). Signals regulating myelination in peripheral nerves and the Schwann cell response to injury. *Current Opinion in Neurobiology*, 23, 1041–1048. <https://doi.org/10.1016/j.conb.2013.06.010>
- Gökbuget, D., Pereira, J. A., Bachofner, S., Marchais, A., Ciaudo, C., Stoffel, M., ... Suter, U. (2015). The Lin28/let-7 axis is critical for myelination in the peripheral nervous system. *Nature Communications*, 6, 8584. <https://doi.org/10.1038/ncomms9584>
- Gomez-Sanchez, J. A., Carty, L., Iruarizaga-Lejarreta, M., Palomo-Irigoyen, M., Varela-Rey, M., Griffith, M., ... Jessen, K. R. (2015). Schwann cell autophagy, myelinophagy, initiates myelin clearance from injured nerves. *Journal of Cell Biology*, 210, 153–168. <https://doi.org/10.1083/jcb.201503019>
- Gomez-Sanchez, J. A., Gomis-Coloma, C., Morenilla-Palao, C., Peiro, G., Serra, E., Serrano, M., & Cabedo, H. (2013). Epigenetic induction of the Ink4a/Arf locus prevents Schwann cell overproliferation during nerve regeneration and after tumorigenic challenge. *Brain*, 136(7), 2262–2278. <http://doi.org/10.1093/brain/awt130>
- Gomez-Sanchez, J. A., Pilch, K. S., van der Lans, M., Fazal, S. V., Benito, C., Wagstaff, L. J., ... Jessen, K. R. (2017). After nerve injury, lineage tracing shows that myelin and Remak Schwann cells elongate extensively and branch to form repair Schwann cells, which shorten radically on remyelination. *Journal of Neuroscience*, 37, 9086–9099. <https://doi.org/10.1523/JNEUROSCI.1453-17.2017>
- Gomis-Coloma, C., Velasco-Aviles, S., Gomez-Sanchez, J. A., Casillas-Bajo, A., Backs, J., & Cabedo, H. (2018). Class IIa histone





- deacetylases link cAMP signaling to the myelin transcriptional program of Schwann cells. *Journal of Cell Biology*, 217, 1249–1268. <https://doi.org/10.1083/jcb.201611150>
- Grande, M. T., Sánchez-Laorden, B., López-Blau, C., De Frutos, C. A., Boutet, A., Arévalo, M., ... Nieto, M. A. (2015). Snail1-induced partial epithelial-to-mesenchymal transition drives renal fibrosis in mice and can be targeted to reverse established disease. *Nature Medicine*, 21, 989–997. <https://doi.org/10.1038/nm.3901>
- Grigore, A. D., Jolly, M. K., Jia, D., Farach-Carson, M. C., & Levine, H. (2016). Tumor budding: The name is EMT. Partial EMT. *Journal of Clinical Medicine*, 5, e51. <https://doi.org/10.3390/jcm5050051>
- Guertin, A. D., Zhang, D. P., Mak, K. S., Alberta, J. A., & Kim, H. A. (2005). Microanatomy of axon/glia signaling during Wallerian degeneration. *Journal of Neuroscience*, 25, 3478–3487. <https://doi.org/10.1523/JNEUROSCI.3766-04.2005>
- Han, S. B., Kim, H., Lee, H., Grove, M., Smith, G. M., & Son, Y. J. (2017). Postinjury induction of activated ErbB2 selectively hyperactivates denervated Schwann cells and promotes robust dorsal root axon regeneration. *Journal of Neuroscience*, 37, 10955–10970. <https://doi.org/10.1523/JNEUROSCI.0903-17.2017>
- Harrisingh, M. C., Perez-Nadales, E., Parkinson, D. B., Malcolm, D. S., Mudge, A. W., & Lloyd, A. C. (2004). The Ras/Raf/ERK signalling pathway drives Schwann cell dedifferentiation. *European Molecular Biology Organisation Journal*, 23, 3061–3071. <https://doi.org/10.1038/sj.emboj.7600309>
- He, X., Zhang, L., Queme, L. F., Liu, X., Lu, A., Waclaw, R. R., ... Lu, Q. R. (2018). A histone deacetylase 3-dependent pathway delimits peripheral myelin growth and functional regeneration. *Nature Medicine*, 24(3), 338–351. <http://doi.org/10.1038/nm.4483>
- Henry, J. J., & Tsonis, P. A. (2010). Molecular and cellular aspects of amphibian lens regeneration. *Progress in Retinal and Eye Research*, 29, 543–555. <https://doi.org/10.1016/j.preteyeres.2010.07.002>
- Hilton, D. A., & Hanemann, C. O. (2014). Schwannomas and their pathogenesis. *Brain Pathology*, 24, 205–220. <https://doi.org/10.1111/bpa.12125>
- Hinz, B., Phan, S. H., Thannickal, V. J., Prunotto, M., Desmoulière, A., Varga, J., ... Gabbiani, G. (2012). Recent developments in myofibroblast biology: Paradigms for connective tissue remodeling. *American Journal of Pathology*, 180, 1340–1355. <https://doi.org/10.1016/j.ajpath.2012.02.004>
- Hirata, K., & Kawabuchi, M. (2002). Myelin phagocytosis by macrophages and nonmacrophages during Wallerian degeneration. *Microscopy Research and Technique*, 57(6), 541–547. <http://doi.org/10.1002/jemt.10108>
- Hirota, H., Kiyama, H., Kishimoto, T., & Taga, T. (1996). Accelerated nerve regeneration in mice by upregulated expression of interleukin (IL) 6 and IL-6 receptor after trauma. *Journal of Experimental Medicine*, 183, 2627–2634.
- Hudson, L. G., Newkirk, K. M., Chandler, H. L., Choi, C., Fossey, S. L., Parent, A. E., & Kusewitt, D. F. (2009). Cutaneous wound reepithelialization is compromised in mice lacking functional slug (Snai2). *Journal of Dermatological Science*, 56, 19–26. <https://doi.org/10.1016/j.jdermsci.2009.06.009>
- Hung, H. A., Sun, G., Keles, S., & Svaren, J. (2015). Dynamic regulation of Schwann cell enhancers after peripheral nerve injury. *Journal of Biological Chemistry*, 290, 6937–6950. <https://doi.org/10.1074/jbc.M114.622878>
- Isaacman-Beck, J., Schneider, V., Franzini-Armstrong, C., & Granato, M. (2015). The lh3 glycosyltransferase directs target-selective peripheral nerve regeneration. *Neuron*, 88(4), 691–703. <http://doi.org/10.1016/j.neuron.2015.10.004>
- Ishii, A., Furusho, M., & Bansal, R. (2013). Sustained activation of ERK1/2 MAPK in oligodendrocytes and Schwann cells enhances myelin growth and stimulates oligodendrocyte progenitor expansion. *Journal of Neuroscience*, 33, 175–186. <https://doi.org/10.1523/JNEUROSCI.4403-12.2013>
- Jacob, C. (2017). Chromatin-remodeling enzymes in control of Schwann cell development, maintenance and plasticity. *Current Opinion in Neurobiology*, 47, 24–30. <https://doi.org/10.1016/j.conb.2017.08.007>
- Jang, S. Y., Shin, Y. K., Park, S. Y., Park, J. Y., Lee, H. J., Yoo, Y. H., ... Park, H. T. (2016). Autophagic myelin destruction by Schwann cells during Wallerian degeneration and segmental demyelination. *Glia*, 64, 730–742. <https://doi.org/10.1002/glia.22957>
- Jessen, K. R., & Mirsky, R. (2005). The origin and development of glial cells in peripheral nerves. *Nature Reviews Neuroscience*, 6, 671–682. <https://doi.org/10.1038/nrn1746>
- Jessen, K. R., & Mirsky, R. (2008). Negative regulation of myelination: Relevance for development, injury, and demyelinating disease. *Glia*, 56, 1552–1565. <https://doi.org/10.1002/glia.20761>
- Jessen, K. R., & Mirsky, R. (2016). The repair Schwann cell and its function in regenerating nerves. *Journal of Physiology*, 594, 3521–3531. <https://doi.org/10.1113/JP270874>
- Jessen, K. R., Mirsky, R., & Arthur-Farraj, P. (2015). The role of cell plasticity in tissue repair: Adaptive cellular reprogramming. *Developmental Cell*, 34, 613–620. <https://doi.org/10.1016/j.devcel.2015.09.005>
- Jessen, K. R., Mirsky, R., & Lloyd, A. C. (2015). Schwann cells: Development and role in nerve repair. *Cold Spring Harbor Perspectives in Biology*, 7, a020487. <https://doi.org/10.1101/cshperspect.a020487>
- Jordan, N. V., Johnson, G. L., & Abell, A. N. (2011). Tracking the intermediate stages of epithelial-mesenchymal transition in epithelial stem cells and cancer. *Cell Cycle*, 10, 2865–2873. <https://doi.org/10.4161/cc.10.17.17188>
- Jung, J., Cai, W., Lee, H. K., Pellegatta, M., Shin, Y. K., Jang, S. Y., ... Park, H. T. (2011). Actin polymerization is essential for myelin sheath fragmentation during Wallerian degeneration. *Journal of Neuroscience*, 31, 2009–2015. <https://doi.org/10.1523/JNEUROSCI.4537-10.2011>
- Kim, D. H., Xing, T., Yang, Z., Dudek, R., Lu, Q., & Chen, Y. H. (2017). Epithelial mesenchymal transition in embryonic development, tissue repair and cancer: A comprehensive overview. *Journal of Clinical Medicine*, 7, e1. <https://doi.org/10.3390/jcm7010001>
- Kim, H. A., Pomeroy, S. L., Whoriskey, W., Pawlitzky, I., Benowitz, L. I., Sicinski, P., ... Roberts, T. M. (2000). A developmentally regulated switch directs regenerative growth of Schwann cells through cyclin D1. *Neuron*, 26, 405–416.
- Kim, S., Maynard, J. C., Strickland, A., Burlingame, A. L., & Milbrandt, J. (2018). Schwann cell O-GlcNAcylation promotes peripheral nerve remyelination via attenuation of the AP-1 transcription factor JUN. *Proceedings of the National Academy of Sciences*, 115(31), 8019–8024. <http://doi.org/10.1073/pnas.1805538115>
- Kioussi, C., Gross, M. K., & Gruss, P. (1995). Pax3: A paired domain gene as a regulator in PNS myelination. *Neuron*, 15, 553–562.
- Kurek, J. B., Austin, L., Cheema, S. S., Bartlett, P. F., & Murphy, M. (1996). Up-regulation of leukaemia inhibitory factor and interleukin-6 in transsected sciatic nerve and muscle following denervation. *Neuromuscular Disorders*, 6, 105–114.
- Kwon, Y. K., Bhattacharyya, A., Alberta, J. A., Giannobile, W. V., Cheon, K., Stiles, C. D., & Pomeroy, S. L. (1997). Activation of ErbB2 during wallerian degeneration of sciatic nerve. *Journal of Neuroscience*, 7, 8293–8299.
- Le, N., Nagarajan, R., Wang, J. Y., Araki, T., Schmidt, R. E., & Milbrand, J. (2005). Analysis of congenital hypomyelinating Egr2Lo/lo nerves identifies Sox2 as an inhibitor of schwann cell differentiation and myelination. *Proceedings of the National Academy of Sciences USA*, 102, 2596–2601. <https://doi.org/10.1073/pnas.0407836102>
- Lee, H. J., Shin, Y. K., & Park, H. T. (2014). Mitogen activated protein kinase family proteins and c-Jun signaling in injury-induced Schwann cell plasticity. *Experimental Neurobiology*, 23, 130–137. <https://doi.org/10.5607/en.2014.23.2.130>
- Liao, T. T., & Yang, M. H. (2017). Revisiting epithelial-mesenchymal transition in cancer metastasis: The connection between epithelial plasticity and stemness. *Molecular Oncology*, 11, 792–804. <https://doi.org/10.1002/1878-0261.12096>
- Lin, H.-P., Oksuz, I., Edward Hurley, E., Wrabetz, L., & Awatramani, R. (2015). Microprocessor complex subunit DiGeorge syndrome critical region gene 8 (Dgcr8) is required for Schwann cell myelination and myelin maintenance. *Journal of Biological Chemistry*, 290, 24294–24307. <https://doi.org/10.1074/jbc.M115.636407>
- Lovisa, S., LeBleu, V. S., Tampe, B., Sugimoto, H., Vadrnagara, K., Carstens, J. L., ... Kalluri, R. (2015). Epithelial-to-mesenchymal transition induces cell cycle arrest and parenchymal damage in renal fibrosis. *Nature Medicine*, 21, 998–1009. <https://doi.org/10.1038/nm.3902>

- Lu, Q. R., Yuk, D., Alberta, J. A., Zhu, Z., Pawlitzky, I., Chan, J., ... Rowitch, D., H. (2000). Sonic hedgehog-regulated oligodendrocyte lineage genes encoding bHLH proteins in the mammalian central nervous system. *Neuron*, 25, 317–332.
- Lubinska, L. (1977). Early course of Wallerian degeneration in myelinated fibres of the rat phrenic nerve. *Brain Research*, 130, 47–63.
- Ma, K. H., Hung, H. A., Srinivasan, R., Xie, H., Orkin, S. H., & Svaren, J. (2015). Regulation of peripheral nerve myelin maintenance by gene repression through Polycomb repressive complex 2. *Journal of Neuroscience*, 35, 8640–8652. <https://doi.org/10.1523/JNEUROSCI.2257-14.2015>
- Ma, K., & Svaren, J. (2016). Epigenomic reprogramming in peripheral nerve injury. *Neural Regeneration Research*, 11(12), 1930–1932. <http://doi.org/10.4103/1673-5374.197133>
- Ma, K. H., & Svaren, J. (2018). Epigenetic control of Schwann cells. *The Neuroscientist*, 24, 206–207. <https://doi.org/10.1177/10738-58417751112>
- Mani, S. A., Guo, W., Liao, M. J., Eaton, E. N., Ayyanan, A., Zhou, A. Y., ... Weinberg, R. A. (2008). The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell*, 133, 704–715. <https://doi.org/10.1016/j.cell.2008.03.027>
- Martini, R., Fischer, S., López-Vales, R., & David, S. (2008). Interactions between Schwann cells and macrophages in injury and inherited demyelinating disease. *Glia*, 56, 1566–1577. <https://doi.org/10.1002/glia.20766>
- Mindos, T., Dun, X. P., North, K., Doddrell, R. D., Schulz, A., Edwards, P., ... Parkinson, D. B. (2017). Merlin controls the repair capacity of Schwann cells after injury by regulating hippo/YAP activity. *Journal of Cell Biology*, 216, 495–510. <https://doi.org/10.1083/jcb.201606052>
- Mizutani, K., Fujioka, M., Hosoya, M., Bramhall, N., Okano, H. J., Okano, H., & Edge, A. S. (2013). Notch inhibition induces cochlear hair cell regeneration and recovery of hearing after acoustic trauma. *Neuron*, 77, 58–69. <https://doi.org/10.1016/j.neuron.2012.10.032>
- Mogha, A., D'Rozario, M., & Monk, K. R. (2016). G protein-coupled receptors in myelinating glia. *Trends in Pharmacological Sciences*, 37, 977–987. <https://doi.org/10.1016/j.tips.2017.01.001>
- Mogha, A., Harty, B. L., Carlin, D., Joseph, J., Sanchez, N. E., Suter, U., ... Monk, K. R. (2016). Gpr126/Adgrg6 has schwann cell autonomous and nonautonomous functions in peripheral nerve injury and repair. *The Journal of Neuroscience*, 36(49), 12351–12367. <http://doi.org/10.1523/JNEUROSCI.3854-15.2016>
- Monje, P. V., Soto, J., Bacallao, K., & Wood, P. M. (2010). Schwann cell dedifferentiation is independent of mitogenic signaling and uncoupled to proliferation: Role of cAMP and JNK in the maintenance of the differentiated state. *Journal of Biological Chemistry*, 285, 31024–31036. <https://doi.org/10.1074/jbc.M110.116970>
- Monk, K. R., Feltri, M. L., & Taveggia, C. (2015). New insights on Schwann cell development. *Glia*, 63, 1376–1393. <https://doi.org/10.1002/glia.22852>
- Monk, K. R., Naylor, S. G., Glenn, T. D., Mercurio, S., Perlin, J. R., Dominguez, C., ... Talbot, W. S. (2009). A G protein-coupled receptor is essential for Schwann cells to initiate myelination. *Science*, 325, 1402–1405. <https://doi.org/10.1126/science.1173474>
- Morel, A. P., Lièvre, M., Thomas, C., Hinkal, G., Ansieau, S., & Puisieux, A. (2008). Generation of breast cancer stem cells through epithelial-mesenchymal transition. *PLoS One*, 3, e2888. <https://doi.org/10.1371/journal.pone.0002888>
- Morgan, L., Jessen, K. R., & Mirsky, R. (1994). Negative regulation of the P0 gene in Schwann cell: suppression of P0 mRNA and protein induction in cultured Schwann cell by FGF2 and TGF beta 1, TGF beta 2 and TGF beta 3. *Development*, 120, 1399–1409.
- Nagarajan, R., Le, N., Mahoney, H., Araki, T., & Milbrandt, J. (2002). Deciphering peripheral nerve myelination by using Schwann cell expression profiling. *Proceedings of the National Academy of Sciences of the USA*, 99, 8998–9003. <https://doi.org/10.1073/pnas.132080999>
- Napoli, I., Noon, L. A., Ribeiro, S., Kerai, A. P., Parrinello, S., Rosenberg, L. H., ... Lloyd, A. C. (2012). A central role for the ERK-signaling pathway in controlling Schwann cell plasticity and peripheral nerve regeneration in vivo. *Neuron*, 73, 729–742. <https://doi.org/10.1016/j.neuron.2011.11.031>
- Negro, S., Bergamin, E., Rodella, U., Duregotti, E., Scorsetto, M., Jalink, K., ... Rigoni, M. (2016). ATP released by injured neurons activates Schwann cells. *Frontiers of Cellular Neuroscience*, 10, 134. <https://doi.org/10.3389/fncel.2016.00134>
- Negro, S., Stazi, M., Marchioretto, M., Tebaldi, T., Rodella, U., Duregotti, E., ... Viero, G. (2018). Hydrogen peroxide is a neuronal alarmin that triggers specific RNAs, local translation of Annexin A2, and cytoskeletal remodeling in Schwann cells. *RNA*, 24, 915–925.
- Newbern, J. M., Li, X., Shoemaker, S. E., Zhou, J., Zhong, J., Wu, Y., ... Snider, W. D. (2011). Specific functions for ERK/MAPK signaling during PNS development. *Neuron*, 69, 91–105. <https://doi.org/10.1016/j.neuron.2010.12.003>
- Niemi, J. P., DeFrancesco-Lisowicz, A., Roldán-Hernández, L., Lindborg, J. A., Mandell, D., & Zigmond, R. E. (2013). A critical role for macrophages near axotomized neuronal cell bodies in stimulating nerve regeneration. *Journal of Neuroscience*, 33, 16236–16248. <https://doi.org/10.1523/JNEUROSCI.3319-12.2013>
- Nieto, M. A., Huang, R. Y., Jackson, R. A., & Thiery, J. P. (2016). EMT: 2016. *Cell*, 166, 21–45. <https://doi.org/10.1016/j.cell.2016.06.028>
- Normén, C., Figlia, G., Pfister, P., Pereira, J. A., Bachofner, S., & Suter, U. (2018). mTORC1 is transiently reactivated in injured nerves to promote c-Jun elevation and Schwann cell dedifferentiation. *Journal of Neuroscience*, 38, 4811–4828. <https://doi.org/10.1523/JNEUROSCI.3619-17.2018>
- Nunan, R., Campbell, J., Mori, R., Pitulescu, M. E., Jiang, W. G., Harding, K. G., ... Martin, P. (2015). Ephrin-Bs drive junctional downregulation and actin stress fiber disassembly to enable wound re-epithelialization. *Cell Reports*, 13, 1380–1395. <https://doi.org/10.1016/j.celrep.2015.09.085>
- Parkinson, D. B., Bhaskaran, A., Arthur-Farraj, P., Noon, L. A., Woodhoo, A., Lloyd, A. C., ... Jessen, K. R. (2008). C-Jun is a negative regulator of myelination. *Journal of Cell Biology*, 181, 625–137. <https://doi.org/10.1083/jcb.200803013>
- Parkinson, D. B., Langner, K., Sharghi Namini, S., Jessen, K. R., & Mirsky, R. (2002).  $\beta$ -Neuregulin and autocrine mediated survival of Schwann cells requires activity of Ets family transcription factors. *Molecular and Cellular Neuroscience*, 20, 154–167. <https://doi.org/10.1006/mcne.2002.1109>
- Parrinello, S., Napoli, I., Ribeiro, S., Wingfield Digby, P., Fedorova, M., Parkinson, D. B., ... Lloyd, A. C. (2010). EphB signaling directs peripheral nerve regeneration through Sox2-dependent Schwann cell sorting. *Cell*, 143, 145–155. <https://doi.org/10.1016/j.cell.2010.08.039>
- Pereira, J. A., Baumann, R., Normén, C., Somandín, C., Mische, M., Jacob, C., ... Suter, U. (2010). Dicer in Schwann cells is required for myelination and axonal integrity. *Journal of Neuroscience*, 30, 6763–6775. <https://doi.org/10.1523/JNEUROSCI.0801-10.2010>
- Petersen, S. C., Luo, R., Liebscher, I., Giera, S., Jeong, S. J., Mogha, A., ... Monk, K. R. (2015). The adhesion GPCR GPR126 has distinct, domain-dependent functions in Schwann cell development mediated by interaction with laminin-211. *Neuron*, 85, 755–769. <https://doi.org/10.1016/j.neuron.2014.12.057>
- Quintes, S., Brinkmann, B. G., Ebert, M., Fröb, F., Kungl, T., Arlt, F. A., ... Nave, K. A. (2016). Zeb2 is essential for Schwann cell differentiation, myelination and nerve repair. *Nature Neuroscience*, 19, 1050–1059. <https://doi.org/10.1038/nn.4321>
- Reynolds, R., & Wolf, C. J. (1992). Terminal Schwann cells elaborate extensive processes following denervation of the motor endplate. *Journal of Neurocytology*, 21, 50–66.
- Rhim, A. D., Mirek, E. T., Aiello, N. M., Maitra, A., Bailey, J. M., McAllister, F., ... Stanger, B. Z. (2012). EMT and dissemination precede pancreatic tumor formation. *Cell*, 148, 349–361. <https://doi.org/10.1016/j.cell.2011.11.025>
- Richardson, R. T., & Atkinson, P. J. (2015). Atoh1 gene therapy in the cochlea for hair cell regeneration. *Expert Opinion on Biological Therapy*, 15, 417–430. <https://doi.org/10.1517/14712598.2015.1009889>
- Roberts, S. L., Dun, X. P., Dee, G., Gray, B., Mindos, T., & Parkinson, D. B. (2017). The role of p38alpha in Schwann cells in regulating peripheral nerve myelination and repair. *Journal of Neurochemistry*, 141(1), 37–47. <https://doi.org/10.1111/jnc.13929>



- Roberts, S. L., Dun, X. P., Doddrell, R. D. S., Mindos, T., Drake, L. K., Onaitis, M. W., ... Parkinson, D. B. (2017). Sox2 expression in Schwann cells inhibits myelination in vivo and induces influx of macrophages to the nerve. *Development*, 144, 3114–3125. <https://doi.org/10.1242/dev.150656>
- Ronchi, G., Gambarotta, G., Di Scipio, F., Salamone, P., Sprio, A. E., Cavallo, F., ... Geuna, S. (2013). ErbB2 receptor over-expression improves post-traumatic peripheral nerve regeneration in adult mice. *PLoS One*, 8, e56282. <https://doi.org/10.1371/journal.pone.0056282>
- Rotshenker, S. (2011). Wallerian degeneration: The innate-immune response to traumatic nerve injury. *Journal of Neuroinflammation*, 8, 109. <https://doi.org/10.1186/1742-2094-8-109>
- Scheib, J., & Höke, A. (2013). Advances in peripheral nerve regeneration. *Nature Reviews of Neurology*, 9, 668–676. <https://doi.org/10.1038/nrneurol.2013.227>
- Scherer, S. S., Kamholz, J., & Jakowlew, S. B. (1993). Axons modulate the expression of transforming growth factor-betas in Schwann cells. *Glia*, 8, 265–276.
- Shaw, T. J., & Martin, P. (2016). Wound repair: A showcase for cell plasticity and migration. *Current Opinion in Cell Biology*, 42, 29–37. <https://doi.org/10.1016/j.cceb.2016.04.001>
- Sheean, M. E., McShane, E., Cheret, C., Walcher, J., Müller, T., Wulf-Goldenberg, A., ... Birchmeier, C. (2014). Activation of MAPK overrides the termination of myelin growth and replaces Nrg1/ErbB3 signals during Schwann cell development and myelination. *Genes and Development*, 28, 290–303. <https://doi.org/10.1101/gad.230045.113>
- Sheu, J. Y., Kulhanek, D. J., & Eckenstein, F. P. (2000). Differential patterns of ERK and STAT3 phosphorylation after sciatic nerve transection in the rat. *Experimental Neurology*, 166, 392–402. <https://doi.org/10.1006/exnr.2000.7508>
- Shin, Y. K., Jang, S. Y., Park, J. Y., Park, S. Y., Lee, H. J., Suh, D. J., & Park, H. T. (2013). The Neuregulin-Rac-MKK7 pathway regulates antagonistic c-Jun/Krox20 expression in Schwann cell dedifferentiation. *Glia*, 61, 892–904.
- Skrypek, N., Goossens, S., De Smedt, E., Vandamme, N., & Bex, G. (2017). Epithelial-to-mesenchymal transition: Epigenetic reprogramming driving cellular plasticity. *Trends in Genetics*, 33, 943–959. <https://doi.org/10.1016/j.tig.2017.08.004>
- Son, Y. J., Trachtenberg, J. T., & Thompson, W. J. (1996). Schwann cells induce and guide sprouting and reinnervation of neuromuscular junctions. *Trends in Neurosciences*, 19, 280–285.
- Stassart, R. M., Fledrich, R., Velanac, V., Brinkmann, B. G., Schwab, M. H., Meijer, D., ... Nave, K. A. (2013). A role for Schwann cell-derived neuregulin-1 in remyelination. *Nature Neuroscience*, 16, 48–54. <https://doi.org/10.1038/nn.3281>
- Stewart, H. J., Rougon, G., Dong, Z., Dean, C., Jessen, K. R., & Mirsky, R. (1995). TGF-betas upregulate NCAM and L1 expression in cultured Schwann cells, suppress cyclic AMP-induced expression of O4 and galactocerebroside, and are widely expressed in cells of the Schwann cell lineage in vivo. *Glia*, 15, 419–436.
- Stoll, G., Griffin, J. W., Li, C. Y., & Trapp, B. D. (1989). Wallerian degeneration in the peripheral nervous system: Participation of both Schwann cells and macrophages in myelin degradation. *Journal of Neurocytology*, 18, 671–683.
- Stolt, C. C., & Wegner, M. (2016). Schwann cells and their transcriptional network: Evolution of key regulators of peripheral myelination. *Brain Research*, 1641, 101–110. <https://doi.org/10.1016/j.brainres.2015.09.025>
- Stratton, J. A., & Shah, P. T. (2016). Macrophage polarization in nerve injury: Do Schwann cells play a role? *Neural Regeneration Research*, 11, 53–57. <https://doi.org/10.4103/1673-5374.175042>
- Sulaiman, W., & Nguyen, D. H. (2016). Transforming growth factor beta 1, a cytokine with regenerative functions. *Neural Regeneration Research*, 11, 1549–1552. <https://doi.org/10.4103/1673-5374.193223>
- Suzuki, K., Lovera, M., Schmachtenberg, O., & Couve, E. (2015). Axonal degeneration in dental pulp precedes human primary teeth exfoliation. *Journal of Dental Research*, 94, 1446–1453. <https://doi.org/10.1177/0022034515593055>
- Syed, N., Reddy, K., Yang, D. P., Taveggia, C., Salzer, J. L., Maurel, P., & Kim, H. A. (2010). Soluble neuregulin-1 has bifunctional, concentration-dependent effects on Schwann cell myelination. *Journal of Neuroscience*, 30, 6122–6131. <https://doi.org/10.1523/JNEUROSCI.1681-09.2010>
- Thiery, J. P., Acloque, H., Huang, R. Y., & Nieto, M. A. (2009). Epithelial-mesenchymal transitions in development and disease. *Cell*, 139, 871–890.
- Thorel, F., Népote, V., Avril, I., Kohno, K., Desgraz, R., Chera, S., & Herrera, P. L. (2010). Conversion of adult pancreatic alpha-cells to beta-cells after extreme beta-cell loss. *Nature*, 464, 1149–1154. <https://doi.org/10.1038/nature08894>
- Truong, K., Ahmad, I., Jason Clark, J., Seline, A., Bertrache, T., Mostaert, B., ... Hansen, M. R. (2018). Nf2 mutation in Schwann cells delays functional neural recovery following injury. *Neuroscience*, 374, 205–213. <https://doi.org/10.1016/j.neuroscience.2018.01.054>
- Vallières, N., Barrette, B., Wang, L. X., Bélanger, E., Thiry, L., Schneider, M. R., ... Lacroix, S. (2017). Betacellulin regulates Schwann cell proliferation and myelin formation in the injured mouse peripheral nerve. *Glia*, 65, 657–669. <https://doi.org/10.1002/glia.23119>
- Varela-Rey, M., Iruarizaga-Lejarreta, M., Lozano, J. J., Aransay, A. M., Fernandez, A. F., Lavin, J. L., ... Woodhoo, A. (2014). S-adenosylmethionine levels regulate the schwann cell DNA methylation. *Neuron*, 81, 1024–1039. <https://doi.org/10.1016/j.neuron.2014.01.037>
- Viader, A., Chang, L. W., Fahrner, T., Nagarajan, R., & Milbrandt, J. (2011). MicroRNAs modulate Schwann cell response to nerve injury by reinforcing transcriptional silencing of dedifferentiation-related genes. *Journal of Neuroscience*, 31, 17358–17369. <https://doi.org/10.1523/JNEUROSCI.3931-11.2011>
- Wagstaff, L., Gomez-Sanchez, J., Mirsky, R., & Jessen, K. R. (2017). The relationship between Schwann cell c-Jun and regeneration failures due to aging and long-term injury. *Glia*, 65, E532.
- Weber, C. E., Li, N. Y., Wai, P. Y., & Kuo, P. C. (2012). Epithelial-mesenchymal transition, TGF- $\beta$ , and osteopontin in wound healing and tissue remodeling after injury. *Journal of Burn Care Research*, 33, 311–318. <https://doi.org/10.1097/BCR.0b013e318240541e>
- Wong, K. M., Babetto, E., & Beirowski, B. (2017). Axon degeneration: Make the Schwann cell great again. *Neural Regeneration Research*, 12, 518–524. <https://doi.org/10.4103/1673-5374.205000>
- Wood, M. D., & Mackinnon, S. E. (2015). Pathways regulating modality-specific axonal regeneration in peripheral nerve. *Experimental Neurology*, 265, 171–175. <https://doi.org/10.1016/j.expneurol.2015.02.001>
- Woodhoo, A., Alonso, M. B., Droggiti, A., Turmaine, M., D'Antonio, M., Parkinson, D. B., ... Jessen, K. R. (2009). Notch controls embryonic Schwann cell differentiation, postnatal myelination and adult plasticity. *Nature Neuroscience*, 12, 839–847. <https://doi.org/10.1038/nn.2323>
- Wu, L. M., Wang, J., Conidi, A., Zhao, C., Wang, H., Ford, Z., ... Lu, Q. R. (2016). Zeb2 recruits HDAC-NuRD to inhibit notch and controls Schwann cell differentiation and remyelination. *Nature Neuroscience*, 19, 1060–1072. <https://doi.org/10.1038/nn.4322>
- Xu, P., Rosen, K. M., Hedstrom, K., Rey, O., Guha, S., Hart, C., & Corfas, G. (2013). Nerve injury induces glial cell line-derived neurotrophic factor (GDNF) expression in Schwann cells through purinergic signalling and the PKC-PKD pathway. *Glia*, 61, 1029–1040. <https://doi.org/10.1002/glia.22491>
- Yang, D. P., Kim, J., Syed, N., Tung, Y. J., Bhaskaran, A., Mindos, T., ... Kim, H. A. (2012). p38 MAPK activation promotes denervated Schwann cell phenotype and functions as a negative regulator of Schwann cell differentiation and myelination. *Journal of Neuroscience*, 32(21), 7158–7168. <https://doi.org/10.1523/JNEUROSCI.5812-11.2012>
- Yang, J., & Weinberg, R. A. (2008). Epithelial-mesenchymal transition: At the crossroads of development and tumor metastasis. *Developmental Cell*, 14, 818–829. <https://doi.org/10.1016/j.devcel.2008.05.009>
- Yanger, K., Zong, Y., Maggs, L. R., Shapira, S. N., Maddipati, R., Aiello, N. M., ... Stanger, B. Z. (2013). Robust cellular reprogramming occurs spontaneously during liver regeneration. *Genes and Development*, 27, 719–724.

- Zhou, S., Gao, R., Hu, W., Qian, T., Wang, N., Ding, G., ... Gu, X. (2014). MiR-9 inhibits Schwann cell migration by targeting Cthrc1 following sciatic nerve injury. *Journal of Cell Science*, 127, 967–976. <https://doi.org/10.1242/jcs.131672>
- Zochodne, D. W. (2012). The challenges and beauty of peripheral nerve regrowth. *Journal of the Peripheral Nervous System*, 17, 1–18. <https://doi.org/10.1111/j.1529-8027.2012.00378.x>

**How to cite this article:** Jessen KR, Arthur-Farraj P. Repair Schwann cell update: Adaptive reprogramming, EMT, and stemness in regenerating nerves. *Glia*. 2018;1–17. <https://doi.org/10.1002/glia.23532>