



## Review

# Inhibiting NF- $\kappa$ B activation by small molecules as a therapeutic strategy

Subash C. Gupta, Chitra Sundaram, Simone Reuter, Bharat B. Aggarwal \*

Cytokine Research Laboratory, Department of Experimental Therapeutics, The University of Texas M. D. Anderson Cancer Center, Houston, TX 77030, USA

## ARTICLE INFO

### Article history:

Received 26 March 2010

Accepted 8 May 2010

Available online 21 May 2010

### Keywords:

Inflammation

NF- $\kappa$ B

Small molecule inhibitors

Therapeutics

## ABSTRACT

Because nuclear factor- $\kappa$ B (NF- $\kappa$ B) is a ubiquitously expressed proinflammatory transcription factor that regulates the expression of over 500 genes involved in cellular transformation, survival, proliferation, invasion, angiogenesis, metastasis, and inflammation, the NF- $\kappa$ B signaling pathway has become a potential target for pharmacological intervention. A wide variety of agents can activate NF- $\kappa$ B through canonical and noncanonical pathways. Canonical pathway involves various steps including the phosphorylation, ubiquitination, and degradation of the inhibitor of NF- $\kappa$ B (I $\kappa$ B $\alpha$ ), which leads to the nuclear translocation of the p50–p65 subunits of NF- $\kappa$ B followed by p65 phosphorylation, acetylation and methylation, DNA binding, and gene transcription. Thus, agents that can inhibit protein kinases, protein phosphatases, proteasomes, ubiquitination, acetylation, methylation, and DNA binding steps have been identified as NF- $\kappa$ B inhibitors. Because of the critical role of NF- $\kappa$ B in cancer and various chronic diseases, numerous inhibitors of NF- $\kappa$ B have been identified. In this review, however, we describe only small molecules that suppress NF- $\kappa$ B activation, and the mechanism by which they block this pathway.

© 2010 Elsevier B.V. All rights reserved.

## 1. Introduction

The nuclear factor- $\kappa$ B (NF- $\kappa$ B) signaling pathway plays a major role in the development, maintenance, and progression of most chronic diseases. NF- $\kappa$ B controls the expression of genes involved in a number of physiological responses, including immune inflammatory responses, acute-phase inflammatory responses, oxidative stress responses, cell adhesion, differentiation, and apoptosis [1]. Recent studies have suggested that NF- $\kappa$ B dysregulation is associated with many diseases including AIDS, atherosclerosis, asthma, arthritis, diabetes, inflammatory bowel disease, stroke, muscle wasting and viral infections. Mounting evidence indicates that NF- $\kappa$ B acts as a link

between inflammation and cancer progression [2–11], making NF- $\kappa$ B essential to and a potential drug target in hematological malignancies and solid tumors [12,13].


NF- $\kappa$ B was first identified in 1986 by Sen and Baltimore [6] in the nucleus bound to an enhancer element of the immunoglobulin kappa light chain gene in B cells [6,14]. It is now known to be ubiquitous in nature present in all the cell types and is evolutionary conserved. It belongs to the family of Rel proteins that includes c-Rel, RelA (p65), RelB, NF- $\kappa$ B1 (p50 and its precursor p105), and NF- $\kappa$ B2 (p52 and its precursor p100) all of which can form hetero- or homodimers [15–17].

NF- $\kappa$ B activation is tightly regulated mainly through its localization. In resting cells, NF- $\kappa$ B proteins are kept in the cytoplasm in association with inhibitory I $\kappa$ B proteins including I $\kappa$ B $\alpha$ , I $\kappa$ B $\beta$ , and I $\kappa$ B $\epsilon$  [16] among which I $\kappa$ B $\alpha$  is the most abundant. NF- $\kappa$ B signaling occurs through the canonical (classical) pathway initiated by NF- $\kappa$ B1 (p50/p105) and a noncanonical (alternative) pathway initiated by NF- $\kappa$ B2 (p52/p100) (Fig. 1). Before the active NF- $\kappa$ B is translocated into the nucleus, NF- $\kappa$ B1 and NF- $\kappa$ B2 are cleaved to the active p50 and p52 subunits, respectively. While the classical pathway depends on IKK complex consisting of IKK $\alpha$ , IKK $\beta$ , IKK $\gamma$  and the inhibitory subunit I $\kappa$ Bs, the alternative pathway depends on IKK $\alpha$  homodimers and NF- $\kappa$ B inducing kinase (NIK) [18–20]. During classical activation, the IKK complex specifically phosphorylates I $\kappa$ Bs on two conserved N-terminal serine residues which target them for E2- and E3-ligase-mediated polyubiquitination and subsequent 26S proteasomal mediated degradation. This process releases and activates NF- $\kappa$ B which now translocates to the nucleus. The activation of alternative pathway, which is commonly associated with RelB results in regulated processing of the p100 precursor protein to p52 and subsequent translocation of p52–RelB

**Abbreviations:** AgR, antigen receptor; ATM, ataxia-telangiectasia mutant; BAFF, B-cell activating factor; BCL, B-cell lymphoma; BCR, B cell receptor; CARMA, CARD-containing MAGUK protein; CD40L, CD40 ligand; CK, casein kinase; DSBS, Double-stranded DNA breaks; ECSIT, evolutionary conserved signaling intermediates on Toll pathways; EGF, epidermal growth factor; EGFR, EGF receptor; ELKS, glutamate, leucine, lysine, serine-rich protein; GSK, glycogen synthase kinase; Hsp90, heat shock protein 90; I $\kappa$ B, inhibitor of NF- $\kappa$ B; IKK, I $\kappa$ B kinase; IRAK, IL-1R-associated kinase; LT $\beta$ , lymphotoxin  $\beta$ ; LPS, lipopolysaccharide; MALT, mucosa-associated lymphoid tissue; MAPK, mitogen activated protein kinase; MAPK/Erk, kinase kinase; MyD88, myeloid differentiation factor; NF- $\kappa$ B, nuclear factor- $\kappa$ B; NIK, NF- $\kappa$ B-inducing kinase; NEMO, NF- $\kappa$ B essential modulator; PDK, Phosphoinositide-dependent kinase; PI3K, phosphatidylinositol 3-kinase; PKC, protein kinase C; PLC, phospholipase C; RANKL, receptor activator of NF- $\kappa$ B ligand; RIP, receptor-interacting protein; Syk, Spleen tyrosine kinase; TAB, TAK1-binding protein; TAK, transforming growth factor- $\beta$ -activated kinase; TCR, T cell receptor; TLR, Toll-like receptor; TNF, tumour necrosis factor; TNFR1, TNF receptor 1; Tpl2, tumour progression locus-2; TRADD, TNF-receptor-associated death domain protein; TRAF, TNF-receptor-associated factor

\* Corresponding author. Tel.: +1 713 794 1817; fax: +1 713 745 6339.

E-mail address: [aggarwal@mdanderson.org](mailto:aggarwal@mdanderson.org) (B.B. Aggarwal).




**Fig. 1.** Schematic representation of major NF- $\kappa$ B activation pathways. In the classical pathway, binding of TNF $\alpha$  to the receptor triggers the sequential recruitment of the adaptors TRADD, TRAF2 and RIP to the membrane. TRAF2 then recruits the IKK complex composed of IKK $\alpha$ , IKK $\beta$  and IKK $\gamma$  (NEMO) through mediation of kinases like TAK1, MEKK1, MEKK3. Activation of the IKK complex leads to the phosphorylation and ubiquitination of I $\kappa$ B $\alpha$  at specific residues followed by its degradation via the proteasome pathway. The p105 subunit of NF- $\kappa$ B then undergoes GSK3 $\beta$  and Tpl2 mediated phosphorylation at S<sup>903</sup> and S<sup>907</sup> and subsequent degradation. The heterodimer p50–p65 is then released and migrates to the nucleus where it undergoes a series of posttranslational modifications including phosphorylation, acetylation and methylation and binds to specific  $\kappa$ B sites and activates NF- $\kappa$ B target genes [49,209,210]. The alternative pathway is IKK $\gamma$  independent and is triggered by binding of the CD40, RANK, LT $\beta$ R, BAFF ligands to their receptor, leading to recruitment of TRAF proteins and the sequential activation of NIK and IKK $\alpha$ . Activation of IKK $\alpha$  then induces the processing of the inhibitory protein p100. p100 proteolysis releases p52 which then translocates to the nucleus and triggers transcription of NF- $\kappa$ B target genes [211]. NF- $\kappa$ B activation in response to bacterial endotoxin LPS involves Toll like receptor and is mediated through recruitment of MyD88, TRAF6 and ECSIT. Recruitment of these adaptors leads to sequential activation of IRAK1/2 and IKK and eventual release of active NF- $\kappa$ B [215]. NF- $\kappa$ B activation by pervanadate and H<sub>2</sub>O<sub>2</sub> induces phosphorylation of I $\kappa$ B $\alpha$  at Tyr<sup>42</sup> by protein tyrosine kinase like Syk. The Tyr phosphorylation does not lead to I $\kappa$ B $\alpha$  degradation but makes the binding weak thereby dissociating the I $\kappa$ B $\alpha$  and releasing active NF- $\kappa$ B to the nucleus [216,217]. Antigen receptor viz., T-cell receptor and B-cell receptor mediated signaling to NF- $\kappa$ B activation depends on recruitment of a trimolecular protein complex CARMA1–BCL10–MALT1. In this pathway PKC $\theta$  (in T cells) and PKC $\zeta$  (in B cells) along with other kinases act upstream to the trimolecular complex to promote IKK $\gamma$  polyubiquitination and consequent IKK activation. Activation of IKK through this pathway involves mediation of TRAF2, TRAF6, TAK1 and TAB1 [218,219]. A novel pathway of NF- $\kappa$ B activation originating from the nucleus is associated with DNA damage. Double-stranded DNA breaks in response to genotoxic agents initiate signals that trigger SUMOylation of nuclear-localized IKK $\gamma$ , preventing its nuclear export. Concomitantly, these breaks activate ATM which phosphorylates SUMO-modified IKK $\gamma$ , promoting the removal of SUMO and enhancing IKK $\gamma$  ubiquitination. Ubiquitinated IKK $\gamma$  then translocates to the cytoplasm, where it activates IKK in cooperation with ATM and the ELKS protein, leading to I $\kappa$ B $\alpha$  phosphorylation and degradation, p65 nuclear translocation and induction of NF- $\kappa$ B dependent target genes [220–223]. NF- $\kappa$ B can also be regulated by phosphatases. WIP1, a Ser/Thr phosphatase was recently shown to negatively regulate NF- $\kappa$ B activation by dephosphorylating p65 at Ser<sup>536</sup> [80].

heterodimers to the nucleus [19]. Although NF- $\kappa$ B activation occurs mainly through canonical and non-canonical pathways, during the past decade a number of pathways for NF- $\kappa$ B activation has been elucidated (Fig. 1).

Once in the nucleus, activated NF- $\kappa$ B undergoes a series of posttranslational modifications, including phosphorylation, acetylation, and methylation. These modifications regulate both the strength and duration of NF- $\kappa$ B activity. RelA/p65 is directly phosphorylated by

cAMP-dependent protein kinase (PKA) at Ser<sup>276</sup>, casein kinase II (CKII) at Ser<sup>529</sup>, and IKK at Ser<sup>536</sup> [21,22]. RelA dephosphorylation by protein phosphatase 2A (PP2A) has been reported to decrease NF- $\kappa$ B activity [23]. RelA is subject to inducible acetylation by p300/CBP, and acetylated RelA interacts weakly, if at all, with I $\kappa$ B $\alpha$  [24,25], but maintains its nuclear localization and NF- $\kappa$ B transcriptional response. RelA is also subject to methylation by lysine methyltransferase Set9 (also called Set7 or KMT7) at Lys<sup>314/315</sup> [26].

**Fig. 2.** A list of gene products regulated by NF- $\kappa$ B. These genes include transcription factors, cell-surface receptors, growth factors, immunoreceptors, acute phase proteins, enzymes, stress response genes, early response genes, viruses, apoptosis regulators, cytokines/chemokines and cell adhesion molecules. For more information, see <http://www.nf-kb.org>.





Activated NF- $\kappa$ B binds to specific DNA sequences in target genes, which are designated as  $\kappa$ B elements, and regulates the transcription of over 500 genes involved in immunoregulation, growth regulation, inflammation, carcinogenesis, and apoptosis (Fig. 2). NF- $\kappa$ B is frequently constitutively activated in patients with chronic inflammatory conditions such as cancer and pulmonary, cardiovascular, autoimmune, skin, and neurodegenerative diseases [27]. NF- $\kappa$ B's ability to control multiple genes involved in human diseases makes the NF- $\kappa$ B signaling pathway a novel target for therapy [28,29].

Due to the various levels of regulation, NF- $\kappa$ B signaling pathway can be potentially targeted at various levels including kinases, phosphatases, ubiquitination, nuclear translocation, DNA binding, protein acetyl transferases and methyl transferases (Fig. 3).

## 2. Inhibitors of the NF- $\kappa$ B activation pathway

Given NF- $\kappa$ B's relevance in human diseases and the fact that many drugs interfere with NF- $\kappa$ B signaling, the NF- $\kappa$ B signaling pathway provides a highly attractive target for the therapeutic development. More than 700 inhibitors of the NF- $\kappa$ B activation pathway, including antioxidants, peptides, small RNA/DNA, microbial and viral proteins, small molecules, and engineered dominant-negative or constitutively active polypeptides have been described (Table 1). Several of these molecules act as general inhibitors of NF- $\kappa$ B activation, while other molecules target specific steps; some molecules possibly target multiple steps in the NF- $\kappa$ B pathway (Fig. 3).

### 2.1. Inhibition of protein kinases

NF- $\kappa$ B activation requires the phosphorylation, polyubiquitination, and subsequent degradation of its inhibitory subunit, I $\kappa$ B $\alpha$ . Hence, inhibiting I $\kappa$ B $\alpha$  phosphorylation ultimately inhibits NF- $\kappa$ B's transcriptional activity [30]. I $\kappa$ B $\alpha$  phosphorylation is carried out by IKK, a serine/threonine protein kinase composed of three basic subunits: the kinases IKK $\alpha$ , IKK $\beta$ , and the regulatory subunit IKK $\gamma$  (NEMO). The IKK activation is usually the first common step in the integration of many NF- $\kappa$ B-activating pathways; therefore, one strategy for inhibiting NF- $\kappa$ B activation is to block IKK activation. However, although more than 150 agents have been shown to inhibit NF- $\kappa$ B activation at the IKK step, few studies have investigated the mechanism by which a given agent can inhibit IKK or its activation. The few IKK inhibitors for which a mechanism of action is known can be divided into three general groups: adenosine triphosphate (ATP) analogs, which show some specificity for interacting with IKK; compounds that have allosteric effects on IKK structure; and compounds that interact with a specific cysteine residue (Cys-179) in the activation loop of IKK $\beta$ . ATP analogs include natural products such

as  $\beta$ -carboline and synthetic compounds such as SC-839, which has an approximately 200-fold preference for IKK $\beta$  compared to IKK $\alpha$  [28,31]. Compounds that have allosteric effects on IKK structure include BMS-345541, a synthetic compound that binds to an allosteric site on both IKK $\alpha$  and IKK $\beta$  and has an approximately 10-fold greater inhibitory effect on IKK $\beta$  than on IKK $\alpha$  [32]. Compounds that interact with Cys-179 IKK $\beta$  include thiol-reactive compounds such as parthenolide, arsenite, and certain epoxyquinoids [33–36]; these compounds' interactions with Cys-179 are believed to interfere with phosphorylation-induced IKK $\beta$  activation because Cys-179 is located between Ser<sup>177</sup> and Ser<sup>181</sup>, which are required for IKK $\beta$  activation in response to upstream signals such as tumor necrosis factor (TNF) and lipopolysaccharide (LPS) [37,38].


Gene-based inhibitors can also block IKK activation. Specifically, mutations at the ATP-binding site or in the kinase activation loop can create dominant-negative IKK $\alpha$  and IKK $\beta$ , which are capable of blocking NF- $\kappa$ B activation [39–43]. Because of their distinct roles in the canonical and non-canonical NF- $\kappa$ B activation pathways, dominant-negative IKK mutants can show stimulus-dependent inhibition [44]. Adenoviral-mediated delivery of an IKK $\beta$  dominant-negative kinase has been shown to have therapeutic potential for airway inflammatory diseases such as asthma [45,46]. NEMO can also serve as a target for inhibiting the IKK complex. In particular, introducing a cell-permeable 10 amino-acid peptide that corresponds to the NEMO-binding domain of IKK $\beta$  can block the binding of NEMO to IKK in response to TNF in the canonical pathway [47].

While activation of NF- $\kappa$ B by many stimuli depends on the phosphorylation of I $\kappa$ Bs at N-terminal sites by the IKK complex, the mechanism of NF- $\kappa$ B activation by ultraviolet (UV) radiation involves the IKK-independent phosphorylation of I $\kappa$ B $\alpha$  at a cluster of C-terminal sites that are recognized by casein kinase II (CKII). CKII activity toward I $\kappa$ B $\alpha$  depends on p38 mitogen-activated protein kinase (MAPK) activation. CKII's role as a key survival signal that activates NF- $\kappa$ B and protects tumor cells from apoptosis suggests that CKII may be an attractive target for the treatment of diverse cancers. Apigenin, a plant flavonoid, and emodin, a plant anthraquinone, are competitive inhibitors of CKII that directly interact with the nucleotide-binding sites of CKII [48].

Besides phosphorylating and subsequently degrading the molecules that inhibit NF- $\kappa$ B, protein kinases can also target the functional domains of NF- $\kappa$ B proteins themselves to optimally activate NF- $\kappa$ B. NF- $\kappa$ B proteins can be phosphorylated in the cytoplasm or nucleus by such kinases as glycogen synthase kinase 3 $\beta$  (GSK3 $\beta$ ) [49], TRAF-associated NF- $\kappa$ B activator (TANK)-binding kinase 1 (TBK1) [50,51], PKAc [21], mitogen- and stress-activated protein kinase-1 (MSK-1) [52], MAP3K NIK [53], Tpl2, PKC- $\theta$  [54], PI3K, Akt [55–57], p38 MAPK [58], protein tyrosine kinase, PKC- $\delta$  [59], RHO-kinase 2 [60], mitogen-activated protein kinase kinase 3 (MEKK3) [61], and receptor tyrosine kinases such as epidermal growth factor receptor, human epidermal growth factor receptor 2 [62]. Antagonistic antibodies or kinase inhibitors that target these molecules may decrease NF- $\kappa$ B activation. Some kinase inhibitors that have the potential to inhibit NF- $\kappa$ B activation include SB203580 and PD0980589 (MAPK inhibitors) [58]; denbinobin (TAK1 inhibitor) [63]; tyrosine kinase inhibitors [62]; rhein, (an MEKK inhibitor) [64,65]; TNAP, betaine (NIK inhibitors) [66–68], epoxyquinol B (a TAK1 crosslinker) [69]; M2L (an extracellular signal-regulated kinase 2 inhibitor) [70,71]; CCK-8 (a p38 kinase kinase inhibitor) [72], KSR2 (an MEKK3 inhibitor) [73], golli BG21 (a PKC inhibitor) [74].

### 2.2. Inhibition of NF- $\kappa$ B activation by protein phosphatases

Because protein phosphorylation is a dynamic process whereby phosphatases counterbalance kinase action, phosphatases may be used to inhibit NF- $\kappa$ B activation. Protein phosphatase 2A is a serine/threonine phosphatase that has been reported to dephosphorylate and modulate the activity of IKK $\beta$  [75]. Cytosine arabinoside, a pyrimidine analogue used to effectively treat acute leukemia, has been reported to induce apoptosis by activating protein phosphatases



**Fig. 3.** Potential targets for inhibition of NF- $\kappa$ B activation pathway. NF- $\kappa$ B can be targeted at various steps including protein kinases, protein phosphatases, ubiquitination step, acetylation step, methylation step, nuclear translocation step and DNA binding step.

Table 1

A list of small molecules as inhibitors of NF-κB pathway.

<b>A. Upstream NF-κB</b>	ox-LDL	Cyclopentenones	<i>Poncirus trifoliata</i> fruit extract	JM34	<b>Natural product</b>
<b>Natural product</b>	Panduratin A	CYL-19 s	Probiotics	KIOM-79	Actinodaphnine
15d-PGJ(2)	PEITC	CYL-26z	Prostaglandin	Leptomycin B	Anthocyanins
Calagualine	Petrosaspongiolide M	Diarylpyridine derivative	Resiniferatoxin	Neomycin	<i>Arnica montana</i> extract
Conophylline	Phytic acid	DPE	Stinging nettle extracts	Nuculing	Artemisinin
Evodiamine	Piceatannol	Epoxyquinone	Thiopental	Oregonin	Baicalein
Geldanamycin	Pinosylvin	Gabexate mesilate	Tipifarnib	OXPAFC	Bambara groundnut
Perrilyl alcohol	<i>Plagius flosculosus</i> extract	Gleevec	Titanium	Paeoniflorin	β-lapachone
PSK	Plumbagin	Hydroquinone	TNP-470	Phallacidin	Biliverdin
Rocaglamides	Pomegranate extract	Ibuprofen	<i>Trichomomas vaginalis</i>	Piperine	Brazilian
<b>Viral protein</b>	Prostaglandin A1	IQCAD	TG-rich lipoproteins	Pitavastatin	Calcitriol
Adenovirus E1A	Quercetin	Indolecarboxamide	Ursodeoxycholic acid	<i>Platycodi radix</i> extract	Camphothecin
NS5A (Hep-C virus)	Rengyolone	Isobutyl nitrite	<i>Xanthium strumarium</i> extract	Probiotics	<i>Sutherlandia frutescens</i>
<b>Protein</b>	Rosmarinic acid	Jesterone dimer	β- PEITC	Rapamycin	Capsiate
Erbin overexpression	Rottlerin	15-deoxyspergualine analog	8-MSO	<i>Rhubarb aqueous</i> extract	Catalposide
Golli BG21	Saikosaponin-d	Methotrexate	β-lapachone	<i>Salvia miltiorrhiza</i> extract	Cat's claw bark
KSR	<i>Salvia miltiorrhiza</i> extract	MLB120**	<b>Peptide</b>	SH extract	Cheongyeolsaseuptang***
MAST205	Sanguinarine	Monochloramine	Penetratin	Selenomethionine	Chitosan
NPM-ALK oncoprotein	SAm extract	MX781 (Retinoid antagonist)	VIP	Shenfu	Chicory root
Hep-C virus protease	Staurosporine	4-HPR	<b>Protein</b>	<i>Sophora radix</i> extract	CSPDP
PDF	Sesquiterpene lactones	Nafamostat mesilate	Activated protein C	Sopoongsan	Clarithromycin
Rituximab	Scoparone	NSAIDs	HSP-70	<i>Sorbus commixta</i> extract	Cloricromene
TNAP	Silibinin	PS-1145 (MLN1145)	Interleukin-13	Sphondin	C-K and Rh(2)
<b>Synthetic</b>	Silymarin	PQD	Intravenous Ig	<i>T. polyglycosides</i>	<i>Cortex cinnamomi</i> extract
Betaine	Sulforaphane	Pyridooxazinone derivative	Murr1 gene product	Younggaechulgum-tang***	Cryptotanshinone
Desloratadine	Sulindac	SC-514**	Neurofibromatosis-2 protein	α-pinene	Cytochalasin D
LY29 and LY30	Tetrandine	Scytonemin	PACAP	<b>Peptide</b>	Black rice extract
MOL 294**	Theaflavin	Sodium salicylate	SAIF	NCPP	Danshenshu
Pefabloc	Thienopyridine	Statins (several)	ST2 (IL-1-like receptor)	PN50	Diterpenoids
Rhein	Tilianin	Sulfasalazine	α-MSH	RelA peptides (P1 & P6)	Ent-kaurane diterpenoids
SMI and FP	Ursolic acid	Sulfasalazine analogs	γ-glutamylcysteine synthetase	<b>Viral Protein</b>	Epinaestine hydrochloride
<b>B. IKK activity and</b>	Vesnarinone	Survanta	<b>Bacterial/Viral Protein</b>	3C protease (EMC virus)	Epoxyquinol A
<b>IkB phosphorylation</b>					Erythromycin
<b>Natural product</b>	Wedelolactone	Thalidomide	K1L (Vaccinia virus protein)	Canine Distemper Virus	Evodiamine
[6]-gingerol	Withanolides	THI 52	Nef (HIV-1)	MNF (myxoma virus)	Fish oil
1'-Acetoxychavicol acetate	Xanthoangelol D	YC-1*	Vpu protein (HIV-1)	<b>Protein</b>	<i>Fomes fomentarius</i> extracts
20(S)-Protopanaxatriol	Zerumbone	<b>Others</b>	YopJ	C5a	Fucoidan
4-Hydroxynonenal	β-carboline	Lead	<b>Synthetic</b>	DQ 65-79	Gallic acid
Acetyl-boswellic acids	γ-mangostin	Mild hypothermia	1-Bromopropane	Fox1j	<i>Ganoderma lucidum</i>
Anandamide	γ-Tocotrienol	Saline (low Na+)	Acetaminophen	GILZ	Garcinol
Anethole	<b>Peptide</b>	<b>C. IkB degradation</b>	Diamide	HSCO	Geranylgeraniol
Apigenin	IKKβ peptide	<b>Natural product</b>	Dobutamine	HSP-72	Ginkgolide B
Artemisia vestita <sup>1</sup>	NEMO CC2-LZ peptide	5'-methylthioadenosine	E-73	Interleukin-10	Glycyrrhizin
Baoganning	<b>Protein</b>	<i>Artemisia iwaiyomogi</i> extract	Ecabet sodium	Interleukin -11	Halofuginone
Betulinic acid	Anti-thrombin III	Alachlor	Gabexate mesilate	Interleukin -13	Hematein
Black raspberry extracts	Chorionic gonadotropin	Amentoflavone	Glimepiride	MTS-SR-1:β	Herbal compound 861
Buddlejasaponin IV	FHIT	<i>Antrodia camphorata</i> #	Hypochlorite	Onconase	Hydroxyethyl starch
Cacospongionolide B	HB-EGF	<i>Artemisia capillaries</i> extract	Losartin	RASSF1A gene	Hydroxyethylpuerarin
Calagualine	Hepatocyte growth factor	Aucubin	LY294002	ROR-α	Hypericin
Cardamomin	Interferon-α	Baicalein	Pervanadate	Surfactant protein A	Kamebakaurin
Cardamonin	Interleukin-10	Blackberry extract	Phenylarsine oxide	TAT-SR-1:β	Linoleic acid
Casparol	PAN1	Buchang-tang***	Phenytol	ZAS3 protein	Lithospermi radix
Cobrotoxin	PTEN	Capsaicin	Ro106-9920**	ZUD protein	Macrolide antibiotics
Cycloepoxydon	SOC51	Catalposide	Sabaeksan	β-amyloid protein	Mediterranean plant extracts
Decursin	<b>Viral Protein</b>	Cyclolinteinone	U0126 (MEK inhibitor)	<b>Synthetic</b>	2-methoxyestradiol
Dehydroascorbic acid	Adenovirus	Dihydroarteannin	<b>Others</b>	BMD	6-MITC
Dexanabinol	Core protein (Hep-C virus)	Docosahexaenoic acid	Vagus nerve stimulation	Carbaryl	Nicotine
Digitoxin	Cytomegalovirus	Emodin	Low level laser therapy	CGS 25462	<i>Ochna macrocalyx</i> bark ext.
Diosgenin	E7 (Papillomavirus)	<i>Ephedrae herba</i> (Mao) extract	Zinc	DHMEQ	Oridonin
Diterpenes	MC159	Equol	<b>D. IkB upregulators/</b>	Diltiazem	PC-SPES (8 herb mixture)
Docosahexaenoic acid	MC160	Erbstatin	<b>NF-κB translocation</b>	Dioxin	PGG
Falcarindol	NS5B (Hep-C virus)	Estrogen	<b>Natural product</b>	Dipyridamole	Pepluanone
Flavopiridol	vIRF3 (KSHV)	Ethacrynic acid	PGG	Disulfiram	<i>Phyllanthus amarus</i> extracts
Furonaphthoquinone	<b>Synthetic</b>	Fosfomycin	15-deoxyspergualin	Enalapril	Plant compound A
Garcinone B	AIDCA derivative	Fungal gliotoxin	2',8"-biapigenin	mEET	Polyozellin
<i>Glossogyne tenuifolia</i> extract	TDZD	Gamisanghyulyunbueum***	5F (from <i>Pteri syemipinnata</i> )	Fluvastatin	Prenylbisabolane 3
Glycine chloramine	TPCA-1	Genipin	<i>Agastache rugosa</i> leaf extract	Indole-3-carbinol	Prostaglandin E2
Guggulsterone	Pyridine derivatives	Genistein	Alginic acid	JSH-23	PSK
Herbimycin A	ACHP	Glabridin	<i>Antrodia camphorata</i> extract	KL-1156	Quinic acid
Honokiol	Acrolein	Glucosamine sulfate	Apigenin	Lefflunomide	Sanggenon C
Hypoestoxide	AGRO100*	Glutamine	Astragaloside IV	Levamisole	Sesamin
Indirubin-3'-oxime	Amino-pyrimidine	Gumiganghwaltang***	AT514 (serratomolide)	MEB	Shen-Fu#
Isorhapontigenin	AS602868**	Isomallotochromanol	Atorvastatin	Moxifloxacin	Silibinin
		Isomallotochromene			
Kahweol	Aspirin	<i>Kochia scoparla</i> fruit extract	Blue honey suckle extract	Omapatrilat	Sinomenine

(continued on next page)

Kava derivatives <sup>2</sup> Licorce extracts Manumycin A Monochloramine N-acetylcysteine Nitric oxide	Azidothymidine BAY-11-7082 BAY-11-7083 Benzoimidazole derivative Benzyl isothiocyanate BMS-345541	L-ascorbic acid Leflunomide metabolite Melatonin Midazolam Momordin I <i>Morinda officinalis</i> extract	<i>Buthus martensi</i> extract Cantharidin Chiisanoside Clarithromycin <i>Cornus officinalis</i> extract Eriocalyxin B	R-etodolac Rolipram SC236 (COX-2 inhibitor) Triflusal Volatile anesthetics <b>E. NF-κB DNA-binding</b> <b>Inorganic Complex</b> Metals (chromium, cadmium, gold, lead, mercury, zinc, arsenic)	Sword brake fern extract <i>Tanacetum larvatum</i> extract Tansinones Taurine + niacine TZD MCC-555 Trichostatin A
Nitrosylcobalamin	Carboplatin	<i>Mosla dianthera</i> extract	Gangliosides		Triptolide
Oleandrin Omega 3 fatty acids Withaferin A Xanthohumol Xylitol Yan-gan-wan <sup>#</sup> Yin-Chen-Hao <sup>#</sup> <i>Yucca schidigera</i> extract <b>Peptide</b> Ghrelin Peptide YY Rapamycin <b>Viral Protein</b> African Swine Fever virus Sendai Virus-CV proteins E1B (Adenovirus) ICP27 (HSV-1) H4/N5 (bracovirus) NS3/4A (Hep-C) <b>Protein</b> Adiponectin AIM2 overexpression Angiopoietin-1 Antithrombin AvrA protein (Salmonella) β-catenin Bromelain	CDDO-Me CHS 828* Compound 5 Compound A <b>Synthetic</b> ADP ribosylation inhibitor 7-amino-4-methylcoumarin Amrinone Atrovastat Benfotiamine Benzamide Bisphenol A Caprofen Carbocisteine Celecoxib Germcitabine Cinnamaldehyde 2-methoxy CNA 2-hydroxy CNA CDS CP Compound Cyanoguanidine HMP α-difluoromethylornithine DTD	<i>Opuntia ficus indica</i> extract <i>Platycodin</i> saponins Polymyxin B Ritonavir Sotiglitazone Roxithromycin DAAS Serotonin derivative Simvastatin SM-7368 <sup>**</sup> T-614 Sulfasalazine SUN C8079 Triclosan plus CPC Tobacco smoke Verapamil <b>Others</b> Heat (fever-like) Hypercapnic acidosis Hyperosmolarity Hypothermia Alcohol <b>E. NF-κB transactivation</b> <b>Natural products</b> 4'-DM-6-Mptox	Glucocorticoids <i>HP</i> extracts Hirsutenone Human breast milk BZLF1 (EBV protein) SH gene products (PMV) <b>Protein</b> Antithrombin NF-kappaB-repression factor PIAS3 PTX-B <b>Synthetic</b> 17-AAG TMFC AQC derivatives 9-aminoacridine derivatives Chromene derivatives D609 Dimethylfumarate EMDPC Histidine HIV-1 PI Mesalamine PEITC Pranlukast	Apocynin Apple juice/extracts Arctigenin Aretemisa p7F Astaxanthin Benidipine bis-eugenol <i>BG</i> compounds BHA CAPE Carnosol Carvedilol Catechol derivatives Celasterol Cepharanthine Chlorogenic acid Chlorophyllin Cocoa polyphenols Curcumin DHEA DHEA sulfate Dehydroevodiamine Demethyltraxillagenin Diethyldithiocarbamate Diferoxamine	Tyrphostin AG-126 Ursolic acid Ligonberries Lupeol Magnolol Maltol <i>Mangifera indica</i> bark extract Mangiferin Melatonin Mn-SOD Mulberry anthocyanins Myricetin N-acetyl-L-cysteine Nacyselyn Naringin N-ethyl-maleimide Nitrosoglutathione NDGA Ochnaflavone Orthophenanthroline Phenylarsine oxide PhIP Phyllanthus urinaria PMC PTX
CaMKK CD43 overexpression FLN29 overexpression FLIP G-120 Gax (homeobox protein) HIV-1 Resistance Factor Interleukin 4 SspH1 and IpaH9.8 NDPP1 (CARD protein) Overexpressed ZIP1 p8 p202a	Fenoldopam FEX Fibrates FK778 Flunixin meglumine Flurbiprofen Hydroquinone IMD-0354 JSH-21 KT-90 Lovastatin Mercaptopyrazine Mevinolin,	Adenosine c-AMP <i>Artemisia sylvatica</i> extract Bifodobacteria Blueberry & berry mix BSASM BF phenylpropanoids cPrG.HC Seaweed extract <i>Fructus benincasae</i> extract Glucocorticoids Gypenoside XLIX Kwei Ling Ko <sup>3</sup>	Tetrathiomolybdate Tranilast Troglitazone <b>Others</b> Low gravity <b>F. Proteasome/protease</b> <b>Natural product</b> Cyclosporin A Lactacystine β-lactone <b>Peptide</b> ALLnL LLM	Dilazep Fenofibric acid DMDTC Dimethylsulfoxide Disulfiram Ebselen Edaravone EGTA EPC-K1 Epigallocatechin-3-gallate Ergothioneine Ethyl pyruvate <i>Ganoderma lucidum</i> polysaccharides Garcinol γ-glutamylcysteine synthetase <i>Ginkgo biloba</i> extract	Pyrrhithione Pyroline dithiocarbamate Quercetin Quinoxolines Rebamipide Red wine Redox factor 1 Resveratrol Ginseng derivative Rotenone Roxithromycin S-allyl-cysteine Sauchinone Spironolactone Strawberry extracts Taxifolin
p21 (Rec) PIAS1	Monoethylfumarate Moxifloxacin	LC root Luteolin	Ubiquitin ligase Z-LLL		Tempol Tepoxaline
Pro-opiomelanocortin	Nicorandil	Manassantins A,B	Z-LLnV		tert-butyl hydroquinone Tetracylic A Vitamin B6 Vitamin C
PYPAF1 protein Raf Kinase inhibitor protein <i>Rhus</i> <i>vermiciflua</i> fruits SLPI Siah2 SIRT1 Deacetylase overexpression Siva-1 <i>Solana nigrum</i> L. Surfactant protein A Tom1 overexpression Transdominant p50 Uterogloblin VEGF	Nilvadipine NO-ASA  Panepoxydone Peptide nucleic acids Perindopril  PAD α-PBN Pioglitazone Pirfenidone PNO derivatives Quinadril Raloxifene Raxofelast Ribavirin Rifamides	<i>MI</i> bark extract Mesuol  Nobiletin Phomol Psychosine  Qingkailing <sup>#</sup> Saucerneol D & E Shuanghuanglian <sup>#</sup> <i>Smilax bockii</i> extract Trilinolein <i>Uncaria tomentosum</i> extract WS extracts Wortmannin α-zearalenol <b>Viral Protein</b>	Boronic acid peptide BTEE 3,4-dichloroisocoumarin  Deoxyspergualin DFP Disulfiram FK506 (Tacrolimus) Bortezomib Salinosporamide A TLCK TPCK <b>G. Antioxidants</b> 23-hydroxyursolic acid Aged garlic extract Anetholdithiolthione	Glutathione Hematein  Hydroquinone Hydroquinone IRF1 042  Iron tetrakis Isovitexin Kangen-karyu extract Ketamine Lacidipine Lazaroids L-cysteine	Vitamin D Vitamin E derivatives Wogonin  xanthohumol Yakuchinone A, B α-lipoic acid α-tocopherol α-torophryl acetate α-torophryl succinate β-Carotene



2A and 2B-A and dephosphorylating the p65 subunit of NF- $\kappa$ B [23,76]. Recently, OspF, a protein phosphatase from *Shigella flexneri*, was found to dephosphorylate MAPK and prevent histone H3 phosphorylation at Ser<sup>10</sup> in a gene-specific manner to block the activation of a subset of NF- $\kappa$ B responsive genes [77]. Our previous studies have shown that protein tyrosine phosphatase (PTP) inhibitors can suppress NF- $\kappa$ B activation and that phenylarsine oxide, a specific PTP inhibitor, can promote tyrosine 42 phosphorylation of I $\kappa$ B $\alpha$  [78]. While some PTPs stimulate NF- $\kappa$ B activation, other PTPs negatively regulate NF- $\kappa$ B activation. For instance, PTEN, a tumor suppressor with phosphatase activity is known to inhibit NF- $\kappa$ B activation [79]. Recently, Chew et al. [80] found that WIP1, a Ser/Thr PP2C family of phosphatases, acts as a negative regulator of NF- $\kappa$ B activation. Overexpression of WIP1 was associated with decreased NF- $\kappa$ B activation, whereas WIP1 knockdown resulted in increased NF- $\kappa$ B activation. The group further showed that WIP1 target Ser<sup>536</sup> of the p65 subunit of NF- $\kappa$ B.

### 2.3. Proteasome inhibitors and I $\kappa$ B ubiquitination blockers

The step before NF- $\kappa$ B leaves the cytoplasm involves the ubiquitination of I $\kappa$ B by the SCF- $\beta$ -TrCP ubiquitin ligase complex followed by the rapid degradation of ubiquitinated I $\kappa$ B by the 26S proteasome [38]. Because I $\kappa$ B $\alpha$  degradation is an important step in the NF- $\kappa$ B activation pathway, inhibiting the proteasomes that degrade I $\kappa$ B $\alpha$  may also serve as a tool for pharmacological intervention. Very specific and potent proteasome inhibitors have been engineered by coupling boronic acid to dipeptides [81]. The dipeptide boronate, bortezomib, the most-studied proteasome inhibitor in clinical development, has been shown to inhibit proliferation and induce apoptosis in head and neck [82–84], prostate

[85], pancreatic [86], gastric [87], and ovarian [88] cancers. Bortezomib's antitumor properties correlate in part with its ability to inhibit I $\kappa$ B $\alpha$  degradation [82,89]. Other well-known proteasome inhibitors include ALLnL, LLM, Z-LLnV, and Z-LLL, lactacystine, N-cbz-Leu-Leu-leucinal (MG132), MG115, and ubiquitin ligase inhibitors [90]. In addition, we recently identified a novel proteasome inhibitor, salinosporamide A (NPI-0052), which can suppress both constitutive and inducible NF- $\kappa$ B activation in a nanomolar range [91].

Several serine protease inhibitors with chymotrypsin-like specificity, including DCIC, TPCK, TLCK, BTEE, APNE, are also able to block proteasome function. However, unlike other protease inhibitors that block only I $\kappa$ B degradation, serine protease inhibitors can block I $\kappa$ B phosphorylation as well as degradation. However, not all serine protease inhibitors can inhibit NF- $\kappa$ B activation [92–94].

Among I $\kappa$ B ubiquitination blockers, the YopJ protein of the bacterial pathogen *Yersinia* deubiquitinates and stabilizes I $\kappa$ B $\alpha$  to prevent NF- $\kappa$ B nuclear translocation [95]. The small molecule R0196-9920 has been reported to inhibit I $\kappa$ B $\alpha$  ubiquitination and oral inflammation in mouse models [96,97]. Yaron et al. [97] blocked TNF $\alpha$ -induced I $\kappa$ B $\alpha$  degradation by microinjecting phosphopeptides that corresponded to I $\kappa$ B $\alpha$ 's signal-dependent phosphorylation site. Presumably these phosphopeptides acted as competitive inhibitors for binding to the ubiquitin ligase complex essential to I $\kappa$ B $\alpha$  degradation. Inhibiting  $\beta$ -TrCP (the recognition subunit of the SCF E3 ligase complex) by specific RNAi treatment or by overexpression of dominant-negative  $\beta$ -TrCP mutants blocked NF- $\kappa$ B activity and sensitized breast cancer cells to chemotherapeutic agents [98]. Recently, A20 (TNFAIP3), a cytoplasmic zinc finger protein, was shown to inhibit NF- $\kappa$ B activation in the TNFR and TLR pathways. The ubiquitin editing property of A20 was shown to be essential for NF- $\kappa$ B inhibition [99].

#### Notes to Table 1:

15d-PGJ(2), 15-deoxy-prostaglandin J(2); 17-AAG, 17-allylamino-17-demethoxygeldanamycin; 20(S)-PPT, 20(S)-Protopanaxatriol; 4'-DM-6-Mptox, 4'-demethyl-6-methoxypodophyllotoxin; 6-MITC, 6-Methylsulfinyl hexyl isothiocyanate;  $\alpha$ -MSH,  $\alpha$ -melanocyte-stimulating hormone;  $\alpha$ -PBN, alpha-phenyl-N-tert-butyl nitron; ACHP, 2-amino-6-[2-(cyclopropylmethoxy)-6-hydroxyphenyl]-4-piperidin-4-yl nicotinonitrile; AGRO100, G-quadruplex oligodeoxynucleotide; AHUP, 8-acetoxy-5-hydroxyumbelliprenin; AIDCA, (Amino)imidazolyl carboxaldehyde derivative; ALLnL, N-acetyl-leucinyll-leucynil-norleucinal; APNE, N-acetyl-DL-phenylalanine-b-naphthylester; AQC, 6-aminoquinazoline; BAY-11-7082, E3((4-methylphenyl)-sulfonyl)-2-propenenitrile; BAY-11-7083, E3((4-t-butylphenyl)-sulfonyl)-2-propenenitrile; BF, *Bupleurum fruticosum*; BG, *Bruguiera gymnorrhiza*; BHA, Butylated hydroxyanisole; BMD, N(1)-Benzyl-4-methylbenzene-1,2-diamine; BMT, o,o'-bismyristoyl thiamine disulfide; BSASM, plant extract mixture; BTEE, N-benzoyl L-tyrosine-ethyl ester; CaMKK, Calcium/calmodulin-dependent kinase kinase; CAPE, Caffeic Acid Phenethyl Ester; CDDO-Me, C-28 methyl ester of 2-cyano-3,12-dioxooolean-1,9-dien-28-oic acid; CDS, Commercial peritoneal dialysis solution; CMP, Chinese medicinal preparations; CNA, Cinnamaldehyde; Compound 5, Uredio-thiophenecarboxamide derivative; CP Compound, 6-Hydroxy-7-methoxychroman-2-carboxylic acid phenylamide; CPC, cetylpyridinium chloride; cPrG.HC, Cycloprodigiosin hydrochloride; CSPDP, Chondrotin sulfate proteoglycan degradation product; CYL-19 s and CYL-26z, two synthetic alpha-methylene-gamma-butyrolactone derivatives; D609, phosphatidylcholine-specific phospholipase C inhibitor; DAAS, diacetoxy acetal derivative of santonin; DFP, diisopropyl fluorophosphates; DHEA, Dehydroepiandrosterone; DHMEQ, Dehydroxymethylepoxyquinomicin; DMDTC, Dimethylthiocarbamates; DPE, 2-(3,4-dihydroxyphenyl)ethanol; DTD, 4,10-dichloropyrido[5,6:4,5]thieno[3,2-d']-1,2,3-dithiazine; EGTA, Ethylene glycol tetraacetic acid; EMC virus, encephalomyocarditis virus; EMDPC, Ethyl 2-[(3-methyl-2,5-dioxo(3-pyrrolinyl)) pyrimidine-5-carboxylate; EPC-K1, phosphodiester compound of vitamin E; FEX, Fexofenadine hydrochloride; FHIT, Fragile histidine triad protein; FLIP, FLICE-Like Inhibitory Protein; G-120, Ulmus davidiana Nakai glycoprotein; GILZ, Glucorticoid-induced leucine zipper protein; HB-EGF, Heparin-binding epidermal growth factor-like growth factor; HMP, 7-(4'-hydroxy-3'-methoxyphenyl)-1-phenylhept-4-en-3-one; HP, *Harpagophytum procumbens*; HSCO, Hepatoma Subtracted-cDNA library Clone One; IQCAD, Imidazolylquinoline-carboxaldehyde derivative; JSH-23, 4-Methyl-(3-phenylpropyl)-benzene-1,2-diamine; KL-1156, 6-Hydroxy-7-methoxychroman-2-carboxylic acid phenylamide; KSHV, Kaposi's sarcoma-associated herpesvirus; KSR, Kinase suppressor of ras; LC, *Ligusticum chuanxiong*; LLM, N-acetyl-leucinyll-leucynil-methional; LY294002, [2-(4-morpholinyl)-8-phenylchromone]; MAST205, a serine/threonine kinase; MC160, Mollusum contagiosum virus; MEB, 2-(4-morpholynl) ethyl butyrate hydrochloride; mEET, Mouse estrogen enhanced transcript; Mn-SOD, Manganese superoxide dismutase; MSO, 8-methylsulphinyloctyl; N-(4-hydroxyphenyl) retinamide [4-HPR; NDGA, Nordihydroguaiaritic acid; NCPP, NLS Cell permeable peptides; NFD-37, 2-Methyl-2-(2-methylpropenyl)-2,3-dihydronaphthoquinone [2,3-b]furan-4,9-dione; NH(2)Cl, monochloramine; NO-ASA, Nitric-oxide-donating aspirin; ox-LDL, Oxidized low density lipoprotein; OXPAPC, oxidized 1-palmitoyl-2-arachidonoyl-sn-glycero-3-phosphorylcholine; PACAP, Pituitary adenylate cyclase-activating polypeptide; PAD, 6(5H)-phenanthridinone; PEDF, pigment epithelium derived factor; PEITC, Phenethyl isothiocyanate; PGG, 1,2,3,4,6-penta-O-galloyl- $\beta$ -D-glucose; PhIP, 2-Amino-1-methyl-6-phenylimidazo[4,5-b]pyridine; PI, protease inhibitor; PIAS1, protein inhibitor of activated STAT1; Plant compound A, plant-derived phenyl aziridine precursor; PMC, (2,2,5,7,8-pentamethyl-6-hydroxychromane); PMV, Paromyxovirus; PNO, Pyridine N-oxide; PQD, Pyrazolo[4,3-c]quinoline derivative; PSK, Protein-bound polysaccharide; PTEN, phosphatase and tensin homolog; PTX, Pentoxifylline (1-(5'-oxohexyl) 3,7-dimethylxanthine); PTX-B, pertussis toxin binding protein; Pyridine derivatives, 2-amino-3-cyano-4-aryl-6-(2-hydroxy-phenyl)pyridine derivatives; RH(2) & C-K, intestinal bacterial metabolites Rh(2) and compound K (C-K); ROR $\alpha$ , Retinoic acid receptor-related orphan receptor-alpha; SH, *Sargassum hemiphyllum*; SAIF, *Saccharomyces boulardii* anti-inflammatory factor; SLPI, Secretory leucoprotease inhibitor; SMI and FP, Salmeterol and Fluticasone propionate; SOCS, suppressor of cytokine signaling proteins; SspH1 and IpaH9.8, Leucine-rich effector proteins of *Salmonella* & *Shigella*; TDZD, 1,2,4-thiadiazolidine derivative; TG, triglyceride; THI 52, 1-naphthylethyl-6,7-dihydroxy-1,2,3,4-tetrahydroisoquinoline; TLCK, N- $\alpha$ -tosyl-L-lysine chloromethyl ketone; TMFC, 3,4,5-trimethoxy-4'-fluorochalcone; TNAP, TRAFs and NIK-associated protein; TP, *Tripterygium polyglycosides*; TPCA-1, 2-[(aminocarbonyl)amino]-5-acetylenyl-3-thiophenecarboxamides; TPCK, N- $\alpha$ -tosyl-L-phenylalanine chloromethyl ketone; TZD, Thiazolidinedione; VEGF, Vascular endothelial growth factor; VIP, Vasoactive intestinal peptide; VVP, Vaccinia virus protein; WS, *Witheringia solanacea*; Z-LLL, N-carbobenzoxyl-L-leucinyll-L-leucinyll-L-norleucinal; Z-LLnV (carbobenzoxyl-leucinyll-leucynil-norvalinal.

<sup>1</sup> Tibetan medicine.

<sup>2</sup> Piper methysticum.

<sup>3</sup> Tortoise shell-Rhizome jelly.

\* Anticancer drugs.

\*\* Small molecules.

\*\*\* Oriental medicines.

# Traditional Chinese Medicine.

## 2.4. Blockage of NF- $\kappa$ B nuclear translocation

One approach for inhibiting NF- $\kappa$ B activation is to use small peptides that cross the cell membrane and block the nuclear translocation of the NF- $\kappa$ B complex [100–102]. For example, SN50, a forty-one-residue synthetic peptide that contains a hydrophobic membrane-translocating region and the nuclear localization sequence of NF- $\kappa$ B p50 [100], can enter cells and compete with NF- $\kappa$ B complexes for the machinery responsible for the nuclear translocation of NF- $\kappa$ B. SN50 effectively inhibits the LPS- and TNF- $\alpha$ -induced nuclear translocation of NF- $\kappa$ B in different cell lines [100, 103–106] and mitigates inflammatory responses *in vivo* [107,108]. However, SN50 also blocks the nuclear translocation of a number of other transcription factors [101]. Dehydroxymethylepoxyquinomicin, a fungal epoxyquinoid that has anti-inflammatory and antitumor activity in several mouse models, has been reported to be a specific inhibitor of NF- $\kappa$ B nuclear translocation [109].

## 2.5. Blocking NF- $\kappa$ B activation by inhibitors of p65 acetylation

The activated p65 subunit of NF- $\kappa$ B undergoes acetylation in the nucleus at multiple lysine residues including K<sup>122</sup>, K<sup>123</sup>, K<sup>218</sup>, K<sup>221</sup> and K<sup>310</sup> [24,110]. The opposing activities of histone acetyltransferases and histone deacetylases (HDACs) regulate p65 complex acetylation [25]. Acetylation of p65 also depends on coactivators such as p300 and CREB-binding protein (CBP) [111]. The K<sup>221</sup> and K<sup>310</sup> acetylation are associated with increased NF- $\kappa$ B target gene transcription [111] and are required for p65 activation [25], which is supported by the observations that SIRT driven deacetylation at K<sup>310</sup> inhibits NF- $\kappa$ B target gene transcription [112]. Additionally, K<sup>122</sup> and K<sup>123</sup> acetylation reduces p65 DNA binding affinity and increases I $\kappa$ B interaction and nuclear export [110]. Site-specific p300-mediated p65 acetylation thus regulates the specificity of NF- $\kappa$ B-dependent gene expression [110,113].

During the last 5 years, a number of compounds have been reported to inhibit NF- $\kappa$ B by inhibiting acetylation. Gallic acid obtained from natural products such as gallnuts, sumac, oak bark, and green tea was recently reported to possess anti-histone acetyltransferase activity, thus showing potential to downregulate NF- $\kappa$ B activation [114]. Daxx, a protein associated with the death domain of Fas receptor, has been reported to suppress NF- $\kappa$ B transcriptional activity by inhibiting p300/CBP-mediated p65 acetylation [115]. Anacardic acid derived from traditional medicinal plants can also inhibit NF- $\kappa$ B activation by inhibiting p65 acetylation [116].

## 2.6. Blocking NF- $\kappa$ B activation by methyltransferases

The RelA subunits of NF- $\kappa$ B undergo various posttranslational modifications that create specific marks to recruit different effectors to control NF- $\kappa$ B's temporal and spatial activation [117]. RelA is subject to monomethylation by lysine methyltransferase Set9 (also called Set7 or KMT7) at Lys<sup>314/315</sup> *in vitro* and *in vivo* in response to stimuli [26]. RelA methylation at these two residues negatively regulates NF- $\kappa$ B function by triggering the ubiquitination and proteasome-mediated degradation of promoter-associated RelA. RelA methylation also serves as a “death” signal for the destruction of DNA-bound, activated NF- $\kappa$ B [26]. Because RelA subunit methylation negatively impacts NF- $\kappa$ B function, designing a molecule that activates Set9 function could potentiate NF- $\kappa$ B inhibition.

## 2.7. Blockage of NF- $\kappa$ B to DNA binding

The most direct strategy for blocking NF- $\kappa$ B activation is to block NF- $\kappa$ B from binding to specific  $\kappa$ B sites on DNA. Some sesquiterpene lactones (SLs) have been reported to inhibit NF- $\kappa$ B [118] by interacting with Cys-38 in the DNA-binding loop of RelA [119,120]. Most SLs can also inhibit DNA binding through an analogous Cys residue in the DNA-binding loops of p50 and c-Rel. Recently, a computer-based structural comparison of 103 SLs predicted that a methylene-carbonyl substructure is important for SL-based inhibition

of RelA at Cys-38 [121]. Some SLs, including parthenolide, have been shown to inhibit IKK $\beta$  through the reactive Cys-179 in the kinase activation loop [34,120]. Thus, SLs, which target both IKK activity and NF- $\kappa$ B subunit DNA binding [36], have multistep inhibitory activity within the NF- $\kappa$ B signaling pathway.

Blocking specific NF- $\kappa$ B-DNA binding can also be accomplished with decoy oligodeoxynucleotides (ODNs). These ODNs have  $\kappa$ B sites and competes for NF- $\kappa$ B dimer binding to specific genomic promoters [122–124]. These oligonucleotides have modifications to increase their stability and their affinity for NF- $\kappa$ B *in vivo* [125–127]. Decoy ODNs have been reported to have therapeutic potential in a number of animal models of inflammation and cancer; for example, directly injecting NF- $\kappa$ B decoy ODNs into implanted adenocarcinoma colon 26 tumors in mice inhibited cachexia without affecting tumor growth [128].

## 2.8. Other mechanisms of NF- $\kappa$ B inhibition

### 2.8.1. By gene transfer

One strategy to block NF- $\kappa$ B activation is through the transfer of genes that code for proteins shown to suppress NF- $\kappa$ B activation. The most direct target is I $\kappa$ B $\alpha$ . I $\kappa$ B $\alpha$  mutation at specific phosphorylation sites (Ser<sup>32</sup> and Ser<sup>36</sup> replaced to alanine) and ubiquitination sites (Lys<sup>21</sup> and Lys<sup>22</sup> mutated to arginine) results in a nondegradable form of I $\kappa$ B $\alpha$ . This results in a stable cytoplasmic pool of I $\kappa$ B $\alpha$ , thereby preventing NF- $\kappa$ B activation [129–131]. Injecting a nonphosphorylatable form of I $\kappa$ B $\alpha$  into bone marrow macrophages has been shown to inhibit osteoclastogenesis and block bone resorption [132]. Additionally, specific C-terminal serine-to-alanine mutations are sometimes included to reduce the constitutive turnover of I $\kappa$ B $\alpha$  [133]. These super-repressor forms of I $\kappa$ B $\alpha$  can still interact with NF- $\kappa$ B dimers to keep the dimers in the cytoplasm permanently [131,133,134]. Such molecules have been used successfully to inhibit NF- $\kappa$ B activity and to study its role in tumor development [135,136] and to sensitize tumor cells to apoptosis-inducing agents [133,134]. Inhibiting NF- $\kappa$ B through the expression of an I $\kappa$ B $\alpha$  super-repressor (I $\kappa$ B $\alpha$ SR) has also been used to sensitize chemoresistant tumors to TNF $\alpha$ - and CPT-11-induced apoptosis, resulting in tumor regression [137], and to inhibit the proliferation of human head and neck carcinoma cells *in vitro* and *in vivo* [138]. However, I $\kappa$ B $\alpha$ SRs have also been shown to interact with and affect the activity of non-NF- $\kappa$ B pathway proteins including p53 [139], cyclin-dependent kinase 4 [140], and HDACs [141]. Furthermore, I $\kappa$ B $\alpha$ SR overexpression has been associated with the spontaneous development of squamous cell carcinoma in a murine model [142].

### 2.8.2. Antioxidants

Antioxidants were suggested as possible NF- $\kappa$ B inhibitors many years ago [143,144]. Treatment with oxidants such as hydrogen peroxide can activate NF- $\kappa$ B in many cell types. In some cell types, antioxidants can inhibit the induction of NF- $\kappa$ B activity in response to a variety of stimuli (e.g., interleukin-1 $\beta$ , LPS, TNF $\alpha$ ) [145,146]. However, using antioxidants as NF- $\kappa$ B inhibitors is now regarded with increasing scepticism because the NF- $\kappa$ B-inhibiting properties of pyrrolidine dithiocarbamate, a thiol-containing compound, cannot be attributed to its antioxidant function but rather to its effects as an inhibitor of I $\kappa$ B ubiquitin ligase activity [147]. The ways in which antioxidants block NF- $\kappa$ B activation remain unclear, but it is likely that they act at different steps in the NF- $\kappa$ B pathway in different cell types. Antioxidants have been suggested to inhibit NF- $\kappa$ B activation by scavenging reactive oxygen intermediates that act as signaling molecules to activate the NF- $\kappa$ B pathway and by directly inhibiting IKK kinase activity by modifying critical Cys residues in the IKK kinase activation loop [145,146]. Mitochondrial electron transport inhibitors that suppress reactive oxygen intermediate production (e.g., rotenone) and overexpression of antioxidantizing enzymes (e.g., manganese superoxide dismutase and catalase) can block TNF $\alpha$ -induced NF- $\kappa$ B activation [148–150]. Caffeic acid phenethyl ester, a phenolic antioxidant, has been



reported to cause direct interference with DNA binding by NF- $\kappa$ B [151] that can be reversed by dithiothreitol [78]. Other antioxidants, viz., N-acetylcysteine, calcium chelators (e.g., EGTA, lacidipine), and vitamin C and E derivatives have been reported to inhibit hydrogen peroxide- or stimulus-induced NF- $\kappa$ B activation.

### 2.8.3. Bacterial, fungal, and viral proteins

Several microorganisms and viruses encode proteins that can inhibit NF- $\kappa$ B activation. Many viruses have developed a number of mechanisms to inhibit NF- $\kappa$ B signaling [152], and three viruses—African swine fever virus (ASFV) [153], rabbit myxoma virus [154], and insect *Microplitis demolitor* bracovirus [155]—encode I $\kappa$ B-like NF- $\kappa$ B inhibitors. The ASFV encodes the A238L I $\kappa$ B-like protein, which can stably interact with RelA to inhibit TNF $\alpha$ -, IFN- $\gamma$ -, and phorbol ester-induced NF- $\kappa$ B–DNA binding [156]. The poliovirus 3C protease cleaves RelA to reduce NF- $\kappa$ B signaling [157]. In addition, several viruses have adaptor-like or small proteins that inhibit IKK activity [152]. For example, the MC160 protein of *Molluscum contagiosum* [158] and the nonstructural 5B protein of the hepatitis C virus [159] appear to be IKK $\alpha$ -specific and thus may specifically inhibit the noncanonical NF- $\kappa$ B pathway.

The YopJ protein, a Src homology 2 domain protein encoded by *Yersinia pseudotuberculosis*, inhibits NF- $\kappa$ B activation by preventing the phosphorylation and degradation of I $\kappa$ B $\alpha$  [160]. YopJ has also been shown to bind directly to IKK $\beta$  *in vitro* and *in vivo* [161]. The *Salmonella typhimurium* AvrA protein also inhibits NF- $\kappa$ B activation, although its mechanism of action may be different than that of the YopJ protein [162].

Glutoxin produced by the fungus *Aspergillus fumigatus* has been reported to inhibit NF- $\kappa$ B activation by preventing I $\kappa$ B degradation [163]. Several other small molecules synthesized by microorganisms or designed derivatives of such compounds that have NF- $\kappa$ B-inhibiting potential include panepoxydone (from *Lentinus crinitus*) [164], 5,6 epoxycyclohexenone compounds (from *Amycolatopsis*), and cycloepoxydon [165]. Such compounds may affect distinct parts of the NF- $\kappa$ B pathway including DNA binding, nuclear translocation, and I $\kappa$ B $\alpha$  phosphorylation and degradation.

### 2.8.4. Anti-inflammatory and immunosuppressive agents

Various anti-inflammatory agents including glucocorticoids, non-steroid anti-inflammatory drugs (NSAIDs), and immunosuppressants have been developed to block NF- $\kappa$ B activation. Glucocorticoids, which are commonly used as anti-inflammatory drugs, strongly inhibit NF- $\kappa$ B activation by mechanisms that are not completely understood but likely include inhibition of DNA binding, IKK activity and transactivation [166]. The glucocorticoids dexamethasone, prednisone and methylprednisolone have been reported to inhibit NF- $\kappa$ B activation. In addition, estrogen and selective estrogen receptor modulators (SERMs) such as raloxifene can act through the estrogen receptor to inhibit NF- $\kappa$ B activation [167,168].

NSAIDs such as sodium salicylate (aspirin) and sulindac have been reported to inhibit NF- $\kappa$ B activation by inhibiting I $\kappa$ B $\alpha$  phosphorylation [169,170]. At higher concentrations, aspirin has been shown to block NF- $\kappa$ B activity by directly binding to and inhibiting the kinase activity of IKK $\beta$  by reducing its ability to bind ATP [171]. More recently, aspirin was reported to inhibit proteasome activity [172]. As such, high-dose aspirin therapy may have applications in treating diseases in which NF- $\kappa$ B activity is involved, including cancer [173], diabetes [174], and heart disease [175]. Other NSAIDs such as ibuprofen and indomethacin have also been reported to inhibit NF- $\kappa$ B activation in cell culture [176–179].

Several well known immunosuppressants are known to target NF- $\kappa$ B by distinct mechanisms, some precluding NF- $\kappa$ B nuclear translocation [180], some through inhibiting calcineurin [181], some by binding heat-shock proteins [182] and some by modulating the DNA binding or transactivation potential of NF- $\kappa$ B [183–186]. Examples of immunosuppressants having inhibitory effect on NF- $\kappa$ B activation include cyclosporin A (CsA) [180], FK506 [187,188], PG490 (diterpene triepoxide) [186] and deoxyspergualin [182].

### 2.8.5. p53 induction

It is known that p53 and NF- $\kappa$ B pathways play opposing roles in human cancer, with p53 acting as a tumor suppressor and NF- $\kappa$ B acting as a tumor activator. The crosstalk between p53 and NF- $\kappa$ B indicates that p53 and NF- $\kappa$ B repress each other's activities owing to their competition for transcriptional coactivator proteins p300 and CBP [189]. A recent study has proposed an additional mechanism of how CBP phosphorylation by IKK $\alpha$  determines whether CBP binds to p53 or NF- $\kappa$ B [190]. Although a number of studies have focused on identifying p53 activators and NF- $\kappa$ B inhibitors individually, few studies have investigated the molecules that target both the pathways simultaneously. Identifying molecules that simultaneously activate p53 and inhibit NF- $\kappa$ B would have great potential in combination therapy for cancer and various other diseases and could provide helpful tools to better understand the crosstalk between the p53 and NF- $\kappa$ B pathways. Quinacrine, an antimalarial drug, and other derivatives of 9 aminoacridine have been shown to simultaneously repress NF- $\kappa$ B and activate p53 in renal cell carcinoma [191]. Other molecules with similar potential include nutlins [192,193], cisplatin [194,195], leptomycin B [196,197], adenosine-2,3-dialdehyde [198], the NSAID JTE-522 [199], and the cyclin-dependent kinase inhibitors R-roscovitine [200,201]; and flavopiridol [202,203]. While most reports indicate antiapoptotic role of NF- $\kappa$ B, some suggest proapoptotic role of this transcription factor depending on the stimulus, cell type and subunit involved [204–206]. In addition, under certain circumstances, the apoptosis induced by p53 may involve the activation of NF- $\kappa$ B [207,208]. Thus, inhibition of NF- $\kappa$ B in tumors that exhibit wild type p53 may diminish, rather than augment a therapeutic response.

## 3. Conclusions and future perspective

NF- $\kappa$ B has been implicated in almost all chronic diseases, and more than 40,000 studies on NF- $\kappa$ B have been published with 9000 on its inhibitors. Although more than 700 different inhibitors (aspirin to I $\kappa$ B $\alpha$  super repressor) of this transcription factor have been reported, yet no NF- $\kappa$ B blocker has been approved for human use. Various steroids and NSAIDs have been found to block NF- $\kappa$ B, but their effects are highly pleiotropic. The molecules that block NF- $\kappa$ B activation lack specificity and thus interfere with NF- $\kappa$ B's physiological roles in immunity, inflammation, and cellular homeostasis. Additionally, whether the concentrations of inhibitors used in tissue culture experiments can be applied *in vivo* is often unclear. Therefore, one of the major challenges facing researchers is to develop NF- $\kappa$ B inhibitors aimed at treating different diseases based on their ability to target specific pathways or cells, thereby avoiding the risk of undesired side effects. Future studies should also focus on validating *in vitro* data *in vivo*.

## Acknowledgments

We thank Joe Munch for carefully proof-reading the manuscript and providing valuable comments. Dr. Aggarwal is the Ransom Horne, Jr., Professor of Cancer Research. This work was supported by a grant from the Clayton Foundation for Research (B.B.A.), a core grant from the National Institutes of Health (CA-16 672), a program project grant from National Institutes of Health (NIH CA-124787-01A2), and grant from Center for Targeted Therapy of M.D. Anderson Cancer Center.

## References

- [1] H.L. Pahl, Activators and target genes of Rel/NF-kappaB transcription factors, *Oncogene* 18 (1999) 6853–6866.
- [2] M. Karin, Y. Cao, F.R. Greten, Z.W. Li, NF-kappaB in cancer: from innocent bystander to major culprit, *Nat. Rev. Cancer* 2 (2002) 301–310.
- [3] B. Haefner, NF-kappa B: arresting a major culprit in cancer, *Drug Discov. Today* 7 (2002) 653–663.

- [4] A. Richmond, NF-kappa B, chemokine gene transcription and tumour growth, *Nat. Rev. Immunol.* 2 (2002) 664–674.
- [5] B.B. Aggarwal, Nuclear factor-kappaB: the enemy within, *Cancer Cell* 6 (2004) 203–208.
- [6] R. Sen, D. Baltimore, Multiple nuclear factors interact with the immunoglobulin enhancer sequences, *Cell* 46 (1986) 705–716.
- [7] H. Clevers, At the crossroads of inflammation and cancer, *Cell* 118 (2004) 671–674.
- [8] F. Balkwill, K.A. Charles, A. Mantovani, Smoldering and polarized inflammation in the initiation and promotion of malignant disease, *Cancer Cell* 7 (2005) 211–217.
- [9] M. Karin, F.R. Greten, NF-kappaB: linking inflammation and immunity to cancer development and progression, *Nat. Rev. Immunol.* 5 (2005) 749–759.
- [10] Q. Li, S. Withoff, I.M. Verma, Inflammation-associated cancer: NF-kappaB is the lynchpin, *Trends Immunol.* 26 (2005) 318–325.
- [11] J.L. Luo, H. Kamata, M. Karin, IKK/NF-kappaB signaling: balancing life and death—a new approach to cancer therapy, *J. Clin. Invest.* 115 (2005) 2625–2632.
- [12] A.L. Shaffer, A. Rosenwald, L.M. Staudt, Lymphoid malignancies: the dark side of B-cell differentiation, *Nat. Rev. Immunol.* 2 (2002) 920–932.
- [13] A. Panwalkar, S. Verstovsek, F. Giles, Nuclear factor-kappaB modulation as a therapeutic approach in hematologic malignancies, *Cancer* 100 (2004) 1578–1589.
- [14] S. Shishodia, B.B. Aggarwal, Nuclear factor-kappaB activation mediates cellular transformation, proliferation, invasion angiogenesis and metastasis of cancer, *Cancer Treat. Res.* 119 (2004) 139–173.
- [15] S. Ghosh, M.J. May, E.B. Kopp, NF-kappa B and Rel proteins: evolutionarily conserved mediators of immune responses, *Annu. Rev. Immunol.* 16 (1998) 225–260.
- [16] S. Ghosh, M. Karin, Missing pieces in the NF-kappaB puzzle, *Cell* 109 (Suppl) (2002) S81–96.
- [17] M.S. Hayden, S. Ghosh, Signaling to NF-kappaB, *Genes Dev.* 18 (2004) 2195–2224.
- [18] U. Sentfelen, Y. Cao, G. Xiao, F.R. Greten, G. Krahn, G. Bonizzi, Y. Chen, Y. Hu, A. Fong, S.C. Sun, M. Karin, Activation by IKKalpha of a second, evolutionary conserved NF-kappa B signaling pathway, *Science* 293 (2001) 1495–1499.
- [19] E. Dejardin, N.M. Droin, M. Delhase, E. Haas, Y. Cao, C. Makris, Z.W. Li, M. Karin, C.F. Ware, D.R. Green, The lymphotoxin-beta receptor induces different patterns of gene expression via two NF-kappaB pathways, *Immunity* 17 (2002) 525–535.
- [20] E. Dejardin, The alternative NF-kappaB pathway from biochemistry to biology: pitfalls and promises for future drug development, *Biochem. Pharmacol.* 72 (2006) 1161–1179.
- [21] H. Zhong, H. SuYang, H. Erdjument-Bromage, P. Tempst, S. Ghosh, The transcriptional activity of NF-kappaB is regulated by the IkappaB-associated PKAc subunit through a cyclic AMP-independent mechanism, *Cell* 89 (1997) 413–424.
- [22] D. Wang, S.D. Westerheide, J.L. Hanson, A.S. Baldwin Jr., Tumor necrosis factor alpha-induced phosphorylation of RelA/p65 on Ser529 is controlled by casein kinase II, *J. Biol. Chem.* 275 (2000) 32592–32597.
- [23] J. Yang, G.H. Fan, B.E. Wadzinski, H. Sakurai, A. Richmond, Protein phosphatase 2A interacts with and directly dephosphorylates RelA, *J. Biol. Chem.* 276 (2001) 47828–47833.
- [24] L. Chen, W. Fischle, E. Verdin, W.C. Greene, Duration of nuclear NF-kappaB action regulated by reversible acetylation, *Science* 293 (2001) 1653–1657.
- [25] L.F. Chen, Y. Mu, W.C. Greene, Acetylation of RelA at discrete sites regulates distinct nuclear functions of NF-kappaB, *EMBO J.* 21 (2002) 6539–6548.
- [26] X.D. Yang, B. Huang, M. Li, A. Lamb, N.L. Kelleher, L.F. Chen, Negative regulation of NF-kappaB action by Set9-mediated lysine methylation of the RelA subunit, *EMBO J.* 28 (2009) 1055–1066.
- [27] G. Sethi, B. Sung, B.B. Aggarwal, Nuclear factor-kappaB activation: from bench to bedside, *Exp. Biol. Med.* (Maywood) 233 (2008) 21–31.
- [28] M. Karin, Y. Yamamoto, Q.M. Wang, The IKK NF-kappa B system: a treasure trove for drug development, *Nat. Rev. Drug Discov.* 3 (2004) 17–26.
- [29] K. Nagashima, V.G. Sasseville, D. Wen, A. Bielecki, H. Yang, C. Simpson, E. Grant, M. Hepperle, G. Harriman, B. Jaffee, T. Ocain, Y. Xu, C.C. Fraser, Rapid TNFR1-dependent lymphocyte depletion in vivo with a selective chemical inhibitor of IKKbeta, *Blood* 107 (2006) 4266–4273.
- [30] M. Karin, How NF-kappaB is activated: the role of the IkappaB kinase (IKK) complex, *Oncogene* 18 (1999) 6867–6874.
- [31] V. Pande, M.J. Ramos, NF-kappaB in human disease: current inhibitors and prospects for de novo structure based design of inhibitors, *Curr. Med. Chem.* 12 (2005) 357–374.
- [32] J.R. Burke, M.A. Pattoli, K.R. Gregor, P.J. Brassil, J.F. MacMaster, K.W. McIntyre, X. Yang, V.S. Iotzova, W. Clarke, J. Strnad, Y. Qiu, F.C. Zusi, BMS-345541 is a highly selective inhibitor of I kappa B kinase that binds at an allosteric site of the enzyme and blocks NF-kappa B-dependent transcription in mice, *J. Biol. Chem.* 278 (2003) 1450–1456.
- [33] P. Kapahi, T. Takahashi, G. Natoli, S.R. Adams, Y. Chen, R.Y. Tsien, M. Karin, Inhibition of NF-kappa B activation by arsenite through reaction with a critical cysteine in the activation loop of Ikappa B kinase, *J. Biol. Chem.* 275 (2000) 36062–36066.
- [34] B.H. Kwok, B. Koh, M.I. Ndubuisi, M. Elofsson, C.M. Crews, The anti-inflammatory natural product parthenolide from the medicinal herb Feverfew directly binds to and inhibits IkappaB kinase, *Chem. Biol.* 8 (2001) 759–766.
- [35] M.C. Liang, S. Bardhan, C. Li, E.A. Pace, J.A. Porco Jr., T.D. Gilmore, Jesterone dimer, a synthetic derivative of the fungal metabolite jesterone, blocks activation of transcription factor nuclear factor kappaB by inhibiting the inhibitor of kappaB kinase, *Mol. Pharmacol.* 64 (2003) 123–131.
- [36] M.C. Liang, S. Bardhan, E.A. Pace, D. Rosman, J.A. Beutler, J.A. Porco Jr., T.D. Gilmore, Inhibition of transcription factor NF-kappaB signaling proteins IKKbeta and p65 through specific cysteine residues by epoxyquinone A monomer: correlation with its anti-cancer cell growth activity, *Biochem. Pharmacol.* 71 (2006) 634–645.
- [37] N.D. Perkins, Post-translational modifications regulating the activity and function of the nuclear factor kappa B pathway, *Oncogene* 25 (2006) 6717–6730.
- [38] C. Scheidereit, IkappaB kinase complexes: gateways to NF-kappaB activation and transcription, *Oncogene* 25 (2006) 6685–6705.
- [39] J.A. DiDonato, M. Hayakawa, D.M. Rothwarf, E. Zandi, M. Karin, A cytokine-responsive IkappaB kinase that activates the transcription factor NF-kappaB, *Nature* 388 (1997) 548–554.
- [40] F. Mercurio, H. Zhu, B.W. Murray, A. Shevchenko, B.L. Bennett, J. Li, D.B. Young, M. Barbosa, M. Mann, A. Manning, A. Rao, IKK-1 and IKK-2: cytokine-activated IkappaB kinases essential for NF-kappaB activation, *Science* 278 (1997) 860–866.
- [41] C.H. Regnier, H.Y. Song, X. Gao, D.V. Goeddel, Z. Cao, M. Rothe, Identification and characterization of an IkappaB kinase, *Cell* 90 (1997) 373–383.
- [42] J.D. Woronicz, X. Gao, Z. Cao, M. Rothe, D.V. Goeddel, IkappaB kinase-beta: NF-kappaB activation and complex formation with IkappaB kinase-alpha and NIK, *Science* 278 (1997) 866–869.
- [43] E. Zandi, D.M. Rothwarf, M. Delhase, M. Hayakawa, M. Karin, The IkappaB kinase complex (IKK) contains two kinase subunits IKKalpha and IKKbeta, necessary for IkappaB phosphorylation and NF-kappaB activation, *Cell* 91 (1997) 243–252.
- [44] Y. Shikama, M. Yamada, T. Miyashita, Caspase-8 and caspase-10 activate NF-kappaB through RIP, NIK and IKKalpha kinases, *Eur. J. Immunol.* 33 (2003) 1998–2006.
- [45] D.H. Broide, T. Lawrence, T. Doherty, J.Y. Cho, M. Miller, K. McElwain, S. McElwain, M. Karin, Allergen-induced peribronchial fibrosis and mucus production mediated by IkappaB kinase beta-dependent genes in airway epithelium, *Proc. Natl. Acad. Sci. U. S. A.* 102 (2005) 17723–17728.
- [46] M.C. Jateley, J.E. Chivers, N.S. Holden, P.J. Barnes, R. Newton, Validation of IKK beta as therapeutic target in airway inflammatory disease by adenoviral-mediated delivery of dominant-negative IKK beta to pulmonary epithelial cells, *Br. J. Pharmacol.* 145 (2005) 114–122.
- [47] M.J. May, F. D'Acquisto, L.A. Madge, J. Glockner, J.S. Pober, S. Ghosh, Selective inhibition of NF-kappaB activation by a peptide that blocks the interaction of NEMO with the IkappaB kinase complex, *Science* 289 (2000) 1550–1554.
- [48] R. Battistutta, S. Sarno, E. De Moliner, E. Papinutto, G. Zanotti, L.A. Pinna, The replacement of ATP by the competitive inhibitor emodin induces conformational modifications in the catalytic site of protein kinase CK2, *J. Biol. Chem.* 275 (2000) 29618–29622.
- [49] F. Demarchi, C. Bertoli, P. Sandy, C. Schneider, Glycogen synthase kinase-3 beta regulates NF-kappa B1/p105 stability, *J. Biol. Chem.* 278 (2003) 39583–39590.
- [50] R.F. Schwabe, D.A. Brenner, Role of glycogen synthase kinase-3 in TNF-alpha-induced NF-kappaB activation and apoptosis in hepatocytes, *Am. J. Physiol.* 283 (2002) G204–211.
- [51] F. Fujita, Y. Taniguchi, T. Kato, Y. Narita, A. Furuya, T. Ogawa, H. Sakurai, T. Joh, M. Itoh, M. Delhase, M. Karin, M. Nakanishi, Identification of NAP1, a regulatory subunit of IkappaB kinase-related kinases that potentiates NF-kappaB signaling, *Mol. Cell. Biol.* 23 (2003) 7780–7793.
- [52] L. Vermeulen, G. De Wilde, P. Van Damme, W. Vanden Berghe, G. Haegeman, Transcriptional activation of the NF-kappaB p65 subunit by mitogen- and stress-activated protein kinase-1 (MSK1), *EMBO J.* 22 (2003) 1313–1324.
- [53] X. Jiang, N. Takahashi, N. Matsui, T. Tetsuka, T. Okamoto, The NF-kappa B activation in lymphotoxin beta receptor signaling depends on the phosphorylation of p65 at serine 536, *J. Biol. Chem.* 278 (2003) 919–926.
- [54] I. Mattioli, A. Sebald, C. Bucher, R.P. Charles, H. Nakano, T. Doi, M. Kracht, M.L. Schmitz, Transient and selective NF-kappa B p65 serine 536 phosphorylation induced by T cell costimulation is mediated by I kappa B kinase beta and controls the kinetics of p65 nuclear import, *J. Immunol.* 172 (2004) 6336–6344.
- [55] N. Sizemore, S. Leung, G.R. Stark, Activation of phosphatidylinositol 3-kinase in response to interleukin-1 leads to phosphorylation and activation of the NF-kappaB p65/RelA subunit, *Mol. Cell. Biol.* 19 (1999) 4798–4805.
- [56] L.V. Madrid, M.W. Mayo, J.Y. Reuther, A.S. Baldwin Jr., Akt stimulates the transactivation potential of the RelA/p65 Subunit of NF-kappa B through utilization of the Ikappa B kinase and activation of the mitogen-activated protein kinase p38, *J. Biol. Chem.* 276 (2001) 18934–18940.
- [57] N. Sizemore, N. Lerner, N. Dombrowski, H. Sakurai, G.R. Stark, Distinct roles of the Ikappa B kinase alpha and beta subunits in liberating nuclear factor kappa B (NF-kappa B) from Ikappa B and in phosphorylating the p65 subunit of NF-kappa B, *J. Biol. Chem.* 277 (2002) 3863–3869.
- [58] W. Vanden Berghe, S. Plaisance, E. Boone, K. De Bosscher, M.L. Schmitz, W. Fiers, G. Haegeman, p38 and extracellular signal-regulated kinase mitogen-activated protein kinase pathways are required for nuclear factor-kappaB p65 transactivation mediated by tumor necrosis factor, *J. Biol. Chem.* 273 (1998) 3285–3290.
- [59] K.H. Bhatt, R.K. Pandey, Y. Dahiya, A. Sodhi, Protein kinase Cdelta and protein tyrosine kinase regulate peptidoglycan-induced nuclear factor-kappaB activation and inducible nitric oxide synthase expression in mouse peritoneal macrophages in vitro, *Mol. Immunol.* 47 (2010) 861–870.
- [60] H. Shimada, L.E. Rajagopalan, RHO kinase-2 activation in human endothelial cells drives LPA-mediated expression of cell adhesion molecules via NF-(kappa)B p65, *J. Biol. Chem.* 285 (2010) 12536–12542.
- [61] W. Sun, N. Ge, Y. Yu, S. Burlingame, X. Li, M. Zhang, S. Ye, S. Fu, J. Yang, Phosphorylation of Thr-516 and Ser-520 in the kinase activation loop of MEKK3 is required for lysophosphatidic acid-mediated optimal I(kappa)B kinase beta (IKK(beta))/nuclear factor-(kappa)B (NF-(kappa)B) activation, *J. Biol. Chem.* 285 (2010) 7911–7918.
- [62] R. Ravi, A. Bedi, NF-kappaB in cancer—a friend turned foe, *Drug Resist. Updat.* 7 (2004) 53–67.

- [63] G. Sanchez-Duffhues, M.A. Calzado, A.G. de Vinuesa, G. Appendino, B.L. Fiebig, U. Looch, A. Lefarth-Risse, K. Krohn, E. Munoz, Denbinobin inhibits nuclear factor-kappaB and induces apoptosis via reactive oxygen species generation in human leukemic cells, *Biochem. Pharmacol.* 77 (2009) 1401–1409.
- [64] D. Martin, A. Daulny, M. Decoville, D. Locker, Mutagenesis analysis of the interaction between the dorsal rel homology domain and HMG boxes of DSP1 protein, *J. Biochem.* 134 (2003) 583–589.
- [65] F. Domagala, G. Martin, P. Bogdanowicz, H. Fichoux, J.P. Pujol, Inhibition of interleukin-1beta-induced activation of MEK/ERK pathway and DNA binding of NF-kappaB and AP-1: potential mechanism for Diacerein effects in osteoarthritis, *Biorheology* 43 (2006) 577–587.
- [66] Y. Hu, H.F. Wang, W.Q. Sun, C.S. Xie, W.N. Wei, J.E. Zheng, J.X. Yao, Regulation of tissue factor expression in brain microvascular endothelial cells by PLA nanoparticles coating NF-kappaB decoy oligonucleotides, *Zhonghua Xue Ye Xue Za Zhi* 26 (2005) 534–538.
- [67] S.K. Manna, C. Bueso-Ramos, F. Alvarado, B.B. Aggarwal, Calagualine inhibits nuclear transcription factors-kappaB activated by various inflammatory and tumor promoting agents, *Cancer Lett.* 190 (2003) 171–182.
- [68] E.K. Go, K.J. Jung, J.M. Kim, H. Lim, H.K. Lim, B.P. Yu, H.Y. Chung, Betaine modulates age-related NF-kappaB by thiol-enhancing action, *Biol. Pharm. Bull.* 30 (2007) 2244–2249.
- [69] H. Kamiyama, T. Usui, H. Sakurai, M. Shoji, Y. Hayashi, H. Kakeya, H. Osada, Epoxyquinol B, a naturally occurring pentaketide dimer, inhibits NF-kappaB signaling by crosslinking TAK1, *Biosci. Biotechnol. Biochem.* 72 (2008) 1894–1900.
- [70] R. Gede, X.L. Jin, O. Hinthong, J.L. Shisler, Poxviral regulation of the host NF-kappaB response: the vaccinia virus M2L protein inhibits induction of NF-kappaB activation via an ERK2 pathway in virus-infected human embryonic kidney cells, *J. Virol.* 80 (2006) 8676–8685.
- [71] O. Hinthong, X.L. Jin, J.L. Shisler, Characterization of wild-type and mutant vaccinia virus M2L proteins' abilities to localize to the endoplasmic reticulum and to inhibit NF-kappaB activation during infection, *Virology* 373 (2008) 248–262.
- [72] Y. Li, N. Chu, A. Hu, B. Gran, A. Rostami, G.X. Zhang, Inducible IL-23p19 expression in human microglia via p38 MAPK and NF-kappaB signal pathways, *Exp. Mol. Pathol.* 84 (2008) 1–8.
- [73] P.L. Channavajhala, V.R. Rao, V. Spaulding, L.L. Lin, Y.G. Zhang, hKSR-2 inhibits MEK3-activated MAP kinase and NF-kappaB pathways in inflammation, *Biochem. Biophys. Res. Commun.* 334 (2005) 1214–1218.
- [74] B. Feng, S. Cheng, W.S. Pear, H.C. Liou, NF-kB inhibitor blocks B cell development at two checkpoints, *Med. Immunol.* 3 (2004) 1.
- [75] S. Barisic, E. Strozzyk, N. Peters, H. Walczak, D. Kulms, Identification of PP2A as a crucial regulator of the NF-kappaB feedback loop: its inhibition by UVB turns NF-kappaB into a pro-apoptotic factor, *Cell Death Differ.* 15 (2008) 1681–1690.
- [76] Y. Sreenivasan, A. Sarkar, S.K. Manna, Mechanism of cytosine arabinoside-mediated apoptosis: role of Rel A (p65) dephosphorylation, *Oncogene* 22 (2003) 4356–4369.
- [77] L. Arbi, D.W. Kim, E. Batsche, T. Pedron, B. Mateescu, C. Muchardt, C. Parsot, P.J. Sansonetti, An injected bacterial effector targets chromatin access for transcription factor NF-kappaB to alter transcription of host genes involved in immune responses, *Nat. Immunol.* 8 (2007) 47–56.
- [78] S. Singh, B.B. Aggarwal, Protein-tyrosine phosphatase inhibitors block tumor necrosis factor-dependent activation of the nuclear transcription factor NF-kappa B, *J. Biol. Chem.* 270 (1995) 10631–10639.
- [79] D. Koul, Y. Yao, J.L. Abbruzzese, W.K. Yung, S.A. Reddy, Tumor suppressor MMAC/PTEN inhibits cytokine-induced NFkappaB activation without interfering with the IkappaB degradation pathway, *J. Biol. Chem.* 276 (2001) 11402–11408.
- [80] J. Chew, S. Biswas, S. Shreeram, M. Humaidi, E.T. Wong, M.K. Dhillon, H. Teo, A. Hazra, C.C. Fang, E. Lopez-Collazo, D.V. Bulavin, V. Tergaonkar, WIP1 phosphatase is a negative regulator of NF-kappaB signalling, *Nat. Cell Biol.* 11 (2009) 659–666.
- [81] M. Iqbal, S. Chatterjee, J.C. Kauer, M. Das, P. Messina, B. Freed, W. Biazio, R. Siman, Potent inhibitors of proteasome, *J. Med. Chem.* 38 (1995) 2276–2277.
- [82] J.B. Sunwoo, Z. Chen, G. Dong, N. Yeh, C. Crowl, Bancroft, E. Sausville, J. Adams, P. Elliott, C. Van Waes, Novel proteasome inhibitor PS-341 inhibits activation of nuclear factor-kappa B, cell survival, tumor growth, and angiogenesis in squamous cell carcinoma, *Clin. Cancer Res.* 7 (2001) 1419–1428.
- [83] M. Lun, P.L. Zhang, P.K. Pellitteri, A. Law, T.L. Kennedy, R.E. Brown, Nuclear factor-kappaB pathway as a therapeutic target in head and neck squamous cell carcinoma: pharmaceutical and molecular validation in human cell lines using Velcade and siRNA/NF-kappaB, *Ann. Clin. Lab. Sci.* 35 (2005) 251–258.
- [84] C. Allen, K. Saigal, L. Nottingham, P. Arun, Z. Chen, C. Van Waes, Bortezomib-induced apoptosis with limited clinical response is accompanied by inhibition of canonical but not alternative nuclear factor-kappaB subunits in head and neck cancer, *Clin. Cancer Res.* 14 (2008) 4175–4185.
- [85] S. Williams, C. Pettaway, R. Song, C. Papandreou, C. Logothetis, D.J. McConkey, Differential effects of the proteasome inhibitor bortezomib on apoptosis and angiogenesis in human prostate tumor xenografts, *Mol. Cancer Ther.* 2 (2003) 835–843.
- [86] S.T. Nawrocki, J.S. Carew, M.S. Pino, R.A. Highshaw, R.H. Andtbacka, K. Dunner Jr., A. Pal, W.G. Bornmann, P.J. Chiao, P. Huang, H. Xiong, J.L. Abbruzzese, D.J. McConkey, Aggressive disruption: a novel strategy to enhance bortezomib-induced apoptosis in pancreatic cancer cells, *Cancer Res.* 66 (2006) 3773–3781.
- [87] M. Adachi, Y. Zhang, X. Zhao, T. Minami, R. Kawamura, Y. Hinoda, K. Imai, Synergistic effect of histone deacetylase inhibitors FK228 and m-carboxycinnamic acid bis-hydroxamide with proteasome inhibitors PSI and PS-341 against gastrointestinal adenocarcinoma cells, *Clin. Cancer Res.* 10 (2004) 3853–3862.
- [88] H. Zhu, L. Zhang, F. Dong, W. Guo, S. Wu, F. Teraishi, J.J. Davis, P.J. Chiao, B. Fang, Btk/Nbk accumulation correlates with apoptosis-induction by bortezomib (PS-341, Velcade) and other proteasome inhibitors, *Oncogene* 24 (2005) 4993–4999.
- [89] M. Lun, P.L. Zhang, N. Siegelmann-Danieli, T.M. Blasick, R.E. Brown, Intracellular inhibitory effects of Velcade correlate with morphoproteomic expression of phosphorylated-nuclear factor-kappaB and p53 in breast cancer cell lines, *Ann. Clin. Lab. Sci.* 35 (2005) 15–24.
- [90] Y. Ishii, S. Waxman, D. Germain, Targeting the ubiquitin–proteasome pathway in cancer therapy, *Anticancer Agents Med. Chem.* 7 (2007) 359–365.
- [91] K.S. Ahn, G. Sethi, T.H. Chao, S.T. Neuteboom, M.M. Chaturvedi, M.A. Palladino, A. Younes, B.B. Aggarwal, Salinosporamide A (NPI-0052) potentiates apoptosis, suppresses osteoclastogenesis, and inhibits invasion through down-modulation of NF-kappaB regulated gene products, *Blood* 110 (2007) 2286–2295.
- [92] M. Higuchi, S. Singh, H. Chan, B.B. Aggarwal, Protease inhibitors differentially regulate tumor necrosis factor-induced apoptosis, nuclear factor-kappa B activation, cytotoxicity, and differentiation, *Blood* 86 (1995) 2248–2256.
- [93] F. D'Acquisto, L. Sautebin, T. Iuvone, M. Di Rosa, R. Carnuccio, Prostaglandins prevent inducible nitric oxide synthase protein expression by inhibiting nuclear factor-kappaB activation in J774 macrophages, *FEBS Lett.* 440 (1998) 76–80.
- [94] A. Rossi, G. Elia, M.G. Santoro, Activation of the heat shock factor 1 by serine protease inhibitors. An effect associated with nuclear factor-kappaB inhibition, *J. Biol. Chem.* 273 (1998) 16446–16452.
- [95] H. Zhou, D.M. Monack, N. Kayagaki, I. Wertz, J. Yin, B. Wolf, V.M. Dixit, Yersinia virulence factor YopJ acts as a deubiquitinase to inhibit NF-kappa B activation, *J. Exp. Med.* 202 (2005) 1327–1332.
- [96] D.C. Swinney, Y.Z. Xu, L.E. Scarafia, I. Lee, A.Y. Mak, Q.F. Gan, C.S. Ramesha, M. A. Mulkins, J. Dunn, O.Y. So, T. Biegel, M. Dinh, P. Volkel, J. Barnett, S.A. Dalrymple, S. Lee, M. Huber, A small molecule ubiquitination inhibitor blocks NF-kappa B-dependent cytokine expression in cells and rats, *J. Biol. Chem.* 277 (2002) 23573–23581.
- [97] A. Yaron, H. Gonen, I. Alkalay, A. Hatzubai, S. Jung, S. Beyth, F. Mercurio, A.M. Manning, A. Ciechanover, Y. Ben-Neriah, Inhibition of NF-kappa-B cellular function via specific targeting of the I-kappa-B-ubiquitin ligase, *EMBO J.* 16 (1997) 6486–6494.
- [98] W. Tang, Y. Li, D. Yu, A. Thomas-Tikhonenko, V.S. Spiegelman, S.Y. Fuchs, Targeting beta-transducin repeat-containing protein E3 ubiquitin ligase augments the effects of antitumor drugs on breast cancer cells, *Cancer Res.* 65 (2005) 1904–1908.
- [99] N. Shembade, A. Ma, E.W. Harhaj, Inhibition of NF-kappaB signaling by A20 through disruption of ubiquitin enzyme complexes, *Science* 327 (2010) 1135–1139.
- [100] Y.Z. Lin, S.Y. Yao, R.A. Veatch, T.R. Torgerson, J. Hawiger, Inhibition of nuclear translocation of transcription factor NF-kappa B by a synthetic peptide containing a cell membrane-permeable motif and nuclear localization sequence, *J. Biol. Chem.* 270 (1995) 14255–14258.
- [101] T.R. Torgerson, A.D. Colosia, J.P. Donahue, Y.Z. Lin, J. Hawiger, Regulation of NF-kappa B, AP-1, NFAT, and STAT1 nuclear import in T lymphocytes by noninvasive delivery of peptide carrying the nuclear localization sequence of NF-kappa B p50, *J. Immunol.* 161 (1998) 6084–6092.
- [102] T. Letoha, C. Somlai, T. Takacs, A. Szabolcs, K. Jarmay, Z. Rakonczay Jr., P. Hegyi, I. Varga, J. Kaszaki, I. Krizbai, I. Boros, E. Duda, E. Kusz, B. Penke, A nuclear import inhibitory peptide ameliorates the severity of cholecystokinin-induced acute pancreatitis, *World J. Gastroenterol.* 11 (2005) 990–999.
- [103] A. Abate, S. Oberle, H. Schroder, Lipopolysaccharide-induced expression of cyclooxygenase-2 in mouse macrophages is inhibited by chloromethylketones and a direct inhibitor of NF-kappa B translocation, *Prostaglandins Other Lipid Mediat* 56 (1998) 277–290.
- [104] V. Kolenko, T. Bloom, P. Rayman, R. Bukowski, E. Hsi, J. Finke, Inhibition of NF-kappa B activity in human T lymphocytes induces caspase-dependent apoptosis without detectable activation of caspase-1 and -3, *J. Immunol.* 163 (1999) 590–598.
- [105] R.R. Mohan, R.R. Mohan, W.J. Kim, S.E. Wilson, Modulation of TNF-alpha-induced apoptosis in corneal fibroblasts by transcription factor NF-kappaB, *Invest. Ophthalmol. Vis. Sci.* 41 (2000) 1327–1336.
- [106] E. Koulich, T. Nguyen, K. Johnson, C. Giardina, S. D'Mello, NF-kappaB is involved in the survival of cerebellar granule neurons: association of IkappaBbeta [correction of Ikappabeta] phosphorylation with cell survival, *J. Neurochem.* 76 (2001) 1188–1198.
- [107] F. D'Acquisto, A. Ialenti, A. Iano, R. Di Vaio, R. Carnuccio, Local administration of transcription factor decoy oligonucleotides to nuclear factor-kappaB prevents carrageenin-induced inflammation in rat hind paw, *Gene Ther.* 7 (2000) 1731–1737.
- [108] F. D'Acquisto, A. Iano, A. Ialenti, P. Maffia, M.C. Maiuri, R. Carnuccio, Transcription factor decoy oligodeoxynucleotides to nuclear factor-kappaB inhibit reverse passive Arthus reaction in rat, *Naunyn-Schmiedeberg's Arch. Pharmacol.* 364 (2001) 422–429.
- [109] K. Umezawa, Inhibition of tumor growth by NF-kappaB inhibitors, *Cancer Sci.* 97 (2006) 990–995.
- [110] R. Kiernan, V. Bres, R.W. Ng, M.P. Coudart, S. El Messaoudi, C. Sardet, D.Y. Jin, S. Emiliani, M. Benkirane, Post-activation turn-off of NF-kappa B-dependent transcription is regulated by acetylation of p65, *J. Biol. Chem.* 278 (2003) 2758–2766.
- [111] L.F. Chen, W.C. Greene, Shaping the nuclear action of NF-kappaB, *Nat. Rev. Mol. Cell Biol.* 5 (2004) 392–401.
- [112] F. Yeung, J.E. Hoberg, C.S. Ramsey, M.D. Keller, D.R. Jones, R.A. Frye, M.W. Mayo, Modulation of NF-kappaB-dependent transcription and cell survival by the SIRT1 deacetylase, *EMBO J.* 23 (2004) 2369–2380.
- [113] W.C. Greene, L.F. Chen, Regulation of NF-kappaB action by reversible acetylation, *Novartis Found. Symp.* 259 (2004) 208–217 discussion 218–225.
- [114] K.C. Choi, Y.H. Lee, M.G. Jung, S.H. Kwon, M.J. Kim, W.J. Jun, J.M. Lee, H.G. Yoon, Gallic acid suppresses lipopolysaccharide-induced nuclear factor-kappaB signaling by preventing RelA acetylation in A549 lung cancer cells, *Mol. Cancer Res.* 7 (2009) 2011–2021.



- [115] J. Park, J.H. Lee, M. La, M.J. Jang, G.W. Chae, S.B. Kim, H. Tak, Y. Jung, B. Byun, J.K. Ahn, C.O. Joe, Inhibition of NF-kappaB acetylation and its transcriptional activity by Daxx, *J. Mol. Biol.* 368 (2007) 388–397.
- [116] B. Sung, M.K. Pandey, K.S. Ahn, T. Yi, M.M. Chaturvedi, M. Liu, B.B. Aggarwal, Anacardic acid (6-nonadecyl salicylic acid), an inhibitor of histone acetyltransferase, suppresses expression of nuclear factor-kappaB-regulated gene products involved in cell survival, proliferation, invasion, and inflammation through inhibition of the inhibitory subunit of nuclear factor-kappaB kinase, leading to potentiation of apoptosis, *Blood* 111 (2008) 4880–4891.
- [117] X.D. Yang, E. Tajkhorshid, L.F. Chen, Functional Interplay between Acetylation and Methylation of the RelA Subunit of NF-(kappa)B, *Mol. Cell. Biol.* 30 (2010) 2170–2180.
- [118] S. Zhang, Y.K. Won, C.N. Ong, H.M. Shen, Anti-cancer potential of sesquiterpene lactones: bioactivity and molecular mechanisms, *Curr. Med. Chem. Anticancer Agents* 5 (2005) 239–249.
- [119] A.J. Garcia-Pineres, V. Castro, G. Mora, T.J. Schmidt, E. Strunck, H.L. Pahl, I. Merfort, Cysteine 38 in p65/NF-kappaB plays a crucial role in DNA binding inhibition by sesquiterpene lactones, *J. Biol. Chem.* 276 (2001) 39713–39720.
- [120] A.J. Garcia-Pineres, M.T. Lindenmeyer, I. Merfort, Role of cysteine residues of p65/NF-kappaB on the inhibition by the sesquiterpene lactone parthenolide and N-ethyl maleimide, and on its transactivating potential, *Life Sci.* 75 (2004) 841–856.
- [121] S. Wagner, A. Hofmann, B. Siedle, L. Terfloth, I. Merfort, J. Gasteiger, Development of a structural model for NF-kappaB inhibition of sesquiterpene lactones using self-organizing neural networks, *J. Med. Chem.* 49 (2006) 2241–2252.
- [122] R. Morishita, T. Sugimoto, M. Aoki, I. Kida, N. Tomita, A. Moriguchi, K. Maeda, Y. Sawa, Y. Kaneda, J. Higaki, T. Ogihara, In vivo transfection of cis element “decoy” against nuclear factor-kappaB binding site prevents myocardial infarction, *Nat. Med.* 3 (1997) 894–899.
- [123] A.R. Khaled, E.J. Butfiloski, E.S. Sobel, J. Schiftenbauer, Use of phosphorothioate-modified oligodeoxynucleotides to inhibit NF-kappaB expression and lymphocyte function, *Clin. Immunol. Immunopathol.* 86 (1998) 170–179.
- [124] C. Kupatt, R. Wichels, M. Deiss, A. Molnar, C. Leberer, P. Raake, G. von Degenfeld, D. Hahnel, P. Boekstegers, Retroinfusion of NFkappaB decoy oligonucleotide extends cardioprotection achieved by CD18 inhibition in a preclinical study of myocardial ischemia and retroinfusion in pigs, *Gene Ther.* 9 (2002) 518–526.
- [125] N. Tomita, T. Ogihara, R. Morishita, Transcription factors as molecular targets: molecular mechanisms of decoy ODN and their design, *Curr. Drug Targets* 4 (2003) 603–608.
- [126] R. Crinelli, M. Bianchi, L. Gentilini, L. Palma, M. Magnani, Locked nucleic acids (LNA): versatile tools for designing oligonucleotide decoys with high stability and affinity, *Curr. Drug Targets* 5 (2004) 745–752.
- [127] I. Isomura, A. Morita, Regulation of NF-kappaB signaling by decoy oligodeoxynucleotides, *Microbiol. Immunol.* 50 (2006) 559–563.
- [128] I. Kawamura, R. Morishita, N. Tomita, E. Lacey, M. Aketa, S. Tsujimoto, T. Manda, M. Tomoi, I. Kida, J. Higaki, Y. Kaneda, K. Shimomura, T. Ogihara, Intratumoral injection of oligonucleotides to the NF kappa B binding site inhibits cachexia in a mouse tumor model, *Gene Ther.* 6 (1999) 91–97.
- [129] D.J. Van Antwerp, I.M. Verma, Signal-induced degradation of I(kappa)B(alpha): association with NF-kappaB and the PEST sequence in I(kappa)B(alpha) are not required, *Mol. Cell. Biol.* 16 (1996) 6037–6045.
- [130] C. Jobin, A. Panja, C. Hellerbrand, Y. Iimuro, J. Didonato, D.A. Brenner, R.B. Sartor, Inhibition of proinflammatory molecule production by adenovirus-mediated expression of a nuclear factor kappaB super-repressor in human intestinal epithelial cells, *J. Immunol.* 160 (1998) 410–418.
- [131] M. Bentires-Alj, A.C. Hellin, M. Ameyar, S. Chouaib, M.P. Merville, V. Bours, Stable inhibition of nuclear factor kappaB in cancer cells does not increase sensitivity to cytotoxic drugs, *Cancer Res.* 59 (1999) 811–815.
- [132] Y. Abu-Amer, S.F. Dowdy, F.P. Ross, J.C. Clohisy, S.L. Teitelbaum, TAT fusion proteins containing tyrosine 42-deleted IkappaBalpha arrest osteoclastogenesis, *J. Biol. Chem.* 276 (2001) 30499–30503.
- [133] D.J. Van Antwerp, S.J. Martin, T. Kafri, D.R. Green, I.M. Verma, Suppression of TNF-alpha-induced apoptosis by NF-kappaB, *Science* 274 (1996) 787–789.
- [134] C.Y. Wang, M.W. Mayo, A.S. Baldwin Jr., TNF- and cancer therapy-induced apoptosis: potentiation by inhibition of NF-kappaB, *Science* 274 (1996) 784–787.
- [135] P.B. Bushdid, D.M. Brantley, F.E. Yull, G.L. Blaeuer, L.H. Hoffman, L. Niswander, L.D. Kerr, Inhibition of NF-kappaB activity results in disruption of the apical ectodermal ridge and aberrant limb morphogenesis, *Nature* 392 (1998) 615–618.
- [136] Y. Kanegae, A.T. Tavares, J.C. Izpisua Belmonte, I.M. Verma, Role of Rel/NF-kappaB transcription factors during the outgrowth of the vertebrate limb, *Nature* 392 (1998) 611–614.
- [137] C.Y. Wang, J.C. Cusack Jr., R. Liu, A.S. Baldwin Jr., Control of inducible chemoresistance: enhanced anti-tumor therapy through increased apoptosis by inhibition of NF-kappaB, *Nat. Med.* 5 (1999) 412–417.
- [138] D.C. Duffey, Z. Chen, G. Dong, F.G. Ondrey, J.S. Wolf, K. Brown, U. Siebenlist, C. Van Waes, Expression of a dominant-negative mutant inhibitor-kappaBalpha of nuclear factor-kappaB in human head and neck squamous cell carcinoma inhibits survival, proinflammatory cytokine expression, and tumor growth in vivo, *Cancer Res.* 59 (1999) 3468–3474.
- [139] N.S. Chang, The non-ankyrin C terminus of Ikappa Balpha physically interacts with p53 in vivo and dissociates in response to apoptotic stress, hypoxia, DNA damage, and transforming growth factor-beta 1-mediated growth suppression, *J. Biol. Chem.* 277 (2002) 10323–10331.
- [140] J. Li, S.H. Joo, M.D. Tsai, An NF-kappaB-specific inhibitor, IkappaBalpha, binds to and inhibits cyclin-dependent kinase 4, *Biochemistry* 42 (2003) 13476–13483.
- [141] C. Aguilera, R. Hoya-Arias, G. Haegeman, L. Espinosa, A. Bigas, Recruitment of IkappaBalpha to the hes1 promoter is associated with transcriptional repression, *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 16537–16542.
- [142] M. van Hogerlinden, B.L. Rozell, L. Ahrlund-Richter, R. Toftgard, Squamous cell carcinomas and increased apoptosis in skin with inhibited Rel/nuclear factor-kappaB signaling, *Cancer Res.* 59 (1999) 3299–3303.
- [143] F.J. Staal, M. Roederer, L.A. Herzenberg, L.A. Herzenberg, Intracellular thiols regulate activation of nuclear factor kappa B and transcription of human immunodeficiency virus, *Proc. Natl. Acad. Sci. U. S. A.* 87 (1990) 9943–9947.
- [144] S. Mihm, J. Ennen, U. Pessara, R. Kurth, W. Droge, Inhibition of HIV-1 replication and NF-kappa B activity by cysteine and cysteine derivatives, *AIDS* 5 (1991) 497–503.
- [145] C. Bubic, S. Papa, K. Dean, G. Franzoso, Mutual cross-talk between reactive oxygen species and nuclear factor-kappa B: molecular basis and biological significance, *Oncogene* 25 (2006) 6731–6748.
- [146] G. Gloire, S. Legrand-Poels, J. Piette, NF-kappaB activation by reactive oxygen species: fifteen years later, *Biochem. Pharmacol.* 72 (2006) 1493–1505.
- [147] M. Hayakawa, H. Miyashita, I. Sakamoto, M. Kitagawa, H. Tanaka, H. Yasuda, M. Karin, K. Kikugawa, Evidence that reactive oxygen species do not mediate NF-kappaB activation, *EMBO J.* 22 (2003) 3356–3366.
- [148] K. Schulze-Osthoff, R. Beyaert, V. Vandevorde, G. Haegeman, W. Fiers, Depletion of the mitochondrial electron transport abrogates the cytotoxic and gene-inductive effects of TNF, *EMBO J.* 12 (1993) 3095–3104.
- [149] S.K. Manna, H.J. Zhang, T. Yan, L.W. Oberley, B.B. Aggarwal, Overexpression of manganese superoxide dismutase suppresses tumor necrosis factor-induced apoptosis and activation of nuclear transcription factor-kappaB and activated protein-1, *J. Biol. Chem.* 273 (1998) 13245–13254.
- [150] S.K. Manna, M.T. Kuo, B.B. Aggarwal, Overexpression of gamma-glutamylcysteine synthetase suppresses tumor necrosis factor-induced apoptosis and activation of nuclear transcription factor-kappa B and activator protein-1, *Oncogene* 18 (1999) 4371–4382.
- [151] K. Natarajan, S. Singh, T.R. Burke Jr., D. Grunberger, B.B. Aggarwal, Caffeic acid phenethyl ester is a potent and specific inhibitor of activation of nuclear transcription factor NF-kappa B, *Proc. Natl. Acad. Sci. U. S. A.* 93 (1996) 9090–9095.
- [152] J. Hiscott, T.L. Nguyen, M. Arguello, P. Nakhaei, S. Paz, Manipulation of the nuclear factor-kappaB pathway and the innate immune response by viruses, *Oncogene* 25 (2006) 6844–6867.
- [153] P.P. Powell, L.K. Dixon, R.M. Parkhouse, An IkappaB homolog encoded by African swine fever virus provides a novel mechanism for downregulation of proinflammatory cytokine responses in host macrophages, *J. Virol.* 70 (1996) 8527–8533.
- [154] C. Camus-Bouclainville, L. Fiette, S. Bouchiha, B. Pignolet, D. Counor, C. Filipe, J. Gelfi, F. Messud-Petit, A virulence factor of myxoma virus colocalizes with NF-kappaB in the nucleus and interferes with inflammation, *J. Virol.* 78 (2004) 2510–2516.
- [155] H. Thoeftiktitkul, M.H. Beck, M.R. Strand, Inhibitor kappaB-like proteins from a polydnavirus inhibit NF-kappaB activation and suppress the insect immune response, *Proc. Natl. Acad. Sci. U. S. A.* 102 (2005) 11426–11431.
- [156] Y. Revilla, M. Callejo, J.M. Rodriguez, E. Culebras, M.L. Nogal, M.L. Salas, E. Vinuela, M. Fresno, Inhibition of nuclear factor kappaB activation by a virus-encoded IkappaB-like protein, *J. Biol. Chem.* 273 (1998) 5405–5411.
- [157] N. Neznanov, K.M. Chumakov, L. Neznanova, A. Almasan, A.K. Banerjee, A.V. Gudkov, Proteolytic cleavage of the p65-RelA subunit of NF-kappaB during poliovirus infection, *J. Biol. Chem.* 280 (2005) 24153–24158.
- [158] D.B. Nichols, J.L. Shisler, The MC160 protein expressed by the dermatotropic poxvirus molluscum contagiosum virus prevents tumor necrosis factor alpha-induced NF-kappaB activation via inhibition of I kappa kinase complex formation, *J. Virol.* 80 (2006) 578–586.
- [159] S.H. Choi, K.J. Park, B.Y. Ahn, G. Jung, M.M. Lai, S.B. Hwang, Hepatitis C virus nonstructural 5B protein regulates tumor necrosis factor alpha signaling through effects on cellular IkappaB kinase, *Mol. Cell. Biol.* 26 (2006) 3048–3059.
- [160] K. Schesser, A.K. Spiik, J.M. Dukuzumuremyi, M.F. Neurath, S. Pettersson, H. Wolf-Watz, The yopJ locus is required for Yersinia-mediated inhibition of NF-kappaB activation and cytokine expression: YopJ contains a eukaryotic SH2-like domain that is essential for its repressive activity, *Mol. Microbiol.* 28 (1998) 1067–1079.
- [161] K. Orth, L.E. Palmer, Z.Q. Bao, S. Stewart, A.E. Rudolph, J.B. Bliska, J.E. Dixon, Inhibition of the mitogen-activated protein kinase kinase superfamily by a Yersinia effector, *Science* 285 (1999) 1920–1923.
- [162] L.S. Collier-Hyams, H. Zeng, J. Sun, A.D. Tomlinson, Z.Q. Bao, H. Chen, J.L. Madara, K. Orth, A.S. Neish, Cutting edge: Salmonella AvrA effector inhibits the key proinflammatory, anti-apoptotic NF-kappa B pathway, *J. Immunol.* 169 (2002) 2846–2850.
- [163] H.L. Pahl, B. Krauss, K. Schulze-Osthoff, T. Decker, E.B. Traenckner, M. Vogt, C. Myers, T. Parks, W. Warring, A. Muhlbacher, A.P. Czernilofsky, P.A. Baueerle, The immunosuppressive fungal metabolite gliotoxin specifically inhibits transcription factor NF-kappaB, *J. Exp. Med.* 183 (1996) 1829–1840.
- [164] G. Erkel, T. Anke, O. Sterner, Inhibition of NF-kappa B activation by panepoxydione, *Biochem. Biophys. Res. Commun.* 226 (1996) 214–221.
- [165] K. Umezawa, A. Ariga, N. Matsumoto, Naturally occurring and synthetic inhibitors of NF-kappaB functions, *Anticancer Drug Des.* 15 (2000) 239–244.
- [166] K. De Bosscher, W. Vanden Berghe, G. Haegeman, Cross-talk between nuclear receptors and nuclear factor kappaB, *Oncogene* 25 (2006) 6868–6886.
- [167] D. Kalaitzidis, T.D. Gilmore, Transcription factor cross-talk: the estrogen receptor and NF-kappaB, *Trends Endocrinol. Metab.* 16 (2005) 46–52.
- [168] S. Olivier, P. Close, E. Castermans, L. de Leval, S. Tabruyn, A. Chariot, M. Malaise, M.P. Merville, V. Bours, N. Franchimont, Raloxifene-induced myeloma cell

- apoptosis: a study of nuclear factor-kappaB inhibition and gene expression signature, *Mol. Pharmacol.* 69 (2006) 1615–1623.
- [169] T.P. McDade, R.A. Perugini, F.J. Vitimberga Jr., R.C. Carrigan, M.P. Callery, Salicylates inhibit NF-kappaB activation and enhance TNF-alpha-induced apoptosis in human pancreatic cancer cells, *J. Surg. Res.* 83 (1999) 56–61.
  - [170] Y. Yamamoto, M.J. Yin, K.M. Lin, R.B. Gaynor, Sulindac inhibits activation of the NF-kappaB pathway, *J. Biol. Chem.* 274 (1999) 27307–27314.
  - [171] M.J. Yin, Y. Yamamoto, R.B. Gaynor, The anti-inflammatory agents aspirin and salicylate inhibit the activity of I(kappa)B kinase-beta, *Nature* 396 (1998) 77–80.
  - [172] P. Dikshit, M. Chatterjee, A. Goswami, A. Mishra, N.R. Jana, Aspirin induces apoptosis through the inhibition of proteasome function, *J. Biol. Chem.* 281 (2006) 29228–29235.
  - [173] M.F. McCarty, K.I. Block, Preadministration of high-dose salicylates, suppressors of NF-kappaB activation, may increase the chemosensitivity of many cancers: an example of proapoptotic signal modulation therapy, *Integr. Cancer Ther.* 5 (2006) 252–268.
  - [174] M. Yuan, N. Konstantopoulos, J. Lee, L. Hansen, Z.W. Li, M. Karin, S.E. Shoelson, Reversal of obesity- and diet-induced insulin resistance with salicylates or targeted disruption of Ikkbeta, *Science* 293 (2001) 1673–1677.
  - [175] J.J. Li, C.H. Fang, C-reactive protein is not only an inflammatory marker but also a direct cause of cardiovascular diseases, *Med. Hypotheses* 62 (2004) 499–506.
  - [176] E. Kopp, S. Ghosh, Inhibition of NF-kappa B by sodium salicylate and aspirin, *Science* 265 (1994) 956–959.
  - [177] M. Grilli, M. Pizzi, M. Memo, P. Spano, Neuroprotection by aspirin and sodium salicylate through blockade of NF-kappaB activation, *Science* 274 (1996) 1383–1385.
  - [178] S.T. Palayoor, E.A. Bump, S.K. Calderwood, S. Bartol, C.N. Coleman, Combined antitumor effect of radiation and ibuprofen in human prostate carcinoma cells, *Clin. Cancer Res.* 4 (1998) 763–771.
  - [179] Y. Takada, A. Bhardwaj, P. Potdar, B.B. Aggarwal, Nonsteroidal anti-inflammatory agents differ in their ability to suppress NF-kappaB activation, inhibition of expression of cyclooxygenase-2 and cyclin D1, and abrogation of tumor cell proliferation, *Oncogene* 23 (2004) 9247–9258.
  - [180] S. Meyer, N.G. Kohler, A. Joly, Cyclosporine A is an uncompetitive inhibitor of proteasome activity and prevents NF-kappaB activation, *FEBS Lett.* 413 (1997) 354–358.
  - [181] B. Frantz, E.C. Nordby, G. Bren, N. Steffan, C.V. Paya, R.L. Kincaid, M.J. Tocci, S.J. O'Keefe, E.A. O'Neill, Calcineurin acts in synergy with PMA to inactivate I kappa B/MAD3, an inhibitor of NF-kappa B, *EMBO J.* 13 (1994) 861–870.
  - [182] M.A. Tepper, S.G. Nadler, J.M. Esselstyn, K.G. Sterbenz, Deoxyspergualin inhibits kappa light chain expression in 70Z/3 pre-B cells by blocking lipopolysaccharide-induced NF-kappa B activation, *J. Immunol.* 155 (1995) 2427–2436.
  - [183] P.G. McCaffrey, P.K. Kim, V.E. Valge-Archer, R. Sen, A. Rao, Cyclosporin A sensitivity of the NF-kappa B site of the IL2R alpha promoter in untransformed murine T cells, *Nucleic Acids Res.* 22 (1994) 2134–2142.
  - [184] A.S. Wechsler, M.C. Gordon, U. Dendorfer, K.P. LeClair, Induction of IL-8 expression in T cells uses the CD28 costimulatory pathway, *J. Immunol.* 153 (1994) 2515–2523.
  - [185] D. Kunz, G. Walker, W. Eberhardt, D. Nitsch, J. Pfeilschifter, Interleukin 1 beta-induced expression of nitric oxide synthase in rat renal mesangial cells is suppressed by cyclosporin A, *Biochem. Biophys. Res. Commun.* 216 (1995) 438–446.
  - [186] D. Qiu, G. Zhao, Y. Aoki, L. Shi, A. Uyei, S. Nazarian, J.C. Ng, P.N. Kao, Immunosuppressant PG490 (triptolide) inhibits T-cell interleukin-2 expression at the level of purine-box/nuclear factor of activated T-cells and NF-kappaB transcriptional activation, *J. Biol. Chem.* 274 (1999) 13443–13450.
  - [187] J. Sen, L. Venkataraman, Y. Shinkai, J.W. Pierce, F.W. Alt, S.J. Burakoff, R. Sen, Expression and induction of nuclear factor-kappa B-related proteins in thymocytes, *J. Immunol.* 154 (1995) 3213–3221.
  - [188] L. Venkataraman, S.J. Burakoff, R. Sen, FK506 inhibits antigen receptor-mediated induction of c-rel in B and T lymphoid cells, *J. Exp. Med.* 181 (1995) 1091–1099.
  - [189] G.A. Webster, N.D. Perkins, Transcriptional cross talk between NF-kappaB and p53, *Mol. Cell. Biol.* 19 (1999) 3485–3495.
  - [190] W.C. Huang, T.K. Ju, M.C. Hung, C.C. Chen, Phosphorylation of CBP by IKKalpha promotes cell growth by switching the binding preference of CBP from p53 to NF-kappaB, *Mol. Cell* 26 (2007) 75–87.
  - [191] K.V. Gurova, J.E. Hill, C. Guo, A. Prokvolit, L.G. Burdelya, E. Samoylova, A.V. Khodyakova, R. Ganapathi, M. Ganapathi, N.D. Tararova, D. Bosykh, D. Lvovskiy, T.R. Webb, G.R. Stark, A.V. Gudkov, Small molecules that reactivate p53 in renal cell carcinoma reveal a NF-kappaB-dependent mechanism of p53 suppression in tumors, *Proc. Natl. Acad. Sci. U. S. A.* 102 (2005) 17448–17453.
  - [192] C. Tovar, J. Rosinski, Z. Filipovic, B. Higgins, K. Kolinsky, H. Hilton, X. Zhao, B.T. Vu, W. Qing, K. Packman, O. Myklebost, D.C. Heimbrook, L.T. Vassilev, Small-molecule MDM2 antagonists reveal aberrant p53 signaling in cancer: implications for therapy, *Proc. Natl. Acad. Sci. U.S.A.* 103 (2006) 1888–1893.
  - [193] A. Dey, E.T. Wong, P. Bist, V. Tergaonkar, D.P. Lane, Nutlin-3 inhibits the NF-kappaB pathway in a p53-dependent manner: implications in lung cancer therapy, *Cell Cycle* 6 (2007) 2178–2185.
  - [194] K.J. Campbell, J.M. Witty, S. Rocha, N.D. Perkins, Cisplatin mimics ARF tumor suppressor regulation of RelA (p65) nuclear factor-kappaB transactivation, *Cancer Res.* 66 (2006) 929–935.
  - [195] A. Dey, V. Tergaonkar, D.P. Lane, Double-edged swords as cancer therapeutics: simultaneously targeting p53 and NF-kappaB pathways, *Nat. Rev. Drug Discov.* 7 (2008) 1031–1040.
  - [196] M.S. Rodriguez, J. Thompson, R.T. Hay, C. Dargemont, Nuclear retention of IkappaBalpha protects it from signal-induced degradation and inhibits nuclear factor kappaB transcriptional activation, *J. Biol. Chem.* 274 (1999) 9108–9115.
  - [197] T.R. Kau, P.A. Silver, Nuclear transport as a target for cell growth, *Drug Discov. Today* 8 (2003) 78–85.
  - [198] A. Dasgupta, K.J. Jung, S.J. Jeong, J.N. Brady, Inhibition of methyltransferases results in induction of g2/m checkpoint and programmed cell death in human T-lymphotropic virus type 1-transformed cells, *J. Virol.* 82 (2008) 49–59.
  - [199] H.L. Li, D.D. Chen, X.H. Li, H.W. Zhang, Y.Q. Lu, C.L. Ye, X.D. Ren, Changes of NF-kB, p53, Bcl-2 and caspase in apoptosis induced by JTE-522 in human gastric adenocarcinoma cell line AGS cells: role of reactive oxygen species, *World J. Gastroenterol.* 8 (2002) 431–435.
  - [200] W. Lu, L. Chen, Y. Peng, J. Chen, Activation of p53 by roscovitine-mediated suppression of MDM2 expression, *Oncogene* 20 (2001) 3206–3216.
  - [201] A. Dey, E.T. Wong, C.F. Cheok, V. Tergaonkar, D.P. Lane, R-Roscovitine simultaneously targets both the p53 and NF-kappaB pathways and causes potentiation of apoptosis: implications in cancer therapy, *Cell Death Differ.* 15 (2008) 263–273.
  - [202] Z.N. Demidenko, M.V. Blagosklonny, Flavopiridol induces p53 via initial inhibition of Mdm2 and p21 and, independently of p53, sensitizes apoptosis-reluctant cells to tumor necrosis factor, *Cancer Res.* 64 (2004) 3653–3660.
  - [203] Y. Takada, B.B. Aggarwal, Flavopiridol inhibits NF-kappaB activation induced by various carcinogens and inflammatory agents through inhibition of IkappaBalpha kinase and p65 phosphorylation: abrogation of cyclin D1, cyclooxygenase-2, and matrix metalloproteinase-9, *J. Biol. Chem.* 279 (2004) 4750–4759.
  - [204] D. Bernard, D. Monte, B. Vandenbunder, C. Abbadié, The c-Rel transcription factor can both induce and inhibit apoptosis in the same cells via the upregulation of MnSOD, *Oncogene* 21 (2002) 4392–4402.
  - [205] A.M. Sheehy, M.S. Schlissel, Overexpression of RelA causes G1 arrest and apoptosis in a pro-B cell line, *J. Biol. Chem.* 274 (1999) 8708–8716.
  - [206] B. Kaltschmidt, C. Kaltschmidt, T.G. Hofmann, S.P. Hehner, W. Dröge, M.L. Schmitz, The pro- or anti-apoptotic function of NF-kappaB is determined by the nature of the apoptotic stimulus, *Eur. J. Biochem.* 267 (2000) 3828–3835.
  - [207] J. Bohuslav, L.F. Chen, H. Kwon, Y. Mu, W.C. Greene, p53 induces NF-kappaB activation by an IkappaB kinase-independent mechanism involving phosphorylation of p65 by ribosomal S6 kinase 1, *J. Biol. Chem.* 279 (2004) 26115–26125.
  - [208] K.M. Ryan, M.K. Ernst, N.R. Rice, K.H. Vousden, Role of NF-kappaB in p53-mediated programmed cell death, *Nature* 404 (2000) 892–897.
  - [209] M. Karin, M. Delhase, The I kappa B kinase (IKK) and NF-kappa B: key elements of proinflammatory signalling, *Semin. Immunol.* 12 (2000) 85–98.
  - [210] M.P. Belich, A. Salmeron, L.H. Johnston, S.C. Ley, TPL-2 kinase regulates the proteolysis of the NF-kappaB-inhibitory protein NF-kappaB1 p105, *Nature* 397 (1999) 363–368.
  - [211] P. Viatour, M.P. Merville, V. Bours, A. Chariot, Phosphorylation of NF-kappaB and IkappaB proteins: implications in cancer and inflammation, *Trends Biochem. Sci.* 30 (2005) 43–52.
  - [212] N. Li, M. Karin, Ionizing radiation and short wavelength UV activate NF-kappaB through two distinct mechanisms, *Proc. Natl. Acad. Sci. U. S. A.* 95 (1998) 13012–13017.
  - [213] T. Kato Jr., M. Delhase, A. Hoffmann, M. Karin, CK2 is a C-terminal IkappaB kinase responsible for NF-kappaB activation during the UV response, *Mol. Cell* 12 (2003) 829–839.
  - [214] G. Sethi, K.S. Ahn, M.M. Chaturvedi, B.B. Aggarwal, Epidermal growth factor (EGF) activates nuclear factor-kappaB through IkappaBalpha kinase-independent but EGF receptor-kinase dependent tyrosine 42 phosphorylation of IkappaBalpha, *Oncogene* 26 (2007) 7324–7332.
  - [215] A.P. West, A.A. Koblansky, S. Ghosh, Recognition and signaling by toll-like receptors, *Annu. Rev. Cell Dev. Biol.* 22 (2006) 409–437.
  - [216] Y. Takada, A. Mukhopadhyay, G.C. Kundu, G.H. Mahabeshwar, S. Singh, B.B. Aggarwal, Hydrogen peroxide activates NF-kappa B through tyrosine phosphorylation of I kappa B alpha and serine phosphorylation of p65: evidence for the involvement of I kappa B alpha kinase and Syk protein-tyrosine kinase, *J. Biol. Chem.* 278 (2003) 24233–24241.
  - [217] V. Imbert, R.A. Rupec, A. Livolsi, H.L. Pahl, E.B. Traenckner, C. Mueller-Dieckmann, D. Farahifar, B. Rossi, P. Auberger, P.A. Baeuerle, J.F. Peyron, Tyrosine phosphorylation of I kappa B-alpha activates NF-kappa B without proteolytic degradation of I kappa B-alpha, *Cell* 86 (1996) 787–798.
  - [218] J. Schulze-Luehmann, S. Ghosh, Antigen-receptor signaling to nuclear factor kappa B, *Immunology* 25 (2006) 701–715.
  - [219] H. Noels, G. van Loo, S. Hagens, V. Broeckx, R. Beyaert, P. Marynen, M. Baens, A Novel TRAF6 binding site in MALT1 defines distinct mechanisms of NF-kappaB activation by API2middle dotMALT1 fusions, *J. Biol. Chem.* 282 (2007) 10180–10189.
  - [220] R.R. Rosato, S.S. Kolla, S.K. Hock, J.A. Almenara, A. Patel, S. Amin, P. Adajia, P. Dent, S. Grant, HDAC inhibitors activate NF-(kappa)B in human leukemia cells through an ATM/nemo-related pathway, *J. Biol. Chem.* 285 (2010) 10064–10077.
  - [221] Z.H. Wu, Y. Shi, R.S. Tibbetts, S. Miyamoto, Molecular linkage between the kinase ATM and NF-kappaB signaling in response to genotoxic stimuli, *Science* 311 (2006) 1141–1146.
  - [222] S. Janssens, A. Tinel, S. Lippens, J. Tschopp, PIDD mediates NF-kappaB activation in response to DNA damage, *Cell* 123 (2005) 1079–1092.
  - [223] G.M. Hur, J. Lewis, Q. Yang, Y. Lin, H. Nakano, S. Nedospasov, Z.G. Liu, The death domain kinase RIP has an essential role in DNA damage-induced NF-kappa B activation, *Genes Dev.* 17 (2003) 873–882.