

NF- κ B Signaling Pathways: Role in Nervous System Physiology and Pathology

Stefka Mincheva-Tasheva and Rosa M. Soler

Neuroscientist 2013 19: 175 originally published online 10 July 2012

DOI: 10.1177/1073858412444007

The online version of this article can be found at:

<http://nro.sagepub.com/content/19/2/175>

Published by:



<http://www.sagepublications.com>

Additional services and information for *The Neuroscientist* can be found at:

Email Alerts: <http://nro.sagepub.com/cgi/alerts>

Subscriptions: <http://nro.sagepub.com/subscriptions>

Reprints: <http://www.sagepub.com/journalsReprints.nav>


Permissions: <http://www.sagepub.com/journalsPermissions.nav>

>> [Version of Record](#) - Mar 13, 2013

[OnlineFirst Version of Record](#) - Jul 10, 2012

[What is This?](#)

NF- κ B Signaling Pathways: Role in Nervous System Physiology and Pathology

The Neuroscientist
19(2) 175–194
© The Author(s) 2012
Reprints and permission: <http://www.sagepub.com/journalsPermissions.nav>
DOI: 10.1177/1073858412444007
<http://nro.sagepub.com>


Stefka Mincheva-Tasheva¹ and Rosa M. Soler¹

Abstract

Intracellular pathways related to cell survival regulate neuronal physiology during development and neurodegenerative disorders. One of the pathways that have recently emerged with an important role in these processes is nuclear factor- κ B (NF- κ B). The activity of this pathway leads to the nuclear translocation of the NF- κ B transcription factors and the regulation of anti-apoptotic gene expression. Different stimuli can activate the pathway through different intracellular cascades (canonical, non-canonical, and atypical), contributing to the translocation of specific dimers of the NF- κ B transcription factors, and each of these dimers can regulate the transcription of different genes. Recent studies have shown that the activation of this pathway regulates opposite responses such as cell survival or neuronal degeneration. These apparent contradictory effects depend on conditions such as the pathway stimuli, the origin of the cells, or the cellular context. In the present review, the authors summarize these findings and discuss their significance with respect to survival or death in the nervous system.

Keywords

NF- κ B, neuronal survival, neurotrophic factors, RelA/p65, neurodegenerative disorders, motoneuron

Introduction

Neurons and non-neuronal cells of the nervous system require for their survival and function extracellular and intracellular signaling molecules such as neurotrophic factors or calcium. These molecules activate intracellular signaling pathways, which in turn regulate the activation or inhibition of gene expression. Intracellular pathways are the mechanisms that regulate cellular events during nervous system development, adulthood, and disease (Airaksinen and Saarma 2002; Chao 2003). New evidences that emerged during the past 10 years have shown the complex regulation of these pathways and expanded the list of those regulating neuronal survival and function. One of these pathways is the nuclear factor- κ B (NF- κ B). The NF- κ B pathway was discovered in 1986 as a transcription modulator of the light chain of B lymphocyte immunoglobulins (Sen and Baltimore 1986). Subsequent studies showed that NF- κ B is a ubiquitously expressed dimeric transcription factor involved in cellular processes such as inflammation, adhesion, proliferation, differentiation, apoptosis, and oncogenesis. This family of transcription factors also has an important role during nervous system development and pathology. The effects caused by NF- κ B activity in the nervous system are usually based on the control of neuronal apoptosis,

neurite outgrowth, and synaptic plasticity. Genes regulated by NF- κ B dimers are mainly responsible for these functions, and the long list of these genes creates an intricate intracellular network that contributes to the complexity of the pathway. The list of NF- κ B functions in the central and peripheral nervous system is apparently not finished, and some of these functions are in fact contradictory. The present review of the NF- κ B pathway places special emphasis on its function in the developing nervous system and in neurological disorders.

NF- κ B Family Members

NF- κ B is a dimer composed of members of the Rel family of transcription factors: RelA (p65), RelB, c-Rel, p50, and p52. All these proteins have in common a highly conserved 300-amino acid domain called Rel homology

¹Neuronal Signaling Unit, Dep. Ciències Mèdiques Bàsiques, Facultat de Medicina, Universitat de Lleida-IRBLLEIDA, Lleida, Spain

Corresponding Author:

Rosa M. Soler, Neuronal Signaling Unit, Dep. Ciències Mèdiques Bàsiques, Facultat de Medicina, Universitat de Lleida-IRBLLEIDA, Montserrat Roig, 2, 25008-Lleida, Spain
Email: rosa.soler@cmb.udl.cat




Figure 1. NF- κ B family members. RelA, RelB, and c-Rel constitute the NF- κ B subfamily I characterized by the presence of Rel homology domain (RHD), responsible for NF- κ B dimerization, I κ B interaction, and association to the DNA. This domain contains the nuclear localization sequence (NLS), responsible for NF- κ B nuclear translocation. The members of NF- κ B subfamily I also contain the transcription activation domain (TAD), responsible for their transcriptional activity. RelB has a leucine zipper motif (LZ). p100 and p105 and their mature forms p52 and p50, respectively, constitute the NF- κ B subfamily II. They also contain the RHD with NLS; however, in their c-terminal end p100 and p105 contain several ankyrin repeats and a death domain (DD). After proteasome degradation, ankyrin repeats and the DD are released and the mature products of these proteins, p52 and p50, are composed only by RHD with NLS.

domain (RHD). The RHD is responsible for NF- κ B dimerization, NF- κ B and I κ B interaction, and NF- κ B dimer association with DNA (Huxford and others 1999) and contains the nuclear localization sequence (NLS) necessary for NF- κ B translocation to the nucleus (Ghosh and others 1998).

The NF- κ B proteins can be classified into two subfamilies depending on their structure (Fig. 1). Subfamily I includes RelA, c-Rel, and RelB; the members of this subfamily contain a transcription activation domain (TAD). Subfamily II includes the proteins p52 and p50, generated from their precursors, p100 and p105, respectively; these do not contain the TAD and are unable to activate gene transcription after NF- κ B activation. The p100 and p105 precursors contain several ankyrin domains that are processed by the proteasome when the pathway is activated. Thus, these immature forms of p52 and p50 may be considered as inhibitors of the NF- κ B

signaling pathway (Ghosh and others 1998). Even though it is possible to find different homo- or heterodimers of NF- κ B members in the cells, in mammals the most abundant form is the RelA/p50 heterodimer.

The NF- κ B Inhibitors: I κ B

In the absence of stimuli NF- κ B homo- and/or heterodimers are present in the cytoplasm and form inactive complexes with their inhibitors, which are members of the I κ B (Inhibitor κ B) protein family (I κ B α , I κ B β , I κ B ϵ , I κ B γ , I κ B δ , and Bcl-3) (Fig. 2A). When the pathway is activated, NF- κ B dimers are released from the inhibitor and translocated to the nucleus where they bind to the κ B sequences of the DNA. The I κ B inhibitors have in common their three-dimensional structure and a variable number of ankyrin-repeat motifs located in their amino-terminal segment. These ankyrin motifs are responsible




Figure 2. Members of the NF- κ B inhibitors (I κ B) and I κ B kinase complex (IKK α , IKK β , and IKK γ). A) The inhibitory κ B (I κ B) family consists in seven members: I κ B α , I κ B β , I κ B γ , I κ B ϵ , Bcl-3 and I κ B ζ . They have in common the conserved ankyrin repeat motifs that are essential for the I κ B/NF- κ B interaction. I κ B α and I κ B β have in their carboxyl-terminal end a domain rich in proline, glutamic acid/aspartic acid, serine and threonine residues (PEST domain). The PEST domain may interact directly with the DNA-binding region of one of the NF- κ B subunits. Bcl-3 also contains a transactivation domain (TAD). B) IKK α and IKK β members of I κ B kinase family contain a kinase domain, a leucine zipper domain, and a helix-loop-helix domain, responsible for their kinase activity and dimerization. Phosphorylation of Ser176 and Ser180 or Ser177 and Ser181 located in the kinase domain induces IKK α and IKK β kinase activity, respectively. IKK α kinase domain contains the NLS responsible for its nuclear localization. The regulatory subunit IKK γ , also called NEMO, contains four domains—two coiled-coil, a leucine zipper, and a zinc finger—required for IKK complex formation and function.

for the I κ B/NF- κ B interaction (Baeuerle 1998). I κ B α and I κ B β have in their carboxyl-terminal end a domain rich in proline, glutamic acid/aspartic acid, serine, and threonine residues (PEST domain). The PEST domain may interact directly with the DNA-binding region of one of the NF- κ B subunits and is required for inhibition of DNA binding (Ernst and others 1995).

I κ B α is the most frequent expressed form of NF- κ B inhibitors in the nervous system. It has been well established that NF- κ B activation induces I κ B α phosphorylation at Ser32 and Ser 36 residues of the ankyrin repeats. This phosphorylation leads to the I κ B release of the complex, its polyubiquitination at Lys21 and Lys22 residues, and its degradation by the 26S proteasome (Chen and others 1995). However, the reduction of I κ B α protein level into the cytoplasm is temporary because I κ B α is one of the earliest genes transcribed after NF- κ B activation (Ito and others 1994). Thus I κ B α protein is reduced after the activation of the pathway, and 1 hour later the levels of the protein increase in response to the newly synthesized I κ B α (Place and others 2001). I κ B α contains NLS, which permits the localization of newly synthesized I κ B α into the nucleus where I κ B α binds with RelA/p50 dimers, removing them from the κ B DNA sequences. Then I κ B α NES (nuclear export sequence) promotes the translocation of this inactive complex from the nucleus to the cytoplasm (Tam and others 2000).

In some cell types, I κ B α can also be removed from NF- κ B by phosphorylation at Tyr42, located in the ankyrin motifs (Takada and others 2003). This phosphorylation induces I κ B α detachment from the complex, but

not its degradation by the proteasome. Thus, I κ B α protein level does not decrease early after Tyr42 phosphorylation (Imbert and others 1996).

The Kinases of the Inhibitors: I κ B Kinases

I κ B phosphorylation require the catalytic activation of a serine-threonine kinase complex called I κ B kinases (IKKs). The IKK complex is composed of three elements: two catalytic subunits, IKK α and IKK β , and a regulatory subunit, NEMO (also called IKK γ) (Fig. 2B). IKK α and IKK β contain a helix-loop-helix domain and a leucine-zipper domain (Häcker and Karin 2006; Woronicz and others 1997; Zandi and others 1997). The leucine-zipper domain is involved in the modulation of the kinase activity, and the helix-loop-helix domain is responsible for IKK homo- or heterodimerization (Zandi and others 1997). Some studies indicate that the IKK α /IKK β heterodimers activate the NF- κ B pathway more efficiently than the IKK homodimers (Huynh and others 2000). The IKK α and IKK β catalytic domains show 65% homology and their kinase activation is induced by the phosphorylation at two serine residues (Ser177/Ser181 of IKK β and Ser176/Ser180 of IKK α) (Kwak and others 2000). IKK α and IKK β contribute differentially to IKK complex activation. Studies using IKK β mutants demonstrated that this kinase is responsible for NF- κ B activation after tumor necrosis factor- α (TNF α), interleukin-1 (IL-1), or lipopolysaccharide (LPS) stimulation (Delhase and others 1999). IKK α and IKK β also have different cellular

distribution. IKK β is distributed predominantly in the cytoplasm, whereas IKK α has both nuclear and cytoplasmic localization. The cytoplasmic localization of IKKs is related to their ability to phosphorylate I κ B α and the consequent RelA/p50 translocation to the nucleus. The presence of IKK α in the nucleus is related to the expression of NF- κ B responsive genes through the phosphorylation and acetylation of histone 3 (Yamamoto and others 2003). These findings demonstrated that IKK α and IKK β kinases have different functions based on their cellular distribution.

NF- κ B Signaling Pathways: Canonical, Non-Canonical, and Atypical

In the nervous system, the NF- κ B pathway can be activated in a variety of ways. The classical IKK-dependent mechanisms include the canonical and non-canonical pathways, but a new IKK-independent mechanism, the atypical pathway, has been described (Bender and others 1998; Kato and others 2003). Canonical and non-canonical pathways are usually distinguished by two main characteristics: the NF- κ B dimer translocated to the nucleus (RelA/p50 and RelB/p52, respectively) and the I κ B contribution to their activation (I κ B dependent and I κ B independent, respectively) (Heissmeyer and others 1999). However, both of them require the presence of the IKK complex for their activation. In contrast, the atypical pathway is IKK-independent but I κ B dependent and induces RelA/p50 nuclear translocation (Perkins 2007).

The proteins and NF- κ B dimers involved in the activation of the three different pathways are summarized in Figure 3 and the principal differences that characterize these pathways in Figure 4. The “canonical pathway” (also called classical pathway) is the most common form of NF- κ B activation in all cell types. This form is characterized by the activation of dimers composed of p50 and RelA or c-Rel. In mammalian cells, the most abundant partner of p50 is RelA. The canonical pathway is generally activated in response to stimuli such as cytokines (TNF- α , TNF- β , IL-1, CNTF, or CT-1) (Barger and others 1995; Middleton and others 2001; Sparacio and others 1992), neurotrophins (Burke and Bothwell 2003), or oxygen-glucose deprivation (Sarnico and others 2009). The activation of this pathway depends on Ser181 and Ser180 phosphorylation of IKK α and IKK β , respectively. Activated IKK-complex phosphorylates I κ B α (Ser32 and/or Ser36), promoting its degradation by the proteasome 26S, and also induces p105 phosphorylation, which in turn promotes the generation of the mature form, p50 (Heissmeyer and others 1999). RelA/p50 heterodimers

are released from the inhibitor and translocated to the nucleus, where they bind to the DNA κ B sites and induce the activation or repression of specific genes.

TNF- α is the most powerful activator of the canonical pathway. TNF associate factor-2 (TRAF-2) is the adaptor protein recruited to the TNFR1 receptor and becomes responsible for the canonical pathway activation. Together with the cellular inhibitor of apoptosis-1/2 (c-IAP1/2), TRAF2 contributes to the polyubiquitination of RIP1, which in turn activates RelA/p50 through the IKK complex. Activation of NF- κ B by TNF- α usually induces protection from cell death, which promotes the transcription of anti-apoptotic target genes such as c-IAP1 and TRAF2 (Chu and others 1997; Wang and others 1998).

IKK α and IKK β have different effects on the canonical NF- κ B pathway activation. IKK β is the predominant kinase responsible for I κ B α and p105 phosphorylation (Li and others 1999c). Studies using IKK β knockout mice show a clear reduction of I κ B α degradation and RelA activation (Li and others 2003). In addition, these mice have a phenotype similar to RelA knockout, reinforcing the central role of IKK β in the canonical pathway activation (Beg and Baltimore 1996; Li and others 1999a; Li and others 1999b). Even though IKK α phosphorylation also stimulates the canonical NF- κ B pathway, this activation occurs in response to certain stimuli such as IL-1 or TNF- α when IKK β is inhibited (Lam and others 2008; Solt and others 2007).

The “atypical pathway” is IKK independent but I κ B α dependent and promotes RelA/p50 nuclear translocation (Fig. 4). In the nervous system, the activation of this pathway has been related to stimuli such as hydrogen peroxide, erythropoietin, or neurotrophic factors (Bui and others 2001; Gallagher and others 2007; Takada and others 2003). The atypical NF- κ B pathway is initiated by tyrosine (Tyr42) phosphorylation at the N-terminus of the I κ B α inhibitor or by serine phosphorylation at its PEST domain (Bender and others 1998; Kato and others 2003; Schwarz and others 1996). I κ B α phosphorylation at Tyr42 is mediated by Syk (spleen tyrosine kinase) in response to CNTF or NGF (Bui and others 2001; Gallagher and others 2007) or by members of the Src family of tyrosine kinases in response to BDNF stimulation (Gavaldà and others 2004). This phosphorylation leads to the release of I κ B α from the RelA/p50 dimer. Liberated I κ B α is not degraded by the proteasome as it occurs during canonical pathway activation (Bui and others 2001; Takada and others 2003). I κ B α can also be serine-phosphorylated by CKII (casein kinase II) at Ser293 located in the PEST domain (Schwarz and others 1996). In contrast to the tyrosine phosphorylation of I κ B α , serine phosphorylation promotes calpain-mediated I κ B α degradation (Wei 2009).

The “non-canonical pathway,” also known as the I κ B-independent pathway, is characterized by the translocation




Figure 3. NF-κB activation pathways. Schematic representation of the three different pathways to activate NF-κB: canonical, non-canonical, and atypical. The canonical and non-canonical are IKK dependent and the atypical depends on casein kinase II (CKII) or tyrosine kinases. Canonical and atypical activation are mediated by IκB phosphorylation in serine or tyrosine residues, inducing IκB release of the complex (Ser293 or Tyr42). IκB phosphorylated at Ser293 is degraded by the proteasome or by the protease calpain, and Tyr42 phosphorylation induces IκB dissociation of the complex without protein degradation. The non-canonical pathway is IκB independent. Its activation is due to p100 phosphorylation induced by NF-κB-inducing kinase (NIK) and IKKα activity. IκBα release or the proteasome processing of p100 unmasks RelA or RelB nuclear localization sequence (NLS), respectively, promoting their translocation to the nucleus. Canonical and atypical activation induce gene expression or repression by the nuclear translocation of the RelA/p50 heterodimer, whereas the non-canonical activation exerts its nuclear function through RelB/p52 translocation.

of the RelB/p52 heterodimer (Fig. 4). The activation of this pathway is mainly mediated by IKKα activity induced by the NF-κB-inducing kinase (NIK) (Bonizzi and others 2004). IKKα phosphorylation leads to a polyubiquitination-dependent degradation of the p100 precursor to the active form p52. The newly formed heterodimers RelB/

p52 translocate to the nucleus, where they target κB elements activating genes related to cellular functions, including Cox-2, Cyclin D, Mn-SOD, and Bcl-x_L (Holley and others 2010; Jacque and others 2005; Maehara and others 2000; Zhang and others 2007). This pathway is activated by a limited number of stimuli,




Figure 4. Schematic description of the principal differences of NF- κ B pathways activation. The convergence and divergence points of the different mechanisms to activate NF- κ B pathways are summarized: non-canonical and canonical are IKK dependent, atypical is IKK independent; canonical and atypical are I κ B dependent, non-canonical is I κ B independent; canonical and atypical induce RelA/p50 nuclear translocation, non-canonical induces RelB/p52 nuclear translocation.

including lymphotoxin B, CD40 ligand, LPS, and neurotrophic factors (Müller and Siebenlist 2003; Bhattacharyya and others 2010).

Regulation of the Transcriptional Activity of NF- κ B by Posttranslational Modifications of RelA

In addition to I κ B α inhibitor degradation, other steps are involved in the control of NF- κ B-mediated gene expression. Several studies have described posttranslational modifications of the NF- κ B members containing TAD domain—RelA, RelB, and c-Rel—that affect both RHD and TAD domains (Fig. 5). When RelA is liberated from the inhibitor I κ B α , it can be phosphorylated at several serine residues by different kinases, promoting conformational changes and favoring RelA binding to co-activators. For example, phosphorylation at Ser276 of the RHD domain by PKA, MSK1, and MSK2 (mitogen- and stress-activated protein kinase 1 and 2) promotes RelA interaction with the transcriptional co-activators CBP (CREB binding protein) and p300 (Olson and others 2007; Zhong and others 1998). PKA and MSK kinases are in turn regulated by the ERK/MAPK signaling pathway in response to different stimuli, including TNF- α (Vermeulen and others 2003). Otherwise, Ser311 phosphorylation of the RHD by PKC ξ regulates CBP and RelA interaction (Duran and others 2003). Within the TAD domain there are several serine residues susceptible to phosphorylation by kinases:

Ser468 phosphorylation occurs predominantly within the nucleus and is induced by the GSK-3 β kinase (Buss and others 2004; Schwabe and Brenner 2002); Ser529 is phosphorylated by the CKII kinase when RelA is liberated from the I κ B α inhibitor in response to IL-1 or TNF- α (Wang and others 2000); Ser535 can be phosphorylated by CAMKIV (Bae and others 2003) and Ser536 by IKKs in response to cytokines and mediated by the PI 3-kinase/Akt pathway (Gutierrez and others 2008; Sizemore and others 2002).

RelA also can be modified by reversible acetylation at different lysine residues. This site-specific acetylation regulates distinct biological activities of the NF- κ B complex. For example, acetylation at Lys310, Lys314, and Lys315 is required for full transcriptional RelA activity (Chen and others 2005b; Rothgiesser and others 2010a; Rothgiesser and others 2010b). However, acetylation at Lys122 and Lys123 exerts negative effects on NF- κ B-mediated transcription (Kiernan and others 2003).

Crosstalk between NF- κ B and Other Transcription Factors

It has been reported that some transcription factors can regulate or be regulated by IKK and/or NF- κ B. Tumor suppressor p53 can promote NF- κ B activation, but this does not occur through the classical activation of the IKKs. The expression of p53 stimulates the ribosomal serine/threonine kinase RSK1, which in turn phosphorylates the RelA subunit of NF- κ B at Ser536. RSK1-phosphorylated RelA is retained into the nucleus and contributes to the pro-apoptotic function of p53 (Bohuslav and others 2004). Thus, p53 and NF- κ B cooperatively induce apoptotic cell death in damaged cells including neurons (Aleyasin and others 2004). On the other hand, RelA activates p53 promoter in response to stress and suppresses cell growth (Wu and Lozano 1994). In microglial cells, a recent study demonstrates the interaction between NF- κ B and FOXO3a during oxygen-glucose deprivation (OGD) (Shang and others 2010). Using knockdown strategies, these studies pointed out that FoxO3a reduction facilitates p65 translocation to the nucleus in microglia during oxidative stress and promotes microglial cell survival. Finally, a novel transcriptional mechanism involving NF- κ B, PKA, and CREB has been described. NF- κ B controls PKA expression and consequently CREB activation; this mechanism is essential to regulate memory formation (Kaltschmidt and others 2006).

The NF- κ B Knockout Mice

Studies using knockout mice for different gene components of the NF- κ B pathway have improved our understanding of the role of these proteins in the physiological




Figure 5. Posttranslational modifications of RelA. Several posttranslational modifications induce increased RelA transcriptional activity. Phosphorylation at different serine residues located in Rel homology domain (RHD) or in transcriptional activation domain (TAD) is mediated by protein kinases from different families. Ser276 phosphorylation at RHD is mainly mediated by ERK/MAPK through the mitogen- and stress-activated kinase-1 (MSK1/2) and protein kinase A (PKA), and Ser311 phosphorylation is mediated by protein kinase C-β (PKCξ). Phosphorylation at TAD is mediated by the following kinases: glycogen synthase kinase-3β (GSK3β) at Ser468, casein kinase II (CKII) at Ser529, Ca²⁺/Calmodulin-Dependent Protein Kinase IV (CaMKIV) at Ser535, and phosphatidylinositol 3-kinase (PI3K)/protein kinase B (Akt) signaling pathway induce activation of IκB kinases (IKKs), which mediate Ser536 phosphorylation of RelA. Acetylation of lysines located outside of RHD or TAD are mediated by CREB binding protein (CBP) or p300 co-activators.

Table 1. Phenotype Characteristics of Knockout Mice of NF-κB Pathway Members

Knockout mice	Phenotype/Alterations	References
RelA	Embryonic death at E15-16; sensitivity to TNF-α; apoptotic death of hepatocytes, defects in the nervous system	(Beg and others 1995; Middleton and others 2000; Niscols and others 2003)
RelB	Multiple pathological lesions; defects in the development of T cell death	(Weih and others 1995; Yilmaz and others 2003)
c-Rel	Defects in lymphocyte proliferation, humoral immunity, neuronal survival, synaptic plasticity	(Ahn and others 2008; Deenick and others 2010; Grumont and others 1998; Köntgen and others 1995; Pizzi and others 2002)
p50	Defects in the immune response; neuronal degeneration	(Lu and others 2006; Sha and others 1995; Yu and others 2000)
p52	Lymphatic nodes abnormality and defects in T cell response	(Beinke and Ley 2004; Franzoso and others 1998)
IKKα	Death 1 day after birth; block the differentiation of keratinocytes and skeletal and epidermal defects	(Li and others 1999)
IKKβ	Embryonic death at E12.5-13.5; sensitivity to TNF-α and hepatic apoptosis; conditional survival of B cells, effects in the nervous system	(Bockhart and others 2009; Li and others 1999; Li and others 2003)
IKKγ (NEMO)	Embryonic death at E10-13; hepatic and B cell apoptosis and developmental defects of lymphocytes T and B cells	(Kim and others 2003; Rudolph and others 2000)

TNF-α = tumor necrosis factor-α.

processes related to development. In Table 1, we summarize the main features of these mice. RelA knockout die during embryonic development (E15-16) due to a massive liver apoptosis (Beg and others 1995). These mice also show defects in the nervous system, such as reduced survival of nodose neurons in response to CNTF and CT-1 cytokines (Middleton and others 2000). The lack of RelA also compromises peripheral myelin formation by Schwann cells (Nickols and others 2003). These findings point out the role of RelA in peripheral nervous system development affecting both neuronal and non-neuronal cells. To further determine the function of RelA during nervous system development, different knockdown strategies have been generated using embryonic neuronal models. These studies have demonstrated that RelA is also involved in neuronal survival of sensorial neurons and spinal cord motoneurons in response to neurotrophic factors (Hamanoue and others 1999; Mincheva and others 2011). RelB knockout mice have shown no defects during embryonic development (Weih and others 1995). These mice show defects in the secondary structure of lymphoid organs and abnormal hematopoiesis throughout life (Yilmaz and others 2003). There are no evidences indicating that RelB is involved in nervous system development. c-Rel knockout mice show defects in neuronal survival (Pizzi and others 2002) and synaptic plasticity related to memory formation (Ahn and others 2008). In the immune system, c-Rel knockout show impaired lymphocyte physiology (Köntgen and others 1995).

p50 knockout mice have a normal development and show changes in both specific and non-specific immune function (Grumont and others 1998; Sha and others 1995). They also show age-related degeneration of neuronal and non-neuronal cells, including caspase-3 activation and apoptotic cell death (Lu and others 2006), increased cell damage in response to the excitotoxic stimuli (Yu and others 1999), and increased apoptosis of striatal neurons in a Huntington disease model (Yu and others 2000). p52 knockout mice show a normal development and no changes in the nervous system have been described, but they have defects in the architecture of the lymphatic ganglia (Beinke and Ley 2004; Franzoso and others 1998).

IKK α knockout mice die shortly after birth because of skin and skeletal abnormalities caused by the blockade of keratinocyte differentiation (Li and others 1999b). No defects during nervous system development have been described in these mice. IKK β knockout embryos die because of a massive liver apoptosis between 12.5 and 13.5 embryonic days due to the reduction of RelA activation in the liver cells (Li and others 1999a). In the nervous system, the specific deletion of IKK β in sensory neurons of the dorsal root ganglia demonstrated that IKK β is a negative modulator of sensory neuron excitability (Bockhart and others 2009). IKK γ knockout mice

die during embryogenesis. Their phenotype is characterized by a massive hepatic apoptosis induced by TNF stimulation (Rudolph and others 2000) and impaired B cells development (Kim and others 2003), but no defects in nervous system development have been reported.

Summarizing the information provided by these knockout mice, we can conclude that RelA, c-Rel, and IKK β have an important role in nervous system development. However, we cannot completely discard the involvement of the rest of the members of the pathway in more specific functions that have not been tested yet. In this context, studies using double knockout strategies are providing new information. For example, IKK α and IKK β double knockout presents an excessive apoptosis in the neural tube, spinal cord, and dorsal root ganglia, leading to the abnormality of neural tube closure (Li and others 2000). Some other double knockout strategies have been generated, such as RelA/p50, but no evidences of neuronal abnormalities have been described in their phenotypes (Franzoso and others 1997; Franzoso and others 1998; Grossmann and others 2000; Horwitz and others 1997; Lo and others 2006; Weih and others 1997).

NF- κ B and the Nervous System

In the early 1990s, several studies showed the presence of RelA/p50 heterodimers in astrocytes (Sparacio and others 1992), Schwann cells (Carter and others 1996), microglia (Nakajima and Kohsaka 1998), and neurons (Kaltschmidt and others 1993; Meffert and others 2003; Schmidt-Ullrich and others 1996) in several regions of the developing and adult nervous system. These observations and subsequent reports suggest the involvement of NF- κ B in physiological processes of the nervous system during development and during adult life. This was the beginning of many studies dedicated to analyzing the functions of NF- κ B in the nervous system. In the following paragraphs, we review previous and recent results that contribute to this hypothesis, but we also reexamine the role of NF- κ B in nervous system pathologies. First of all, we will cite the specific activators of NF- κ B and the specific genes and proteins regulated by NF- κ B in the nervous system.

Activators of the NF- κ B Pathway in the Nervous System

There are many reports showing the specific activators of NF- κ B pathway in the nervous system (reviewed in Kaltschmidt and others 2005). NF- κ B is activated in both neuronal and non-neuronal cells, and the physiological response of these cells depends on the stimulus and the cellular origin. These responses include cell

Table 2. Neuroprotective Effects Induced by NF- κ B Pathway Activation in the Nervous System

NF- κ B activators	Cell type	Effects	Reference
CNTF	Sensory neurons	Survival and neurite growth	(Gallagher and others 2007)
CT-I	Sensory neurons	Survival	(Middleton and others 2000)
GDNF	Astrocytes	Neuroprotection in cerebral ischemia	(Chu and others 2008)
BDNF	Neurons	Survival and neurite growth	(Gutierrez and others 2005)
	PC12	Neurite outgrowth	(Sole and others 2004)
NGF	Neurons	Survival	(Maggirwar et al, 1998; Carter and others 1996)
	Schwann cells	Myelin formation	
IGF-I	Neurons	Survival	(Heck and others 1999)
Neurotrophic factors cocktail	Neurons	Survival	(Mincheva and others 2011)
TNF- α	Neurons	Synaptic plasticity	(Albensi and Mattson 2000)
cAMP	Schwann cell	Myelin formation	(Yoon and others 2008)

TNF- α = tumor necrosis factor- α .

Table 3. Neuronal degeneration NF- κ B Pathway Activation in the Nervous System

NF- κ B activators	Cell type	Effects	Reference
TNF- α	Neurons; glia	Neuronal degeneration and death	(Mir and others 2008)
Focal cerebral ischemia	Neurons	Neuronal death	(Stephenson and others 2000)
Amyloid	Astrocytes; oligodendrocytes	NO production increased; apoptosis	(Akama and others 1998; Xu and others 2001)
NMDA	Neurons	Apoptotic death	(Kitaoka and others 2004)
LPS	Microglia	Cox-2 expression	(Bauer and others 1997)
Kainic acid	Neurons	Apoptotic death	(Nakai and others 2000)
Oxidative stress	Neurons	Neuronal death	(Shou and others 2002)
Glutamate	Neurons	Apoptotic death	(Grilli and Memo 1999a)
IL-1 β	Neurons	Depressive-like behaviors	(Koo and others 2010)
H ₂ O ₂	Oligodendrocytes	Cell death	(Vollgraf and others 1999)

TNF- α = tumor necrosis factor- α ; IL-1 β = interleukin-1 β .

survival, apoptotic cell death, neurite outgrowth, neuronal differentiation and plasticity, or cell proliferation. Table 2 lists the stimuli that cause “positive effects” on neuronal and non-neuronal cells. These include survival, neurite outgrowth, differentiation, proliferation, and plasticity. For example, activation of NF- κ B mediated by neurotrophic factors promotes neuronal survival, neurite outgrowth, myelin formation, and axonal regeneration in various experimental models *in vivo* and *in vitro*, both in mature and developing cells (Mincheva and others 2011). However, NF- κ B activation can also induce “negative effects” such as cell death and toxicity (Akama and others 1998; Bauer and others 1997; Grilli and Memo 1999; Kitaoka and others

2004; Koo and others 2010; Mir and others 2008; Nakai and others 2000; Shou and others 2002; Stephenson and others 2000; Vollgraf and others 1999; Xu and others 2001). Table 3 lists some of the known stimuli that cause neuronal negative effects. For example, in Alzheimer disease (AD), β -amyloid protein induces NF- κ B activation in neurons, which produces cell toxicity because of nitric oxide production and release from astrocytes (Akama and others 1998). In fact, these different stimuli cause different events in the different types of cells present in the nervous system. This could explain the “apparently contradictory” effects caused by NF- κ B activity that were mentioned at the beginning of this review.

Genes and Proteins Regulated after NF- κ B Activation in the Nervous System

Independent studies over the past 10 years have proposed NF- κ B as one of the main pathways controlling neuronal survival. This regulation is based on the ability of NF- κ B to exert transcriptional control (activation or inhibition) over several pro- or anti-apoptotic genes (Table 4). For example, Bcl-2 and Bcl-x_L promoters have binding sites for NF- κ B dimers (Tamatani and others 1999) or NF- κ B positively regulates cIAP gene expression (Baud and Karin 2001). Moreover, TNFR1/2 adaptor proteins TRAF1 and TRAF2 increase when NF- κ B is activated, and TRAF1 and TRAF2 themselves are able to induce IKK phosphorylation, which in turn activates the NF- κ B pathway (Häcker and Karin 2006; Wang and others 1998). Along the same line, we have recently demonstrated that the NF- κ B pathway also regulates the level of one protein essential for motoneuron physiology, Survival Motor Neuron (SMN) (Mincheva and others 2011).

On the other hand, NF- κ B activation can promote apoptosis and cell death in the nervous system (Kaltschmidt and others 2000; Kaltschmidt and others 2002; Pizzi and others 2002). For example, in response to oxidative stress, NMDA receptor activation or apoptotic stimuli of the NF- κ B pathway increases the expression of the pro-apoptotic genes Bax and Bcl-x_s in cortical neurons (Shou and others 2002). In some diseases, such as cerebral ischemia, RelA increases the expression of Bim and Noxa pro-apoptotic genes (Inta and others 2006). Even though NO participation in neuronal apoptosis is still under debate, the transcriptional control of the inducible nitric oxide synthase (iNOS) is also regulated by NF- κ B (Xie and others 1994). The cell cycle regulator Cyclin D is also controlled by NF- κ B and has been related to neuronal apoptosis in striate and cortical neurons (Liang and others 2007).

Finally, under NF- κ B transcriptional control other factors related to neuronal homeostasis are also regulated, such as antioxidant enzymes (Maehara and others 2000; Rojo and others 2004), adhesion molecules (Simpson and Morris 2000), transcription factors (Khoroshii and others 2008; Qin and others 1999; Wu and Lozano 1994), and neurotrophic factors (Saha and others 2006).

Role of NF- κ B in the Nervous System during Development and in the Adult

Control of Apoptosis

As mentioned above, there are defects in neural tube closure in IKK α and IKK β double knockout caused by the apoptosis of the neuroepithelium (Li Q and others 2000). During embryogenesis NF- κ B transcriptional

activity is detected in the central nervous system, particularly in the spinal cord and brain nuclei (Schmidt-Ullrich and others 1996). One of the first observations of NF- κ B activity in the spinal cord coincides with the beginning of motoneuron neurotrophic factor dependence and the programmed cell death period (Schmidt-Ullrich and others 1996; Yeo and Gautier 2004). Our recent results demonstrated that reduction of some NF- κ B pathway members causes apoptotic cell death of embryonic spinal cord motoneurons, even in the presence of neurotrophic factors (Mincheva and others 2011). NF- κ B activation can also be required for peripheral nervous system neuronal survival. Studies using embryonic sympathetic and sensory neurons indicated that NF- κ B activation is necessary for NGF-induced cell survival (Maggirwar et al., 1998 and Mincheva-Tasheva unpublished results). All these observations highlight the important role of NF- κ B signaling pathway in the regulation of apoptotic cell death during neuronal development (Fig. 6).

Involvement in Neurite Outgrowth

NF- κ B signaling has been implicated in control of axon initiation, branching and elongation, and dendrite density in the adult. Studies using different neuronal models including PC12 cells (Azoitei and others 2005; Sole and others 2004), sensory neurons (Gallagher and others 2007; Gutierrez and others 2005; Gutierrez and others 2008), and hippocampal neurons (Sanchez-Ponce and others 2011) have demonstrated that NF- κ B blockade reduces neurite length and branching. A recent review extensively discusses the role of NF- κ B pathway in modulating the growth and morphology of neuronal processes, including axon and dendrites (Gutierrez and Davies 2011).

Involvement of NF- κ B in Synaptic Plasticity, Memory, and Learning

NF- κ B plays an important role in synaptic signaling and learning in the mature nervous system (Kaltschmidt and Kaltschmidt 2009). Independent genetic studies have demonstrated that RelA, c-Rel, and p50 are involved in memory formation. RelA/TNFR1 double knockout mice have defects in spatial learning memory (Meffert and others 2003), c-RelA knockouts have impaired hippocampus-dependent memory formation (Ahn and others 2008), and p50 knockout present deficit in short-term memory (Denis-Donini and others 2008). The importance of NF- κ B in these processes may be related to the presence and function of NF- κ B family members in the pre- and postsynaptic space, which contributes to transduction of synaptic signals to transcriptional changes (Meffert and others 2003). In these processes of learning and memory, NF- κ B also cooperates with other transcription factors and signaling pathways. For example, NF- κ B controls synaptic plasticity

Table 4. Target Genes Regulated by NF- κ B Pathways and their Cellular Effects in the Nervous System

Target genes	Pathway	Cellular function	References
<i>Bcl-2, Bcl-x</i>	C	Anti-apoptotic	(Tamatani and others 1999)
<i>Bcl-xL</i>	A, C, non-C	Anti-apoptotic	(Bui and others 2001)
<i>c-FLIP</i>	X	Anti-apoptotic	(Chu and others 1997)
<i>TRAF1, 2</i>	X	Anti-apoptotic	(Wang and others 1998)
<i>IAPs</i>	C	Anti-apoptotic	(Chu and others 1997)
<i>Bax, Bcl-xs</i>	X	Pro-apoptotic	(Shou and others 2002)
<i>Bim, Nova</i>	C	Pro-apoptotic	(Inta and others 2006)
<i>Smn</i>	C	Survival Motor Neuron	(Mincheva and others 2011)
<i>CREB</i>	C	Transcription factor	(Mincheva and others 2011)
<i>p53</i>	C	Transcription factor	(Wu and Lozano 1994)
<i>STAT2</i>	X	Transcription factor	(Khorrooshi and others 2008)
<i>c-Myc</i>	C	Transcription factor	(Qin and others 1999)
<i>Mn-SOD</i>	C	Antioxidant enzyme	(Maehara and others 2000)
<i>Cu/Zn-SOD</i>	C	Antioxidant enzyme	(Rojo and others 2004)
<i>Cox-2</i>	Non-C	Enzyme responsible for inflammation	(Kaltschmidt and others 2002)
<i>iNOS</i>	C	Nitric oxide synthesis	(Xie and others 1994)
<i>IκBα</i>	C	NF- κ B inhibitor	(Bui and others 2001)
<i>BDNF</i>	C	Neurotrophic factor	(Saha and others 2006)
<i>NCAM</i>	C	Neural adhesion molecule	(Simpson and Morris 2000)
<i>Cyclin D1</i>	Non-C	Cell cycle regulator	(Liang and others 2007)

C = canonical NF- κ B pathway; non-C = non-canonical NF- κ B pathway; A = atypical NF- κ B pathway; X = no information.

by regulating the expression of the catalytic subunit of PKA, an essential memory regulator, and phosphorylation of CREB (Kaltschmidt and others 2006).

NF- κ B and Neurodegenerative Disorders

To better understand the molecular mechanisms of the pathology of neurodegenerative disorders, it is important to understand the signal transduction processes that regulate neuronal survival and differentiation during development. These mechanisms could be involved in the pathology of these diseases, and there are evidences suggesting this hypothesis. Several studies associate the alteration of NF- κ B activity with neurodegenerative diseases such as Alzheimer, Parkinson, Huntington, or amyotrophic lateral sclerosis (ALS), and in processes related to injury or ischemia. In general terms, authors propose that NF- κ B activation in neurons exerts a neuroprotective role in the degenerative process of these diseases (Cardoso and Oliveira 2003; Fridmacher and others 2003; Smith and others 2009). However, there are also evidences indicating that it causes neuronal death. Analysis of brain samples showed an increase of activated NF- κ B in AD patients compared with healthy individuals (Kaltschmidt and others 1997; Kaltschmidt and others 1999). This NF- κ B activation can contribute to the pathological changes observed in AD-inducing proinflammatory and cytotoxic genes, or can be a part of

the cellular protection mechanism depending on the genetic program and the time of exposure to the stimulus. In mice models of AD (Tg2576), the increase of NF- κ B activity has been related to neuronal apoptosis and the onset and development of the disease (Niu and others 2010). Two independent studies have demonstrated the involvement of NF- κ B in A β 42 oligomer production (Buggia-Prevot and others 2008; Valerio and others 2006). Taking into consideration these reports, NF- κ B can be considered as a pharmacological target for AD treatment by directly inhibiting the production of A β peptides (Paris and others 2007). More detailed studies have shown that the specific inhibition of NF- κ B activity in microglia, but not in neurons, blocks A β -induced neurotoxicity (Chen and others 2005a). Along the same line and based on the ability of NF- κ B to block caspase activation in neurons, it has been proposed that the pharmacological induction of NF- κ B activity can be a beneficial therapeutic target in AD therapy (Cardoso and Oliveira 2003).

There are also evidences that NF- κ B may participate in Parkinson disease (PD) although its role in dopaminergic neurons survival is still controversial. Some authors have shown that the activation of NF- κ B pathway is essential for the apoptotic death induced by dopamine in PC12 cells (Panet and others 2001); and NF- κ B activation contributes to 6-hydroxydopamine-induced apoptosis of dopaminergic neurons (Li and others 2008). However, other authors suggest the neuroprotective role of NF- κ B




Figure 6. Effects of NF- κ B activation in neurons. (A) Confocal images of RelA subcellular localization in cultured motoneurons (MNs) in the absence (no NTFs) or the presence (NTFs) of neurotrophic factors. NTFs treatment induces RelA translocation to the nucleus indicating the activation of the pathway (RelA, red; Hoechst nuclear dye, blue). (B) and (C) show the effect of inactivating NF- κ B pathway using an RNA interference approach. (B) Phase contrast images of cultured dorsal root ganglion (DRG) neurons (left) and MNs (right) in the presence of NTFs under control (top) or RelA interference (bottom) conditions. (C) Nuclear apoptotic morphology of MNs cultured in RelA interference conditions (bottom).

activation based on its ability to block the apoptotic cell death caused by auto-oxidized dopamine (Lee and others 2001). In the same line of evidences, it has been described that the exogenous administration of the neurotrophic

factor GDNF induces RelA/p52 nuclear translocation and protects dopaminergic neurons in an early rat model of PD (Cao and others 2008).

Although not widely studied, NF- κ B pathway may also participate in Huntington disease. p50-deficient mice show an increased loss of striatal neurons and consequent motor dysfunction (Yu and others 2000), and in mammalian cells human huntingtin (Htt) co-immunoprecipitates with p50 (Takano and Gusella 2002). Other members of NF- κ B signaling can also contribute to neurodegeneration induced by mutated Htt. It has been reported that inhibition of IKK γ or IKK β activity, or I κ B α degradation, can reduce Htt-induced toxicity and promote striatal neuronal survival (Khoshnaw and others 2004). In later studies, the same authors demonstrated an antagonistic effect for the different IKKs. IKK α overexpression or IKK β inhibition protects neurons from death induced by mutant Htt (Khoshnaw and others 2009).

Indirect effects of NF- κ B pathway have also been related to neurodegenerative disorders of the spinal cord motoneurons. In spinal cords of ALS patients, NF- κ B is activated in glial cells but not in motoneurons (Migheli and others 1997; Pyo and others 2010). Studies using SOD transgenic mice demonstrated that pharmacological induction of NF- κ B pathway in motoneurons prolongs survival by blocking apoptosis (DeL Signore and others 2009; Ryu and others 2005). However, some authors suggest a beneficial effect of NF- κ B pathway pharmacological inhibitors in the spinal cord of ALS mutant mice (Xu and others 2006). From all these results, it is feasible to conclude that although NF- κ B activity might have a role in the pathology and therapy of these degenerative disorders, additional data are required to establish the clinical relevance, as we discuss below.

NF- κ B and Neuronal Injury

NF- κ B activation has been observed in response to brain and spinal cord ischemia and trauma (Xu and others 2005). The analysis of experimental models of stroke and injury reveals that the NF- κ B pathway may modulate neuronal degeneration or protection. There are several examples of these opposite effects. Mice lacking the p50 subunit develop significantly smaller infarct size (Nurmi and others 2004), and there are evidences that IKK/NF- κ B signaling contributes to ischemic brain damage (Schwaninger and others 2006). On the other hand, some data in hippocampus and striatum provide evidences that NF- κ B participates in survival signaling following temporary focal ischemia (Duckworth and others 2006). A recent study demonstrates that transgenic inhibition of glial NF- κ B attenuates pain and the inflammatory response following chronic constriction of the sciatic nerve (Fu and others 2010).




Figure 7. Hypothesis for NF- κ B function in nervous system physiology and pathology. A) Model for NF- κ B homeostasis suggested by Kaltschmidt. This model proposes that under physiological conditions RelA binds to nuclear co-activators which promote cell survival by activating the transcription of anti-apoptotic genes. RelA-induced I κ B protein expression controls NF- κ B homeostasis inactivating the pathway. Under pathologic conditions deregulation of NF- κ B activity leads RelA binding to nuclear co-repressors blocking the anti-apoptotic gene expression and promotes neuronal cell death. B) Opposite effects of NF- κ B activation on neuronal and on glial cells under pathologic conditions. As suggested by Mattson and Camandola's hypothesis, the same NF- κ B activating stimulus can induce opposite effects on neurons. NF- κ B activation in neurons under some pathologic conditions such as injury or stroke promotes the expression of anti-apoptotic genes, neurotrophic factors, or adhesion molecules, which in turn have a neuroprotection role. The same pathological stimuli in astrocytes or microglia induce the expression of nitric oxide, TNF- α , and/or excitatory amino acids, leading to neuronal degeneration. Representative images of the NF- κ B activation effect on motoneurons (MNs) (left) and glial cells (right) when both types of cells are cocultured: NF- κ B activation in MNs induces cell survival (arrows) whereas NF- κ B activation in glial cells induces cell death.

All together these results illustrate the controversial role of NF- κ B pathway in neuronal damage and disorders. To find an answer to these apparently contradictory results, two hypotheses have been generated. The most recent, proposed by Kaltschmidt in 2005, is the model for NF- κ B homeostasis (Kaltschmidt and others 2005) (Fig. 7A). This model defends the premise that low levels or high levels of NF- κ B activation maintained over a long time period (weeks to months) cause neuronal death. Under physiological conditions, nuclear RelA promotes survival by activating the transcription of anti-apoptotic genes through its binding to co-activators. The maintenance of "physiological" NF- κ B activation depends on

I κ B protein expression that is responsible for bringing RelA back to the cytoplasm and deactivating the pathway. However, pathologic conditions can lead to low or hyperactivation of NF- κ B, which can switch RelA from a selective promoter to a dominant repressor of anti-apoptotic genes by recruiting corepressors and inducing neuronal cell death. The more supported hypothesis was proposed by Mattson and Camandola in 2001 (Mattson and Camandola 2001). These authors suggest that NF- κ B activation in neurons induces anti-apoptotic genes that mediate cell survival, whereas NF- κ B activation in glial cells results in the production of proinflammatory cytokines that mediate neuronal death (Fig. 7B). Thus, the same stimulus produces opposite responses by activating NF- κ B in different cell types. One of the cytokines that can be produced by glial cells is TNF- α . Binding of TNF- α to TNFR1 and TNFR2 receptor activates the NF- κ B pathway. TNFR1 activation in glial cells induces the production of NO, which may lead to neuronal death, whereas TNFR2 activation in neurons increases the expression of anti-apoptotic genes. This hypothesis has been supported by several studies demonstrating that TNF- α toxicity in neurons is only observed in the presence of glial cells (Mir and others 2008; Taylor and others 2005; Tolosa and others 2011). These studies showed that microglia, macrophages, and astrocytes produce proinflammatory cytokines, free radicals, and excitotoxins in response to TNF- α -induced NF- κ B activation and thereby promote neuronal death. Furthermore, in neuronal injury models the inhibition of the pathway only in astrocytes demonstrated that neurons are protected from cell death when NF- κ is silenced in astrocytes (Meunier and others 2007). Similar results were obtained in hippocampal neurons in response to kainic acid excitotoxicity, when the *IKK β* gene was specifically deleted in microglia (Cho and others 2008). Based on these observations, the use of glial-specific NF- κ B pathway inhibitors has been proposed as a potential therapy in neurodegenerative disorders such as AD (Chen and others 2005a) or ALS (Crosio and others 2011).

Concluding Remarks

This review describes a number of important functions of the NF- κ B pathway in the nervous system. The pathway's function in the regulation of apoptosis, neurite outgrowth, and synaptic plasticity during development and memory formation and learning in adult life has been well studied; however, new evidences are emerging about its role in neurodegenerative disorders and neuronal injury. Although some of the results could lead to confusion about the function of the pathway during these processes, these recent studies confirm Mattson and Camandola's hypothesis concerning the importance of the stimuli, context, and cellular origin

of NF- κ B activation, but among the context and the stimuli, cell type appears to be the decisive factor determining NF- κ B function during development and neurodegeneration. It remains to be determined whether NF- κ B up- or down-regulation in a neuron under pathologic conditions is the origin of cell degeneration or is a neuroprotective reaction. When these issues have been clarified, NF- κ B activity regulation can be proposed as a therapeutic target in some neurodegenerative disorders.

Acknowledgment

We thank Elaine Lilly, PhD (Writer's First Aid), for English language revision of the article.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by grants to RMS from Instituto de Salud Carlos III, Fondo de Investigaciones Sanitarias (PI080267/PI1101047), and Consolider-Ingenio 2010 (CSD2007-00020, Ministerio de Ciencia e Innovación), and by the Generalitat de Catalunya (Suport Grups de Recerca Emergents) 2009SGR740. SMT holds a fellowship from Consolider-Ingenio 2010 (CSD2007-00020).

References

- Ahn HJ, Hernandez CM, Levenson JM, Lubin FD, Liou HC, Sweatt JD. 2008. c-Rel, an NF-kappaB family transcription factor, is required for hippocampal long-term synaptic plasticity and memory formation. *Learn Mem* 15:539–49.
- Airaksinen MS, Saarma M. 2002. The GDNF family: signaling, biological functions and therapeutic value. *Nat Rev Neurosci* 3:383–94.
- Akama KT, Albanese C, Pestell RG, Van Eldik LJ. 1998. Amyloid beta-peptide stimulates nitric oxide production in astrocytes through an NFkappaB-dependent mechanism. *Proc Natl Acad Sci U S A* 95:5795–800.
- Aleyasin H, Cregan SP, Iyirhiano G, O'Hare MJ, Callaghan SM, Slack RS, and others. 2004. Nuclear factor-(kappa)B modulates the p53 response in neurons exposed to DNA damage. *J Neurosci* 24:2963–73.
- Azoitei N, Wirth T, Baumann B. 2005. Activation of the IkappaB kinase complex is sufficient for neuronal differentiation of PC12 cells. *J Neurochem* 93:1487–501.
- Bae JS, Jang MK, Hong S, An WG, Choi YH, Kim HD, and others. 2003. Phosphorylation of NF-kappa B by calmodulin-dependent kinase IV activates anti-apoptotic gene expression. *Biochem Biophys Res Commun* 305:1094–8.
- Baeuerle PA. 1998. IkappaB-NF-kappaB structures: at the interface of inflammation control. *Cell* 95:729–31.
- Barger SW, Hörster D, Furukawa K, Goodman Y, Kriegstein J, Mattson MP. 1995. Tumor necrosis factors alpha and beta protect neurons against amyloid beta-peptide toxicity: evidence for involvement of a kappa B-binding factor and attenuation of peroxide and Ca²⁺ accumulation. *Proc Natl Acad Sci U S A* 92:9328–32.
- Baud V, Karin M. 2001. Signal transduction by tumor necrosis factor and its relatives. *Trends Cell Biol* 11:372–7.
- Bauer MK, Lieb K, Schulze-Osthoff K, Berger M, Gebicke-Haerter PJ, Bauer J, and others. 1997. Expression and regulation of cyclooxygenase-2 in rat microglia. *Eur J Biochem* 243:726–31.
- Beg AA, Baltimore D. 1996. An essential role for NF-kappaB in preventing TNF-alpha-induced cell death. *Science* 274:782–4.
- Beg AA, Sha WC, Bronson RT, Ghosh S, Baltimore D. 1995. Embryonic lethality and liver degeneration in mice lacking the RelA component of NF-kappa B. *Nature* 376:167–70.
- Beinke S, Ley SC. 2004. Functions of NF-kappaB1 and NF-kappaB2 in immune cell biology. *Biochem J* 382:393–409.
- Bender K, Göttlicher M, Whiteside S, Rahmsdorf HJ, Herrlich P. 1998. Sequential DNA damage-independent and -dependent activation of NF-kappaB by UV. *EMBO J* 17:5170–81.
- Bhattacharyya S, Borthakur A, Dudeja PK, Tobacman JK. 2010. Lipopolysaccharide-induced activation of NF-kB non-canonical pathway requires BCL10 serine 138 and NIK phosphorylations. *Exp Cell Res* 316:3317–27.
- Bockhart V, Constantin CE, Häussler A, Wijnvoord N, Kanngiesser M, Myrczek T, and others. 2009. Inhibitor kappaB kinase beta deficiency in primary nociceptive neurons increases TRP channel sensitivity. *J Neurosci* 29:12919–29.
- Bohuslav J, Chen LF, Kwon H, Mu Y, Greene WC. 2004. p53 induces NF-kappaB activation by an IkappaB kinase-independent mechanism involving phosphorylation of p65 by ribosomal S6 kinase 1. *J Biol Chem* 279:26115–25.
- Bonizzi G, Bebiën M, Otero DC, Johnson-Vroom KE, Cao Y, Vu D, and others. 2004. Activation of IKKalpha target genes depends on recognition of specific kappaB binding sites by RelB:p52 dimers. *EMBO J* 23:4202–10.
- Buggia-Prevot V, Sevalle J, Rossner S, Checler F. 2008. NFkappaB-dependent control of BACE1 promoter transactivation by Abeta42. *J Biol Chem* 283:10037–47.
- Bui NT, Livolsi A, Peyron JF, Prehn JH. 2001. Activation of nuclear factor kappaB and Bcl-x survival gene expression by nerve growth factor requires tyrosine phosphorylation of IkappaBalpha. *J Cell Biol* 152:753–64.
- Burke MA, Bothwell M. 2003. p75 neurotrophin receptor mediates neurotrophin activation of NF-kappa B and induction of iNOS expression in P19 neurons. *J Neurobiol* 55:191–203.
- Buss H, Dörrie A, Schmitz ML, Frank R, Livingstone M, Resch K, and others. 2004. Phosphorylation of serine 468 by GSK-

- 3beta negatively regulates basal p65 NF-kappaB activity. *J Biol Chem* 279:49571–4.
- Cao JP, Wang HJ, Yu JK, Liu HM, Gao DS. 2008. The involvement of NF-kappaB p65/p52 in the effects of GDNF on DA neurons in early PD rats. *Brain Res Bull* 76:505–11.
- Cardoso SM, Oliveira CR. 2003. Inhibition of NF-kB renders cells more vulnerable to apoptosis induced by amyloid beta peptides. *Free Radic Res* 37:967–73.
- Carter BD, Kaltschmidt C, Kaltschmidt B, Offenhäuser N, Böhm-Matthaei R, Baeuerle PA, and others. 1996. Selective activation of NF-kappa B by nerve growth factor through the neurotrophin receptor p75. *Science* 272:542–5.
- Chao MV. 2003. Neurotrophins and their receptors: a convergence point for many signalling pathways. *Nat Rev Neurosci* 4:299–309.
- Chen J, Zhou Y, Mueller-Steiner S, Chen LF, Kwon H, Yi S, and others. 2005a. SIRT1 protects against microglia-dependent amyloid-beta toxicity through inhibiting NF-kappaB signaling. *J Biol Chem* 280:40364–74.
- Chen LF, Williams SA, Mu Y, Nakano H, Duerr JM, Buckbinder L, and others. 2005b. NF-kappaB RelA phosphorylation regulates RelA acetylation. *Mol Cell Biol* 25:7966–75.
- Chen Z, Hagler J, Palombella VJ, Melandri F, Scherer D, Ballard D, and others. 1995. Signal-induced site-specific phosphorylation targets I kappa B alpha to the ubiquitin-proteasome pathway. *Genes Dev* 9:1586–97.
- Cho IH, Hong J, Suh EC, Kim JH, Lee H, Lee JE, and others. 2008. Role of microglial IKKbeta in kainic acid-induced hippocampal neuronal cell death. *Brain* 131:3019–33.
- Chu LF, Wang WT, Ghanta VK, Lin CH, Chiang YY, Hsueh CM. 2008. Ischemic brain cell-derived conditioned medium protects astrocytes against ischemia through GDNF/ERK/NF-kB signaling pathway. *Brain Res* 1239:24–35.
- Chu ZL, McKinsey TA, Liu L, Gentry JJ, Malim MH, Ballard DW. 1997. Suppression of tumor necrosis factor-induced cell death by inhibitor of apoptosis c-IAP2 is under NF-kappaB control. *Proc Natl Acad Sci U S A* 94:10057–62.
- Crosio C, Valle C, Casciati A, Iaccarino C, Carri MT. 2011. Astroglial inhibition of NF-kB does not ameliorate disease onset and progression in a mouse model for amyotrophic lateral sclerosis (ALS). *PLoS One* 6:e17187.
- Del Signore SJ, Amante DJ, Kim J, Stack EC, Goodrich S, Cormier K, and others. 2009. Combined riluzole and sodium phenylbutyrate therapy in transgenic amyotrophic lateral sclerosis mice. *Amyotroph Lateral Scler* 10:85–94.
- Delhase M, Hayakawa M, Chen Y, Karin M. 1999. Positive and negative regulation of IkappaB kinase activity through IKKbeta subunit phosphorylation. *Science* 284:309–13.
- Denis-Donini S, Dellarole A, Crociara P, Francese MT, Bortolotto V, Quadrato G, and others. 2008. Impaired adult neurogenesis associated with short-term memory defects in NF-kappaB p50-deficient mice. *J Neurosci* 28:3911–9.
- Duckworth EA, Butler T, Collier L, Collier S, Pennypacker KR. 2006. NF-kappaB protects neurons from ischemic injury after middle cerebral artery occlusion in mice. *Brain Res* 1088:167–75.
- Duran A, Diaz-Meco MT, Moscat J. 2003. Essential role of RelA Ser311 phosphorylation by zetaPKC in NF-kappaB transcriptional activation. *EMBO J* 22:3910–8.
- Ernst MK, Dunn LL, Rice NR. 1995. The PEST-like sequence of I kappa B alpha is responsible for inhibition of DNA binding but not for cytoplasmic retention of c-Rel or RelA homodimers. *Mol Cell Biol* 15:872–82.
- Franzoso G, Carlson L, Poljak L, Shores EW, Epstein S, Leonardi A, and others. 1998. Mice deficient in nuclear factor (NF)-kappa B/p52 present with defects in humoral responses, germinal center reactions, and splenic microarchitecture. *J Exp Med* 187:147–59.
- Franzoso G, Carlson L, Xing L, Poljak L, Shores EW, Brown KD, and others. 1997. Requirement for NF-kappaB in osteoclast and B-cell development. *Genes Dev* 11:3482–96.
- Fridmacher V, Kaltschmidt B, Goudeau B, Ndiaye D, Rossi FM, Pfeiffer J, and others. 2003. Forebrain-specific neuronal inhibition of nuclear factor-kappaB activity leads to loss of neuroprotection. *J Neurosci* 23:9403–8.
- Fu ES, Zhang YP, Sagen J, Candiotti KA, Morton PD, Liebl DJ, and others. 2010. Transgenic inhibition of glial NF-kappa B reduces pain behavior and inflammation after peripheral nerve injury. *Pain* 148:509–18.
- Gallagher D, Gutierrez H, Gavalda N, O’Keeffe G, Hay R, Davies AM. 2007. Nuclear factor-kappaB activation via tyrosine phosphorylation of inhibitor kappaB-alpha is crucial for ciliary neurotrophic factor-promoted neurite growth from developing neurons. *J Neurosci* 27:9664–9.
- Gavalda N, Pérez-Navarro E, Gratacòs E, Comella J, Alberch J. 2004. Differential involvement of phosphatidylinositol 3-kinase and p42/p44 mitogen activated protein kinase pathways in brain-derived neurotrophic factor-induced trophic effects on cultured striatal neurons. *Mol Cell Neurosci* 25:460–8.
- Ghosh S, May MJ, Kopp EB. 1998. NF-kappa B and Rel proteins: evolutionarily conserved mediators of immune responses. *Annu Rev Immunol* 16:225–60.
- Grilli M, Memo M. 1999. Nuclear factor-kappaB/Rel proteins: a point of convergence of signalling pathways relevant in neuronal function and dysfunction. *Biochem Pharmacol* 57:1–7.
- Grossmann M, O’Reilly LA, Gugasyan R, Strasser A, Adams JM, Gerondakis S. 2000. The anti-apoptotic activities of Rel and RelA required during B-cell maturation involve the regulation of Bcl-2 expression. *EMBO J* 19:6351–60.
- Grumont RJ, Rourke IJ, O’Reilly LA, Strasser A, Miyake K, Sha W, and others. 1998. B lymphocytes differentially use the Rel and nuclear factor kappaB1 (NF-kappaB1) transcription factors to regulate cell cycle progression and apoptosis in quiescent and mitogen-activated cells. *J Exp Med* 187:663–74.

- Gutierrez H, Davies AM. 2011. Regulation of neural process growth, elaboration and structural plasticity by NF- κ B. *Trends Neurosci* 34:316–25.
- Gutierrez H, Hale VA, Dolcet X, Davies A. 2005. NF-kappaB signalling regulates the growth of neural processes in the developing PNS and CNS. *Development* 132:1713–26.
- Gutierrez H, O'Keeffe GW, Gavalda N, Gallagher D, Davies AM. 2008. Nuclear factor kappa B signaling either stimulates or inhibits neurite growth depending on the phosphorylation status of p65/RelA. *J Neurosci* 28:8246–56.
- Häcker H, Karin M. 2006. Regulation and function of IKK and IKK-related kinases. *Sci STKE* 2006:re13.
- Hamanoue M, Middleton G, Wyatt S, Jaffray E, Hay RT, Davies AM. 1999. p75-mediated NF-kappaB activation enhances the survival response of developing sensory neurons to nerve growth factor. *Mol Cell Neurosci* 14:28–40.
- Heck S, Lezoualc'h F, Engert S, Behl C. 1999. Insulin-like growth factor-I-mediated neuroprotection against oxidative stress is associated with activation of nuclear factor kappaB. *J Biol Chem* 274:9828–35.
- Heissmeyer V, Krappmann D, Wulczyn FG, Scheidereit C. 1999. NF-kappaB p105 is a target of IkappaB kinases and controls signal induction of Bcl-3-p50 complexes. *EMBO J* 18:4766–78.
- Holley AK, Xu Y, St Clair DK, St Clair WH. 2010. RelB regulates manganese superoxide dismutase gene and resistance to ionizing radiation of prostate cancer cells. *Ann N Y Acad Sci* 1201:129–36.
- Horwitz BH, Scott ML, Cherry SR, Bronson RT, Baltimore D. 1997. Failure of lymphopoiesis after adoptive transfer of NF-kappaB-deficient fetal liver cells. *Immunity* 6:765–72.
- Huxford T, Malek S, Ghosh G. 1999. Structure and mechanism in NF-kappa B/I kappa B signaling. *Cold Spring Harb Symp Quant Biol* 64:533–40.
- Huynh QK, Boddupalli H, Rouw SA, Koboldt CM, Hall T, Sommers C, and others. 2000. Characterization of the recombinant IKK1/IKK2 heterodimer. Mechanisms regulating kinase activity. *J Biol Chem* 275:25883–91.
- Imbert V, Rupec RA, Livolsi A, Pahl HL, Traenckner EB, Mueller-Dieckmann C, and others. 1996. Tyrosine phosphorylation of I kappa B-alpha activates NF-kappa B without proteolytic degradation of I kappa B-alpha. *Cell* 86:787–98.
- Inta I, Paxian S, Maegele I, Zhang W, Pizzi M, Spano P, and others. 2006. Bim and Noxa are candidates to mediate the deleterious effect of the NF-kappa B subunit RelA in cerebral ischemia. *J Neurosci* 26:12896–903.
- Ito CY, Kazantsev AG, Baldwin AS. 1994. Three NF-kappa B sites in the I kappa B-alpha promoter are required for induction of gene expression by TNF alpha. *Nucleic Acids Res* 22:3787–92.
- Jacque E, Tchenio T, Piton G, Romeo PH, Baud V. 2005. RelA repression of RelB activity induces selective gene activation downstream of TNF receptors. *Proc Natl Acad Sci U S A* 102:14635–40.
- Kaltschmidt B, Heinrich M, Kaltschmidt C. 2002. Stimulus-dependent activation of NF-kappaB specifies apoptosis or neuroprotection in cerebellar granule cells. *Neuromolecular Med* 2:299–309.
- Kaltschmidt B, Kaltschmidt C. 2009. NF-kappaB in the nervous system. *Cold Spring Harb Perspect Biol* 1:a001271.
- Kaltschmidt B, Widera D, Kaltschmidt C. 2005. Signaling via NF-kappaB in the nervous system. *Biochim Biophys Acta* 1745:287–99.
- Kaltschmidt B, Uherek M, Volk B, Baeuerle PA, Kaltschmidt C. 1997. Transcription factor NF-kappaB is activated in primary neurons by amyloid beta peptides and in neurons surrounding early plaques from patients with Alzheimer disease. *Proc Natl Acad Sci U S A* 94:2642–7.
- Kaltschmidt B, Uherek M, Wellmann H, Volk B, Kaltschmidt C. 1999. Inhibition of NF-kappaB potentiates amyloid beta-mediated neuronal apoptosis. *Proc Natl Acad Sci U S A* 96:9409–14.
- Kaltschmidt B, Kaltschmidt C, Hofmann TG, Hehner SP, Dröge W, Schmitz ML. 2000. The pro- or anti-apoptotic function of NF-kappaB is determined by the nature of the apoptotic stimulus. *Eur J Biochem* 267:3828–35.
- Kaltschmidt B, Ndiaye D, Korte M, Pothion S, Arbibe L, Prüllage M, and others. 2006. NF-kappaB regulates spatial memory formation and synaptic plasticity through protein kinase A/CREB signaling. *Mol Cell Biol* 26:2936–46.
- Kaltschmidt C, Kaltschmidt B, Baeuerle PA. 1993. Brain synapses contain inducible forms of the transcription factor NF-kappa B. *Mech Dev* 43:135–47.
- Kato T Jr, Delhase M, Hoffmann A, Karin M. 2003. CK2 is a C-terminal IkappaB kinase responsible for NF-kappaB activation during the UV response. *Mol Cell* 12:829–39.
- Khoroshii R, Babcock AA, Owens T. 2008. NF-kappaB-driven STAT2 and CCL2 expression in astrocytes in response to brain injury. *J Immunol* 181:7284–91.
- Khoshnan A, Ko J, Tescu S, Brundin P, Patterson PH. 2009. IKKalpha and IKKbeta regulation of DNA damage-induced cleavage of huntingtin. *PLoS One* 4:e5768.
- Khoshnan A, Ko J, Watkin EE, Paige LA, Reinhart PH, Patterson PH. 2004. Activation of the IkappaB kinase complex and nuclear factor-kappaB contributes to mutant huntingtin neurotoxicity. *J Neurosci* 24:7999–8008.
- Kiernan R, Brès V, Ng RW, Coudart MP, El Messaoudi S, Sardet C, and others. 2003. Post-activation turn-off of NF-kappa B-dependent transcription is regulated by acetylation of p65. *J Biol Chem* 278:2758–66.
- Kim S, La Motte-Mohs RN, Rudolph D, Zuniga-Pflucker JC, Mak TW. 2003. The role of nuclear factor-kappaB essential modulator (NEMO) in B cell development and survival. *Proc Natl Acad Sci U S A* 100:1203–8.
- Kitaoka Y, Kumai T, Lam TT, Munemasa Y, Isenoumi K, Motoki M, and others. 2004. Nuclear factor-kappa B p65 in

- NMDA-induced retinal neurotoxicity. *Brain Res Mol Brain Res* 131:8–16.
- Köntgen F, Grumont RJ, Strasser A, Metcalf D, Li R, Tarlinton D, and others. 1995. Mice lacking the c-rel proto-oncogene exhibit defects in lymphocyte proliferation, humoral immunity, and interleukin-2 expression. *Genes Dev* 9:1965–77.
- Koo JW, Russo SJ, Ferguson D, Nestler EJ, Duman RS. 2010. Nuclear factor-kappaB is a critical mediator of stress-impaired neurogenesis and depressive behavior. *Proc Natl Acad Sci U S A* 107:2669–74.
- Kwak YT, Guo J, Shen J, Gaynor RB. 2000. Analysis of domains in the IKKalpha and IKKbeta proteins that regulate their kinase activity. *J Biol Chem* 275:14752–9.
- Lam LT, Davis RE, Ngo VN, Lenz G, Wright G, Xu W, and others. 2008. Compensatory IKKalpha activation of classical NF-kappaB signaling during IKKbeta inhibition identified by an RNA interference sensitization screen. *Proc Natl Acad Sci U S A* 105:20798–803.
- Lee HJ, Kim SH, Kim KW, Um JH, Lee HW, Chung BS, and others. 2001. Antiapoptotic role of NF-kappaB in the auto-oxidized dopamine-induced apoptosis of PC12 cells. *J Neurochem* 76:602–9.
- Li LY, Zhao XL, Fei XF, Gu ZL, Qin ZH, Liang ZQ. 2008. Bilobalide inhibits 6-OHDA-induced activation of NF-kappaB and loss of dopaminergic neurons in rat substantia nigra. *Acta Pharmacol Sin* 29:539–47.
- Li Q, Estepa G, Memet S, Israel A, Verma IM. 2000. Complete lack of NF-kappaB activity in IKK1 and IKK2 double-deficient mice: additional defect in neurulation. *Genes Dev* 14:1729–33.
- Li Q, Lu Q, Hwang JY, Buscher D, Lee KF, Izpisua-Belmonte JC, and others. 1999b. IKK1-deficient mice exhibit abnormal development of skin and skeleton. *Genes Dev* 13:1322–8.
- Li Q, Van Antwerp D, Mercurio F, Lee KF, Verma IM. 1999a. Severe liver degeneration in mice lacking the IkappaB kinase 2 gene. *Science* 284:321–5.
- Li ZW, Chu W, Hu Y, Delhase M, Deerinck T, Ellisman M, and others. 1999c. The IKKbeta subunit of IkappaB kinase (IKK) is essential for nuclear factor kappaB activation and prevention of apoptosis. *J Exp Med* 189:1839–45.
- Li ZW, Omori SA, Labuda T, Karin M, Rickert RC. 2003. IKK beta is required for peripheral B cell survival and proliferation. *J Immunol* 170:4630–7.
- Liang ZQ, Wang X, Li LY, Wang Y, Chen RW, Chuang DM, and others. 2007. Nuclear factor-kappaB-dependent cyclin D1 induction and DNA replication associated with N-methyl-D-aspartate receptor-mediated apoptosis in rat striatum. *J Neurosci Res* 85:1295–309.
- Lo JC, Basak S, James ES, Quiambo RS, Kinsella MC, Alegre ML, and others. 2006. Coordination between NF-kappaB family members p50 and p52 is essential for mediating LTbetaR signals in the development and organization of secondary lymphoid tissues. *Blood* 107:1048–55.
- Lu Z, Yu S, Wei J, Wei L. 2006. Age-related neural degeneration in nuclear-factor kappaB p50 knockout mice. *Neuroscience* 139:965–78.
- Maehara K, Hasegawa T, Isobe KI. 2000. A NF-kappaB p65 subunit is indispensable for activating manganese superoxide: dismutase gene transcription mediated by tumor necrosis factor-alpha. *J Cell Biochem* 77:474–86.
- Mattson MP, Camandola S. 2001. NF-kappaB in neuronal plasticity and neurodegenerative disorders. *J Clin Invest* 107:247–54.
- Meffert MK, Chang JM, Wiltgen BJ, Fanselow MS, Baltimore D. 2003. NF-kappa B functions in synaptic signaling and behavior. *Nat Neurosci* 6:1072–8.
- Meunier A, Latrémolière A, Dominguez E, Mauborgne A, Philippe S, Hamon M, and others. 2007. Lentiviral-mediated targeted NF-kappaB blockade in dorsal spinal cord glia attenuates sciatic nerve injury-induced neuropathic pain in the rat. *Mol Ther* 15:687–97.
- Middleton G, Hamanoue M, Enokido Y, Wyatt S, Pennica D, Jaffray E, and others. 2000. Cytokine-induced nuclear factor kappa B activation promotes the survival of developing neurons. *J Cell Biol* 148:325–32.
- Middleton G, Wyatt S, Ninkina N, Davies AM. 2001. Reciprocal developmental changes in the roles of Bcl-w and Bcl-x(L) in regulating sensory neuron survival. *Development* 128:447–57.
- Migheli A, Piva R, Atzori C, Troost D, Schiffer D. 1997. c-Jun, JNK/SAPK kinases and transcription factor NF-kappa B are selectively activated in astrocytes, but not motor neurons, in amyotrophic lateral sclerosis. *J Neuropathol Exp Neurol* 56:1314–22.
- Mincheva S, Garcera A, Gou-Fabregas M, Encinas M, Dolcet X, Soler RM. 2011. The canonical nuclear factor-kappaB pathway regulates cell survival in a developmental model of spinal cord motoneurons. *J Neurosci* 31:6493–503.
- Mir M, Tolosa L, Asensio VJ, Lladó J, Olmos G. 2008. Complementary roles of tumor necrosis factor alpha and interferon gamma in inducible microglial nitric oxide generation. *J Neuroimmunol* 204:101–9.
- Müller JR, Siebenlist U. 2003. Lymphotoxin beta receptor induces sequential activation of distinct NF-kappa B factors via separate signaling pathways. *J Biol Chem* 278:12006–12.
- Nakai M, Qin ZH, Chen JF, Wang Y, Chase TN. 2000. Kainic acid-induced apoptosis in rat striatum is associated with nuclear factor-kappaB activation. *J Neurochem* 74:647–58.
- Nakajima K, Kohsaka S. 1998. Functional roles of microglia in the central nervous system. *Hum Cell* 11:141–55.
- Nickols JC, Valentine W, Kanwal S, Carter BD. 2003. Activation of the transcription factor NF-kappaB in Schwann cells is required for peripheral myelin formation. *Nat Neurosci* 6:161–7.
- Niu YL, Zhang WJ, Wu P, Liu B, Sun GT, Yu DM, and others. 2010. Expression of the apoptosis-related proteins caspase-3

- and NF-kappaB in the hippocampus of Tg2576 mice. *Neurosci Bull* 26:37–46.
- Nurmi A, Lindsberg PJ, Koistinaho M, Zhang W, Juettler E, Karjalainen-Lindsberg ML, and others. 2004. Nuclear factor-kappaB contributes to infarction after permanent focal ischemia. *Stroke* 35:987–91.
- Olson CM, Hedrick MN, Izadi H, Bates TC, Olivera ER, Anguita J. 2007. p38 mitogen-activated protein kinase controls NF-kappaB transcriptional activation and tumor necrosis factor alpha production through RelA phosphorylation mediated by mitogen- and stress-activated protein kinase 1 in response to *Borrelia burgdorferi* antigens. *Infect Immun* 75:270–7.
- Panet H, Barzilai A, Daily D, Melamed E, Offen D. 2001. Activation of nuclear transcription factor kappa B (NF-kappaB) is essential for dopamine-induced apoptosis in PC12 cells. *J Neurochem* 77:391–8.
- Paris D, Patel N, Quadros A, Linan M, Bakshi P, Ait-Ghezala G, and others. 2007. Inhibition of Abeta production by NF-kappaB inhibitors. *Neurosci Lett* 415:11–6.
- Perkins ND. 2007. Integrating cell-signalling pathways with NF-kappaB and IKK function. *Nat Rev Mol Cell Biol* 8:49–62.
- Pizzi M, Goffi F, Boroni F, Benarese M, Perkins SE, Liou HC, and others. 2002. Opposing roles for NF-kappa B/Rel factors p65 and c-Rel in the modulation of neuron survival elicited by glutamate and interleukin-1beta. *J Biol Chem* 277:20717–23.
- Place RF, Haspeslagh D, Hubbard AK, Giardina C. 2001. Cytokine-induced stabilization of newly synthesized I(kappa) B-alpha. *Biochem Biophys Res Commun* 283:813–20.
- Pyo JS, Ko YS, Kim WH, Kim M, Lee KW, Nam SY, and others. 2010. Impairment of nuclear factor-kappaB activation increased glutamate excitotoxicity in a motoneuron-neuroblastoma hybrid cell line expressing mutant (G93A) Cu/Zn-superoxide dismutase. *J Neurosci Res* 88:2494–503.
- Qin ZH, Chen RW, Wang Y, Nakai M, Chuang DM, Chase TN. 1999. Nuclear factor kappaB nuclear translocation upregulates c-Myc and p53 expression during NMDA receptor-mediated apoptosis in rat striatum. *J Neurosci* 19:4023–33.
- Rojo AI, Salinas M, Martín D, Perona R, Cuadrado A. 2004. Regulation of Cu/Zn-superoxide dismutase expression via the phosphatidylinositol 3 kinase/Akt pathway and nuclear factor-kappaB. *J Neurosci* 24:7324–34.
- Rothgiesser KM, Erenner S, Waibel S, Lüscher B, Hottiger MO. 2010b. SIRT2 regulates NF-κB dependent gene expression through deacetylation of p65 Lys310. *J Cell Sci* 123:4251–8.
- Rothgiesser KM, Fey M, Hottiger MO. 2010a. Acetylation of p65 at lysine 314 is important for late NF-kappaB-dependent gene expression. *BMC Genomics* 11:22.
- Rudolph D, Yeh W, Wakeham A, Rudolph B, Nallainathan D, Potter J, and others. 2000. Severe liver degeneration and lack of NF-kappaB activation in NEMO/IKKgamma-deficient mice. *Genes Dev* 14:854–62.
- Ryu H, Smith K, Camelo SI, Carreras I, Lee J, Iglesias AH, and others. 2005. Sodium phenylbutyrate prolongs survival and regulates expression of anti-apoptotic genes in transgenic amyotrophic lateral sclerosis mice. *J Neurochem* 93:1087–98.
- Saha RN, Liu X, Pahan K. 2006. Up-regulation of BDNF in astrocytes by TNF-alpha: a case for the neuroprotective role of cytokine. *J Neuroimmune Pharmacol* 1:212–22.
- Sanchez-Ponce D, Muñoz A, Garrido JJ. 2011. Casein kinase 2 and microtubules control axon initial segment formation. *Mol Cell Neurosci* 46:222–34.
- Sarnico I, Lanzillotta A, Boroni F, Benarese M, Alghisi M, Schwaninger M, and others. 2009. NF-kappaB p50/RelA and c-Rel-containing dimers: opposite regulators of neuron vulnerability to ischaemia. *J Neurochem* 108:475–85.
- Schmidt-Ullrich R, Mémet S, Lilienbaum A, Feuillard J, Raphaël M, Israel A. 1996. NF-kappaB activity in transgenic mice: developmental regulation and tissue specificity. *Development* 122:2117–28.
- Schwabe RF, Brenner DA. 2002. Role of glycogen synthase kinase-3 in TNF-alpha-induced NF-kappaB activation and apoptosis in hepatocytes. *Am J Physiol Gastrointest Liver Physiol* 283:G204–G211.
- Schwaninger M, Inta I, Herrmann O. 2006. NF-kappaB signaling in cerebral ischaemia. *Biochem Soc Trans* 34:1291–4.
- Schwarz EM, Van Antwerp D, Verma IM. 1996. Constitutive phosphorylation of IkappaBalpha by casein kinase II occurs preferentially at serine 293: requirement for degradation of free IkappaBalpha. *Mol Cell Biol* 16:3554–9.
- Sen R, Baltimore D. 1986. Multiple nuclear factors interact with the immunoglobulin enhancer sequences. *Cell* 46:705–16.
- Sha WC, Liou HC, Tuomanen EI, Baltimore D. 1995. Targeted disruption of the p50 subunit of NF-kappa B leads to multifocal defects in immune responses. *Cell* 80:321–30.
- Shang YC, Chong ZZ, Hou J, Maiese K. 2010. Wnt1, FoxO3a, and NF-kappaB oversee microglial integrity and activation during oxidant stress. *Cell Signal* 22:1317–29.
- Shou Y, Li N, Li L, Borowitz JL, Isom GE. 2002. NF-kappaB-mediated up-regulation of Bcl-X(S) and Bax contributes to cytochrome c release in cyanide-induced apoptosis. *J Neurochem* 81:842–52.
- Simpson CS, Morris BJ. 2000. Regulation of neuronal cell adhesion molecule expression by NF-kappa B. *J Biol Chem* 275:16879–84.
- Sizemore N, Lerner N, Dombrowski N, Sakurai H, Stark GR. 2002. Distinct roles of the Ikappa B kinase alpha and beta subunits in liberating nuclear factor kappa B (NF-kappa B) from Ikappa B and in phosphorylating the p65 subunit of NF-kappa B. *J Biol Chem* 277:3863–69.
- Smith D, Tweed C, Fernyhough P, Glazner GW. 2009. Nuclear factor-kappaB activation in axons and Schwann cells in experimental sciatic nerve injury and its role in modulating axon regeneration: studies with etanercept. *J Neuropathol Exp Neurol* 68:691–700.

- Sole C, Dolcet X, Segura MF, Gutierrez H, Diaz-Meco MT, Gozzelino R, and others. 2004. The death receptor antagonist FAIM promotes neurite outgrowth by a mechanism that depends on ERK and NF-kappa B signaling. *J Cell Biol* 167:479–92.
- Solt LA, Madge LA, Orange JS, May MJ. 2007. Interleukin-1-induced NF-kappaB activation is NEMO-dependent but does not require IKKbeta. *J Biol Chem* 282:8724–33.
- Sparacio SM, Zhang Y, Vilcek J, Benveniste EN. 1992. Cytokine regulation of interleukin-6 gene expression in astrocytes involves activation of an NF-kappa B-like nuclear protein. *J Neuroimmunol* 39:231–42.
- Stephenson D, Yin T, Smalstig EB, Hsu MA, Panetta J, Little S, and others. 2000. Transcription factor nuclear factor-kappa B is activated in neurons after focal cerebral ischemia. *J Cereb Blood Flow Metab* 20:592–603.
- Takada Y, Mukhopadhyay A, Kundu GC, Mahabeleshwar GH, Singh S, Aggarwal BB. 2003. Hydrogen peroxide activates NF-kappa B through tyrosine phosphorylation of I kappa B alpha and serine phosphorylation of p65: evidence for the involvement of I kappa B alpha kinase and Syk protein-tyrosine kinase. *J Biol Chem* 278:24233–41.
- Takano H, Gusella JF. 2002. The predominantly HEAT-like motif structure of huntingtin and its association and coincident nuclear entry with dorsal, an NF-kB/Rel/dorsal family transcription factor. *BMC Neurosci* 3:15.
- Tam WF, Lee LH, Davis L, Sen R. 2000. Cytoplasmic sequestration of rel proteins by IkappaBalpha requires CRM1-dependent nuclear export. *Mol Cell Biol* 20:2269–84.
- Tamatani M, Che YH, Matsuzaki H, Ogawa S, Okado H, Miyake S, and others. 1999. Tumor necrosis factor induces Bcl-2 and Bcl-x expression through NFkappaB activation in primary hippocampal neurons. *J Biol Chem* 274:8531–8.
- Taylor DL, Jones F, Kubota ES, Pocock JM. 2005. Stimulation of microglial metabotropic glutamate receptor mGlu2 triggers tumor necrosis factor alpha-induced neurotoxicity in concert with microglial-derived Fas ligand. *J Neurosci* 25:2952–64.
- Tolosa L, Caraballo-Miralles V, Olmos G, Lladó J. 2011. TNF- α potentiates glutamate-induced spinal cord motoneuron death via NF- κ B. *Mol Cell Neurosci* 46:176–86.
- Valerio A, Boroni F, Benarese M, Sarnico I, Ghisi V, Bresciani LG, and others. 2006. NF-kappaB pathway: a target for preventing beta-amyloid (A β)-induced neuronal damage and A β 42 production. *Eur J Neurosci* 23:1711–20.
- Vermeulen L, De Wilde G, Van Damme P, Vanden Berghe W, Haegeman G. 2003. Transcriptional activation of the NF-kappaB p65 subunit by mitogen- and stress-activated protein kinase-1 (MSK1). *EMBO J* 22:1313–24.
- Vollgraf U, Wegner M, Richter-Landsberg C. 1999. Activation of AP-1 and nuclear factor-kappaB transcription factors is involved in hydrogen peroxide-induced apoptotic cell death of oligodendrocytes. *J Neurochem* 73:2501–9.
- Wang CY, Mayo MW, Korneluk RG, Goeddel DV, Baldwin AS. 1998. NF-kappaB antiapoptosis: induction of TRAF1 and TRAF2 and c-IAP1 and c-IAP2 to suppress caspase-8 activation. *Science* 281:1680–3.
- Wang D, Westerheide SD, Hanson JL, Baldwin AS Jr. 2000. Tumor necrosis factor alpha-induced phosphorylation of RelA/p65 on Ser529 is controlled by casein kinase II. *J Biol Chem* 275:32592–7.
- Wei X, Zhang X, Flick LM, Drissi H, Schwarz EM, O'Keefe RJ. 2009. Titanium particles stimulate COX-2 expression in synovial fibroblasts through an oxidative stress-induced, calpain-dependent, NF-kappaB pathway. *Am J Physiol Cell Physiol* 297:C310–320.
- Weih F, Carrasco D, Durham SK, Barton DS, Rizzo CA, Ryseck RP, and others. 1995. Multiorgan inflammation and hematopoietic abnormalities in mice with a targeted disruption of RelB, a member of the NF-kappa B/Rel family. *Cell* 80:331–40.
- Weih F, Durham SK, Barton DS, Sha WC, Baltimore D, Bravo R. 1997. p50-NF-kappaB complexes partially compensate for the absence of RelB: severely increased pathology in p50(–/–)relB(–/–) double-knockout mice. *J Exp Med* 185:1359–70.
- Woronicz JD, Gao X, Cao Z, Rothe M, Goeddel DV. 1997. IkappaB kinase-beta: NF-kappaB activation and complex formation with IkappaB kinase-alpha and NIK. *Science* 278:866–9.
- Wu H, Lozano G. 1994. NF-kappa B activation of p53. A potential mechanism for suppressing cell growth in response to stress. *J Biol Chem* 269:20067–74.
- Xie QW, Kashiwabara Y, Nathan C. 1994. Role of transcription factor NF-kappa B/Rel in induction of nitric oxide synthase. *J Biol Chem* 269:4705–8.
- Xu J, Chen S, Ahmed SH, Chen H, Ku G, Goldberg MP, and others. 2001. Amyloid-beta peptides are cytotoxic to oligodendrocytes. *J Neurosci* 21:RC118.
- Xu W, Chi L, Xu R, Ke Y, Luo C, Cai J, and others. 2005. Increased production of reactive oxygen species contributes to motor neuron death in a compression mouse model of spinal cord injury. *Spinal Cord* 43:204–13.
- Xu Z, Chen S, Li X, Luo G, Li L, Le W. 2006. Neuroprotective effects of (–)-epigallocatechin-3-gallate in a transgenic mouse model of amyotrophic lateral sclerosis. *Neurochem Res* 31:1263–9.
- Yamamoto Y, Verma UN, Prajapati S, Kwak YT, Gaynor RB. 2003. Histone H3 phosphorylation by IKK-alpha is critical for cytokine-induced gene expression. *Nature* 423:655–9.
- Yeo W, Gautier J. 2004. Early neural cell death: dying to become neurons. *Dev Biol* 274:233–44.
- Yilmaz ZB, Weih DS, Sivakumar V, Weih F. 2003. RelB is required for Peyer's patch development: differential regulation of p52-RelB by lymphotoxin and TNF. *EMBO J* 22:121–30.

- Yoon C, Korade Z, Carter BD. 2008. Protein kinase A-induced phosphorylation of the p65 subunit of nuclear factor-kappaB promotes Schwann cell differentiation into a myelinating phenotype. *J Neurosci* 28:3738–46.
- Yu Z, Zhou D, Bruce-Keller AJ, Kindy MS, Mattson MP. 1999. Lack of the p50 subunit of nuclear factor-kappaB increases the vulnerability of hippocampal neurons to excitotoxic injury. *J Neurosci* 19:8856–65.
- Yu Z, Zhou D, Cheng G, Mattson MP. 2000. Neuroprotective role for the p50 subunit of NF-kappaB in an experimental model of Huntington's disease. *J Mol Neurosci* 15:31–44.
- Zandi E, Rothwarf DM, Delhase M, Hayakawa M, Karin M. 1997. The IkappaB kinase complex (IKK) contains two kinase subunits, IKKalpha and IKKbeta, necessary for IkappaB phosphorylation and NF-kappaB activation. *Cell* 91:243–52.
- Zhang J, Warren MA, Shoemaker SF, Ip MM. 2007. NFkappaB1/p50 is not required for tumor necrosis factor-stimulated growth of primary mammary epithelial cells: implications for NFkappaB2/p52 and RelB. *Endocrinology* 148:268–78.
- Zhong H, Voll RE, Ghosh S. 1998. Phosphorylation of NF-kappa B p65 by PKA stimulates transcriptional activity by promoting a novel bivalent interaction with the coactivator CBP/p300. *Mol Cell* 1:661–71.