

## XM531 Problem Set 4

## Problem 1

The *gamma function* is denoted by  $\Gamma(p)$  and is defined by the integral

$$\Gamma(p+1) = \int_0^\infty e^{-x} x^p dx.$$

- Show that, for  $p > 0$ ,  $\Gamma(p+1) = p\Gamma(p)$ .
- Show that  $\Gamma(1) = 1$ .
- If  $p$  is a positive integer  $n$ , show that  $\Gamma(n+1) = n!$ . Since  $\Gamma(p)$  is also defined when  $p$  is not an integer, this function provides an extension of the factorial function to nonintegral values of the independent variable. Note that it is also consistent to define  $0! = 1$ .
- Show that, for  $p > 0$ ,

$$p(p+1)(p+2)\dots(p+n+1) = \frac{\Gamma(p+n)}{\Gamma(p)}.$$

- e. Thus  $\Gamma(p)$  can be determined for all positive values of  $p$  if  $\gamma(p)$  is known in a single interval of unit length, say  $0 < p \leq 1$ . It is possible to show that  $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$ . Find  $\Gamma\left(\frac{3}{2}\right)$  and  $\Gamma\left(\frac{11}{2}\right)$ .

[illegible]

## Problem 2

Use the Laplace transform to solve the given initial value problem.

$$y'' + 3y' + 2y = 0; \quad y(0) = 1, \quad y'(0) = 0.$$

### Problem 3

Let  $\alpha$  be a real number. Express the solution of the given initial value problem in terms of a convolution integral.

$$y'' + 2y' + 2y = \sin(\alpha t); \quad y(0) = 0, \quad y'(0) = 0.$$

This image shows a full page of white paper with horizontal dotted lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

## Problem 4

Let  $\alpha$  be a real number. Find the Laplace transform of the following:

a.  $f(t) = t^2 \sinh(\alpha t)$

$$\text{b. } g(t) = \begin{cases} t, & 0 \leq t < 1, \\ 2 - t, & 1 \leq t < 2, \text{ and} \\ 0, & 2 \leq t < \infty \end{cases}$$

[illegible]

