

Figure 6. Phenotype paper table (15 rows = 3 cohorts x 5 phenotype clusters)
spatial_k2: focal vs diffuse; burden_k3: low/mid/high Agatston tertiles

Cohort	Partition	Cluster	N	Agatston med	[IQR]	%Qr36d/2	%low burden	Hennig
full	spatial_k2	focal	195	31	[11-65]	48%	60%	0.86
full	spatial_k2	diffuse	225	425	[185-925]	56%	1%	0.89
full	burden_k3	low	140	16	[9-34]	46%	82%	
full	burden_k3	mid	140	125	[80-202]	52%	3%	
full	burden_k3	high	140	766	[484-1397]	59%	0%	
Qr36d/2	spatial_k2	focal	111	49	[15-94]	100%	49%	0.92
Qr36d/2	spatial_k2	diffuse	109	675	[269-1215]	100%	0%	0.93
Qr36d/2	burden_k3	low	74	25	[9-46]	100%	73%	
Qr36d/2	burden_k3	mid	73	166	[115-256]	100%	0%	
Qr36d/2	burden_k3	high	73	921	[675-1624]	100%	0%	
I30f/3	spatial_k2	focal	93	26	[10-57]	0%	67%	0.89
I30f/3	spatial_k2	diffuse	107	319	[135-696]	0%	3%	0.90
I30f/3	burden_k3	low	67	14	[7-26]	0%	91%	
I30f/3	burden_k3	mid	66	92	[63-137]	0%	6%	
I30f/3	burden_k3	high	67	638	[351-1191]	0%	0%	