

Other laboratory strains (a, d, e) or non-ITP control groups (b, c)
The four UM-HET3 founder strains
ITP controls
a ITP males were heavier than every founder strain and almost every laboratory strain
ITP lightest quartile 34.4 g
ITP mean 40.6 g
ITP heaviest quartile 46.9 g
C3H/HeJ
C57BL/6J
DBA/2J
BALB/cByJ
1 of 27 strains reaches the ITP mean
20
25
30
35
40
45
Male body weight (g)
b Male ITP controls lived as long as other control groups
ITP median 801 days
DBA/2J
C3H/HeJ
BALB/cByJ
C57BL/6J
10 of 20 non-ITP groups reach the ITP median
400
500
600
700
800
900
1000
Male control-group lifespan (days)
c Female ITP controls outlived every non-ITP control group
ITP median 891 days
C3H/HeJ
BALB/cByJ
DBA/2J
C57BL/6J
0 of 17 non-ITP groups reach the ITP median
400
500
600
700
800
900
1000
Female control-group lifespan (days)
d ITP males carried more fat than 38 of 40 strains
ITP 35.8%
C57BL/6J
C3H/HeJ
BALB/cByJ
DBA/2J
2 of 40 strains reach the ITP value
15
20
25
30
35
40
45
Male body fat (%)
e ITP females carried more fat than 39 of 40 strains
ITP 43.4%
BALB/cByJ
DBA/2J
C3H/HeJ
C57BL/6J
1 of 40 strains reaches the ITP value
15
20
25
30
35
40
45
Female body fat (%)