

Compound (experiments)
All 14 compounds (26)
+5.0%
+20.2%
Shared treatment × weight HR 0.90 (0.85–0.95); P = 1.1 × 10⁻⁴
Lifespan gain in the lightest mice (≥5%)
16-hydroxyestradiol (1)
estradiol metabolite
+17.7%
+25.7%
Acarbose (4)
α-glucosidase inhibitor
+7.9%
+24.8%
Canagliflozin (2)
SGLT2 inhibitor
+6.1%
+23.2%
17α-estradiol (4)
non-feminizing estrogen
+5.9%
+22.9%
Rapamycin (6)
mTOR inhibitor
+7.3%
+21.0%
MIF098 (1)
MIF antagonist
+6.5%
+14.7%
Glycine (1)
amino acid
+1.5%
+8.3%
Little or no lifespan gain in the lightest mice
Meclizine (1)
antihistamine, mTORC1 inhibitor
+3.8%
+29.1%
Candesartan (1)
angiotensin receptor blocker
+2.2%
+24.1%
NDGA (1)
antioxidant
+3.2%
+19.1%
Geranylgeranylacetone (1)
heat-shock protein inducer
−2.8%
+19.9%
Resveratrol (1)
plant polyphenol
−5.0%
+11.5%
HBX (1)
amyloid-binding compound
−2.6%
+7.9%
Fisetin (1)
senolytic flavonoid
−7.4%
+6.8%
Lightest quartile
Heaviest quartile
−10
0
10
20
30
Increase in median lifespan versus controls (%)