Laboratory strains (a, d, e) or non-ITP control groups (b, c)
UM-HET3 founder strains
ITP reference values

a Body weight, ♂

Survey mean 30.3 g
ITP lightest quartile 34.4
ITP overall mean 40.6
ITP heaviest quartile 46.9
DBA/2J
C57BL/6J
BALB/cByJ
C3H/HeJ
1 of 27 strains at or above the ITP value; 7 reach the lightest ITP quartile, none the heaviest.
20
25
30
35
40
45
Body weight (g)

b Control survival, ♂

ITP median 801 days
BALB/cByJ
C57BL/6J
DBA/2J
C3H/HeJ
11 of 24 non-ITP groups reach the ITP median
400
600
800
1000
Control-group lifespan (days)

c Control survival, ♀

ITP median 891 days
DBA/2J
BALB/cByJ
C3H/HeJ
C57BL/6J
0 of 20 non-ITP groups reach the ITP median
400
600
800
1000
Control-group lifespan (days)

d Body fat, ♂

ITP UM-HET3 35.8
C57BL/6J
DBA/2J
C3H/HeJ
BALB/cByJ
2 of 40 strains at or above the ITP value
20
25
30
35
40
Body fat (%)

e Body fat, ♀

ITP UM-HET3 43.4
C57BL/6J
DBA/2J
C3H/HeJ
BALB/cByJ
1 of 40 strains at or above the ITP value
20
25
30
35
40
Body fat (%)