
IDENTIFYING FINANCIAL RISK FACTORS WITH A LOW-RANK/SPARSE DECOMPOSITION

Alexander D. Shkolnik

UC Berkeley

ads2@berkeley.edu

Risk Seminar. April 12th, 2016.

Joint work with Lisa Goldberg, UC Berkeley.

Factor models in finance

- Market model and CAPM
 - market equilibrium theory of asset prices under conditions of risk,
 - Sharpe (1964) and Treynor (1962).
 - Arbitrage pricing theory and statistical models
 - the market portfolio plays no special role,
 - e.g. GNP and other factors (Ross 1976).
 - Fundamental models
 - reverse the roles of the known and unknown variables to estimate factor returns; Rosenberg (1984) and Rosenberg (1984).
-

The factor model

- Returns to securities are driven by a relatively small number of risk factors, plus security specific returns: R is a $T \times N$ matrix

$$R = \psi Y + \epsilon \tag{1}$$

- ψ is a $T \times K$ matrix of factor returns.
 - Y is a $K \times N$ matrix of factor exposures.
 - ϵ is a $T \times N$ matrix of security specific returns.
- N is the number of securities, T is the number of observations and K is (a relatively small) number of factors,
 - To identify factors we rely on a universe of securities whose returns depend linearly on returns to factors.
-

Security covariance matrix

- In R the factor and specific returns ψ and ϵ are random while Y (the exposures) are to be estimated.

$$R = \psi Y + \epsilon \quad (2)$$

- Take ψ and ϵ to be mean zero and uncorrelated. Then the factor covariance matrix takes the form

$$\Sigma = Y^{\top} F Y + S \quad (3)$$

- F is a factor covariance (diagonal if factors are uncorrelated),
 - S is the specific risk covariance matrix (diagonal if the specific returns are uncorrelated).
-

A taxonomy of risk factors

- **Broad:** market, equity styles, interest rates, credit
 - **Thin:** industries, countries, currencies, credit
 - Persistent: all of the above, in some cases
 - Emerging or transient: new industries (internet), new sensitivities (housing bubble, climate), equity styles, liquidity, credit
-

Thin and broad risk factors

- If the factor and specific returns are uncorrelated:

$$\Sigma = L + S \quad (4)$$

- $L = Y^{\top} F Y$ is a rank K matrix,
 - S is the specific risk covariance matrix.
 - The K **broad factors** are contained in L ,
 - i.e. factors that affect most/all of the securities.
 - The **thin factors** may be found in S ,
 - i.e. factors that affect a “thin” (smaller) number of securities.
 - (S diagonal means there are no thin factors)
-

Extracting risk factors

- Determine the matrices $L = Y^\top F Y$ and S in the decomposition

$$\Sigma = L + S. \quad (5)$$

- An elementary example (identifiability)
 - Suppose we observe R where $R = \psi + \epsilon$ and it is known that $\psi \sim N(0, \sigma_\psi)$ and $\epsilon \sim N(0, \sigma_\epsilon)$.
 - Is it possible to recover σ_ψ^2 and σ_ϵ^2 if we know only their sum?
 - What conditions are required for L and S to be **identifiable**?
 - There is **estimation error** (given sample covariance $\hat{\Sigma}$ only).
-

A number of approaches

- Fundamental models; requires many human analysts; factors are included as long as model designers think of them.
 - Algebraic approach (Drton, Sturmfels & Sullivant 2007); has yet to yield scalable algorithms or address practical issues.
 - **ML factor analysis** (statistical model, latent factors):
 - Distribution assumptions address estimation errors.
 - Identifiability depends on initialization/constraints.
 - **Dimensionality reduction** (PCA, distribution free):
 - Identifiability is inherently not an issue.
 - Estimation errors treated by regularization (asymptotic analysis).
 - **Convex optimization:**
 - Low rank plus sparse decompositions
-

ML factor analysis

- We assume a distribution, e.g. let (R, ψ) be a Gaussian random vector in \mathbb{R}^{N+K} with covariance matrix

$$\begin{bmatrix} \Sigma & Y^\top \\ Y & F \end{bmatrix} \quad (6)$$

(R are observed variables; ψ are hidden (latent) variables).

- Σ may be decomposed as (Anderson 1968):

$$\Sigma = Y^\top F Y + S \quad (S = \Sigma_{R|\psi}). \quad (7)$$

- Here, S is the conditional covariance (given the latent variables).
 - The fact that it is constant is a feature of the Gaussian distribution.
 - Plays the role of the specific covariance.
-

Some controversy

- Given the sample covariance $\hat{\Sigma}$, we maximize the likelihood

$$G(Y, F, S) = \log \det (Y^{\top} F Y + S)^{-1} - \text{tr} \left((Y^{\top} F Y + S)^{-1} \hat{\Sigma} \right) .$$

Jöreskog (1967, grad), Rubin & Thayer (1982, EM algorithm), etc.

- Rubin & Thayer (1982) claim their EM algorithm found “multiple local maxima of the likelihood”.
- Bentler & Tanaka (1983) point out “problems with EM algorithms for factor analysis”:

“Rather than highlight the limitations of ML factor analysis, the example demonstrates weaknesses in the EM algorithm ...”

- Rubin & Thayer (1983) respond that the EM algorithm can find local maxima, is complementary to other methods, etc ...
-

Decomposition multiplicity

- Letting $Q = (Y^\top FY + S)^{-1}$ we may write $G(S, Y, F) = G(Q)$,

$$G(Q) = \log \det(Q) - \text{tr}(Q\hat{\Sigma}).$$

- First order conditions (Petersen & Pedersen 2008) may be stated to prove the unique solution is $Q = \hat{\Sigma}^{-1}$.
- The identifiability issues come in the multiplicity of decompositions

$$Q^{-1} = \hat{\Sigma} = L + S. \tag{8}$$

- Multiplicity also occurs in $L = Y^\top FY$, e.g. when F is diagonal $Y^\top FY$ is unique upto multiplication by an orthogonal matrix.
- Multiplicity of (8) persists even when $S = \Delta$, a diagonal matrix (setting of Rubin & Thayer (1982)).

Identifiability

- “How much can we reduce the rank of a symmetric (positive definite) matrix by changing only its diagonal entries?” (Shapiro 1982)

$$\hat{\Sigma} = L + \Delta. \quad (9)$$

- No solution in the 1-dim case; a unique solution of rank 1 in the 2-dim case (4 equations and 4 unknowns).
 - Wilson & Worcester (1939) give example of a 6×6 correlation matrix and reduce it to rank 3 with two unique diagonals.
 - Basic analysis was carried out in Shapiro (1982), Shapiro (1985) and Shapiro & Ten Berge (2002).
 - Recently: Saunderson, Chandrasekaran, Parrilo & Willsky (2012).
- Even harder for $\hat{\Sigma} = L + S$ with non-zero off-diagonal entries in S .
-

ML factor analysis (summary)

- Under some assumed statistical model (typically Gaussian) we maximize a likelihood function.
 - In the Gaussian case the solution is the inverse of the sample covariance (the concentration or precision matrix).
 - **Identification** issues arise when a certain structure (e.g. low rank plus diagonal) on the covariance matrix is assumed.
 - Hence, convergence may occur at multiple points.
 - More constraints are required to resolve the indeterminacy.
 - Implies assumptions on broad/thin factors.
 - **Estimation error** issues are dealt with by assuming a statistical model (i.e. maximize the likelihood given observations).
-

Dimensionality reduction

- Seeks a low dimensional approximation of a given $\hat{\Sigma}$.
- Classical principal component analysis (PCA)

$$\text{minimize } \|\hat{\Sigma} - L\| \quad (10)$$

$$\text{subject to } \text{rank}(L) \leq K \quad (11)$$

- Solution given by truncated SVD

$$\hat{\Sigma} = U \Lambda U^{\top}; \quad L = \sum_{i=1}^K \lambda_i u_i u_i^{\top}. \quad (12)$$

extract components (factors) with largest variance.


- Many variants (LS, PFA, etc) (Jolliffe 2002).

Identifying factors (PCA)

- The output of PCA (truncated SVD) L retains the (broad) factors.
 - The remainder may be diagonalized, i.e. $S = \text{diag}(\hat{\Sigma} - L)$.
 - Strict factor model (Ross 1976).
 - Justified (Gershgorin theorem) when off-diag entries are small.
 - The remainder may be kept $S = \hat{\Sigma} - L$
 - Approximate factor model (Chamberlain & Rothschild 1982).
 - No issues with **identification**:
 - The “largest” variance is always explained by the broad factors.
 - The thin factors (if any) always have smaller variance.
-

Estimation errors (PCA)

- PCA is very sensitive to outliers in $\hat{\Sigma}$.



Remedies for PCA sensitivity

- Robust PCA (not poly-time with performance guarantees)
 - multivariate trimming (Gnanadesikan & Kettenring 1972),
 - random sampling (Fischler & Bolles 1981),
 - influence function techniques (De La Torre & Black 2003)
 - alternating minimization (Ke & Kanade 2005),
- Covariance regularization (Bickel & Levina 2008):
 - approximate, thresholded covariance matrix for asymptotics:

$$\log(N)/T = o(1) .$$

- Random matrix theory (spectra) (Karoui 2008): $T \sim N$.
-

Traditional methods (summary)

- **ML factor analysis:**
 - address estimation error by requiring a statistical model,
 - do not deal well with identifiability (thin vs broad factors).
 - **Dimentionality reduction (PCA):**
 - reduce identifiability to the claim that latent (broad) factors have the largest variance (picture for thin factors is unclear),
 - do not deal well with estimation error.
 - It is possible to address both issues? (with or without distributional assumptions and imposing as little structure as possible.)
-

Low rank plus sparse decomposition

- Suppose the true covariance matrix satisfies

$$\Sigma = L + S \quad (13)$$

- L is low rank (contains broad factors);
 - S is sparse (correlation due to thin factors).
- Given a sample covariance $\hat{\Sigma}$ we wish to recover estimates of (L, S) .
 - recover (L, S) given the true covariance Σ ?
 - approximate (L, S) given the sample covariance $\hat{\Sigma}$?
-

Is the problem well posed?

- Surprisingly, the answer is **yes**; but some assumptions are required on structure of the matrices L and S .
- **Identifiability** (eigenvectors of L and sparsity pattern of S)
 - deterministic conditions (Chandrasekaran, Sanghavi, Parrilo & Willsky 2011)
 - assumption on randomness (Candès, Li, Ma & Wright 2011)
- Both solve Principal Component Pursuit (PCP)

$$\text{minimize } \|L\|_* + \gamma \|S\|_1 \quad (14)$$

$$\text{subject to } \Sigma = L + S \quad (15)$$

- Ironically, γ is data-dependent in Chandrasekaran et al. (2011) and a universal constant ($1/\sqrt{N}$) in Candès et al. (2011).
-

Convex optimization

- Goes back to Shapiro (1982): “*how much can we reduce the rank of a symmetric matrix by changing only its diagonal entries*”.

$$\begin{aligned} & \text{minimize} \quad \text{rank}(L) \\ & \text{subject to} \quad \Sigma = L + \Delta, \\ & \quad \quad \quad L \geq 0, \Delta \text{ diagonal.} \end{aligned} \tag{16}$$

- Problem (16) is not convex; Shapiro (1982) considered the relaxation:

$$\begin{aligned} & \text{minimize} \quad \text{tr}(L) \\ & \text{subject to} \quad \Sigma = L + \Delta, \\ & \quad \quad \quad L \geq 0, \Delta \text{ diagonal.} \end{aligned} \tag{17}$$

- Problem (17), minimum trace factor analysis (MTFA), is convex.
-

MTFA

- If MTFA is feasible, (L, Δ) it has a unique optimal solution.

$$\begin{aligned} & \text{minimize} \quad \text{tr}(L) \\ & \text{subject to} \quad \Sigma = L + \Delta, \\ & \quad \quad \quad L \geq 0, \Delta \text{ (block) diagonal.} \end{aligned} \tag{18}$$

- Saunderson et al. (2012) study conditions under which the solution (L, Δ) is “correct” (they extend Δ to a block diagonal structure).
- **Factor extraction:** L contains the broad factors; blocks of Δ constitute thin factors (industries, countries, etc)
- If $\hat{\Sigma}$ is the input, we may not wish to preserve the equality constraint.

Principal component pursuit (PCP)

- Solves the convex optimization problem

$$\begin{aligned} & \text{minimize} \quad \|L\|_* + \gamma \|S\|_1 \\ & \text{subject to} \quad \Sigma = L + S \end{aligned} \tag{19}$$

- $\|L\|_*$ is the nuclear norm, the sum of singular values (equals the trace whenever L is symmetric):
 - attempts to reduce rank of L .
 - $\|S\|_1$ is the ℓ_1 -vector norm with S viewed as a vector.
 - encourages sparsity of S .
 - Its simple to preserve symmetry; less so for positive definiteness.
-

LRPS recovery

- Suppose $\Sigma = L + S$ is $N \times N$ for (L, S) such that
 - support of S is uniformly distributed over sets of cardinality m .
 - L obeys *incoherence conditions* with parameter μ (Candès & Recht 2009); i.e. singular vectors of L are reasonably spread out.
 - For some constants ρ_s and ρ_r we have

$$\text{rank}(L) \leq \frac{\rho_r N}{\mu (\log N)^2} \quad \text{and} \quad m \leq \rho_s N^2. \quad (20)$$

- Then^a (Candès et al. 2011) for $\gamma = 1/\sqrt{N}$ there exists constant c such that solving PCP recovers (L, S) exactly with probability

$$1 - cN^{-10}. \quad (21)$$

^asee (Chandrasekaran et al. 2011) for a related result.

Violating the equality constraint

- Suppose $\hat{\Sigma}$ is the sample covariance of the observed Gaussian returns.
- Since $\hat{\Sigma}$ may not be close to Σ we may wish to violate the constraint

$$\hat{\Sigma} = L + S \tag{22}$$

and instead maximize the Gaussian likelihood

$$G(L, S) = \log \det (L + S)^{-1} - \text{tr} \left((L + S)^{-1} \hat{\Sigma} \right).$$

LEMMA. *Given the decomposition $\Sigma = L + S$ we have a decomposition $\Sigma^{-1} = S^{-1} - \mathcal{L}$ such that $\text{rank}(L) = \text{rank}(\mathcal{L})$ with*

$$S = S^{-1} \quad \text{and} \quad \mathcal{L} = S^{-1} L \Sigma. \tag{23}$$

Latent variable convex optimization

- Let (R, ψ) be Gaussian with R the observed security returns and ψ the (unobserved) latent factors.
- If we believe Σ has a LRPS decomposition $L + S$ and $S = S^{-1}$ is **sparse** we may solve the convex optimization problem

$$\begin{aligned} &\text{minimize} && -G(\mathcal{L}, S) + \lambda (\|\mathcal{L}\|_* + \gamma \|S\|_1) \\ &\text{subject to} && S - \mathcal{L} \succ 0, \mathcal{L} \succeq 0 \end{aligned} \quad (24)$$

- For example, if S is block diagonal (or any permutation) so it S .
 - Problem (24) was proposed in Chandrasekaran, Parrilo & Willsky (2010) and analyzed in Chandrasekaran, Parrilo & Willsky (2012).
-

Parameters

- The problem parameters (λ, γ) appearing in

$$\begin{aligned} & \text{minimize} \quad -G(\mathcal{L}, \mathcal{S}) + \lambda (\|\mathcal{L}\|_* + \gamma \|\mathcal{S}\|_1) \\ & \text{subject to} \quad \mathcal{S} - \mathcal{L} \succ 0, \mathcal{L} \succeq 0 \end{aligned} \quad (25)$$

are data dependent and harder to select than for PCP.

- λ may be set in proportion to $\sqrt{N/T}$ and γ must be chosen by trial and error (Chandrasekaran et al. 2012).
- The solution $(\mathcal{L}, \mathcal{S})$ depends on λ to the degree that


$$\hat{\Sigma}^{-1} \approx \mathcal{S} - \mathcal{L}. \quad (26)$$

- We solve (25) by using an algorithm of Ma, Xue & Zou (2013).


Numerical studies (synthetic data)

- $N = 32$ securities
 - $T = 260$ observations (one year of daily data)
 - $K = 2$ broad factors:
 - The market, with annualized volatility %20
(long only: all securities have positive exposure).
 - Creditworthiness, with annualized volatility of %10
(long/short factor: half creditworthy; half close to default)
 - $\kappa = 4$ thin factors (e.g. countries)
-


Algorithm input: sample covariance matrix




Low-rank part (true vs recovered)



Sparse part (true vs recovered)




Eigenvalues (true vs recovered)




Numerical studies (real data)

- $N = 32$ securities
 - $T = 260$ observations (one year of daily data)
 - $K = ?$ broad factors
 - Securities drawn from $\kappa = 4$ countries
 - China
 - Argentina
 - India
 - Saudi Arabia
-


Algorithm input: sample covariance, Oct. 2015




LRPS decomposition (covariance)




Sample correlation matrix, Oct. 2015




LRPS decomposition (correlations)



Recovered eigenvalues



PCA solution (K=4) and remainder



Questions.
(ads2@berkeley.edu)

References

- Anderson, TW (Theodore Wilbur) (1968), *An introduction to multivariate statistical analysis*, John Wiley & Sons.
- Bentler, Peter M & Jeffrey S Tanaka (1983), ‘Problems with em algorithms for ml factor analysis’, *Psychometrika* **48**(2), 247–251.
- Bickel, Peter J & Elizaveta Levina (2008), ‘Covariance regularization by thresholding’, *The Annals of Statistics* pp. 2577–2604.
- Candès, Emmanuel J & Benjamin Recht (2009), ‘Exact matrix completion via convex optimization’, *Foundations of Computational mathematics* **9**(6), 717–772.
- Candès, Emmanuel J, Xiaodong Li, Yi Ma & John Wright (2011), ‘Robust principal component analysis?’, *Journal of the ACM (JACM)* **58**(3), 11.
- Chamberlain, Gary & Michael Rothschild (1982), ‘Arbitrage, factor structure, and mean-variance analysis on large asset markets’.
-

- Chandrasekaran, Venkat, Pablo A Parrilo & Alan S Willsky (2010), Latent variable graphical model selection via convex optimization, *in* ‘Communication, Control, and Computing (Allerton), 2010 48th Annual Allerton Conference on’, IEEE, pp. 1610–1613.
- Chandrasekaran, Venkat, Pablo A Parrilo & Alan S Willsky (2012), ‘Latent variable graphical model selection via convex optimization’, *The Annals of Statistics* **40**(4), 1935–1967.
- Chandrasekaran, Venkat, Sujay Sanghavi, Pablo A Parrilo & Alan S Willsky (2011), ‘Rank-sparsity incoherence for matrix decomposition’, *SIAM Journal on Optimization* **21**(2), 572–596.
- De La Torre, Fernando & Michael J Black (2003), ‘A framework for robust subspace learning’, *International Journal of Computer Vision* **54**(1-3), 117–142.
- Drton, Mathias, Bernd Sturmfels & Seth Sullivant (2007), ‘Algebraic factor analysis: tetrads, pentads and beyond’, *Probability Theory and Related Fields* **138**(3-4), 463–493.
-

- Fischler, Martin A & Robert C Bolles (1981), ‘Random sample consensus: a paradigm for model fitting with applications to image analysis and automated cartography’, *Communications of the ACM* **24**(6), 381–395.
- Gnanadesikan, Ramanathan & John R Kettenring (1972), ‘Robust estimates, residuals, and outlier detection with multiresponse data’, *Biometrics* pp. 81–124.
- Jolliffe, Ian (2002), *Principal component analysis*, Wiley Online Library.
- Jöreskog, Karl G (1967), ‘A general approach to confirmatory maximum likelihood factor analysis’, *ETS Research Bulletin Series* **1967**(2), 183–202.
- Karoui, Nouredine El (2008), ‘Spectrum estimation for large dimensional covariance matrices using random matrix theory’, *The Annals of Statistics* pp. 2757–2790.
- Ke, Qifa & Takeo Kanade (2005), Robust l_1 norm factorization in the presence of outliers and missing data by alternative convex programming, in ‘Computer Vision and Pattern Recognition, 2005. CVPR 2005. IEEE Computer Society Conference on’, Vol. 1, IEEE, pp. 739–746.
-

- Ma, Shiqian, Lingzhou Xue & Hui Zou (2013), ‘Alternating direction methods for latent variable gaussian graphical model selection’, *Neural computation* **25**(8), 2172–2198.
- Petersen, Kaare Brandt & Michael Syskind Pedersen (2008), ‘The matrix cookbook’, *Technical University of Denmark* **7**, 15.
- Rosenberg, Barr (1984), ‘Prediction of common stock investment risk’, *The Journal of Portfolio Management* **11**(1), 44–53.
- Ross, Stephen A (1976), ‘The arbitrage theory of capital asset pricing’, *Journal of economic theory* **13**(3), 341–360.
- Rubin, Donald B & Dorothy T Thayer (1982), ‘Em algorithms for ml factor analysis’, *Psychometrika* **47**(1), 69–76.
- Rubin, Donald B & Dorothy T Thayer (1983), ‘More on em for ml factor analysis’, *Psychometrika* **48**(2), 253–257.
- Saunderson, James, Venkat Chandrasekaran, Pablo A Parrilo & Alan S Willsky (2012), ‘Diagonal and low-rank matrix decompositions, correlation
-

matrices, and ellipsoid fitting', *SIAM Journal on Matrix Analysis and Applications* **33**(4), 1395–1416.

Shapiro, Alexander (1982), 'Rank-reducibility of a symmetric matrix and sampling theory of minimum trace factor analysis', *Psychometrika* **47**(2), 187–199.

Shapiro, Alexander (1985), 'Identifiability of factor analysis: Some results and open problems', *Linear Algebra and its Applications* **70**, 1–7.

Shapiro, Alexander & Jos MF Ten Berge (2002), 'Statistical inference of minimum rank factor analysis', *Psychometrika* **67**(1), 79–94.

Sharpe, William F (1964), 'Capital asset prices: A theory of market equilibrium under conditions of risk', *The journal of finance* **19**(3), 425–442.

Treynor, Jack L (1962), Toward a theory of market value of risky assets. Presented to the MIT Finance Faculty Seminar.

Wilson, Edwin B & Jane Worcester (1939), 'The resolution of six tests into three general factors', *Proceedings of the National Academy of Sciences of the United States of America* **25**(2), 73.
