

**Q1.**

$$\begin{aligned} \text{(a)} \quad \frac{d}{dx} \int_{4x}^{2x} \cos(e^{2t}) dt &= \frac{d}{dx} \int_{4x}^0 \cos(e^{2t}) dt + \frac{d}{dx} \int_0^{2x} \cos(e^{2t}) dt \\ &= -\frac{d}{du} \int_0^u \cos(e^{2t}) dt \cdot \frac{du}{dx} + \frac{d}{dw} \int_0^w \cos(e^{2t}) dt \cdot \frac{dw}{dx} \quad (\text{where } u = 4x, w = 2x) \\ &= -\cos(e^{2u})(-4) + \cos(e^{2w})(2) = 2\cos(e^{4x}) - 4\cos(e^{8x}). \end{aligned}$$

**[8 marks]**

(b) Let  $\Delta x = \frac{1}{n}$ ,  $x_i = \frac{i}{n}$ , and  $f(x) = \sqrt{x + 3x^2}$ . Then  $a = 0$ ,  $b = 1$ , and

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{\frac{ni+3i^2}{n^4}} = \lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{\frac{i}{n} + 3\left(\frac{i}{n}\right)^2} \cdot \frac{1}{n} = \int_0^1 \sqrt{x + 3x^2} dx$$

**[8 marks]**

**Q2.**

(a) Solving for the intersection points of the curve  $y = x^2$  and  $y = 4x - x^2$ , the area  $R$  is bounded above by  $y = 4x - x^2$ , bounded below by  $y = x^2$ , for  $0 \leq x \leq 2$ . We can use the cylindrical shell method. Since the axis is vertical ( $x = 4$ ), a typical vertical shell at  $x$  has height  $(4x - x^2) - x^2 = 4x - 2x^2$ . The radius of the shell is  $4 - x$ .

Hence, the volume  $V$  of the solid generated by revolving  $R$  about  $x = 4$  is

$$\begin{aligned} V &= 2\pi \int_0^2 (4-x)(4x-2x^2) dx = 2\pi \int_0^2 16x - 12x^2 + 2x^3 dx \\ &= 2\pi \left[ 8x^2 - 4x^3 + \frac{x^4}{2} \right]_0^2 = 2\pi [32 - 32 + 8] = 16\pi. \end{aligned}$$

**[10 marks]**

(b) Let  $u = \sqrt{k}x$ . Then  $du = \sqrt{k} dx$ .

$$\begin{aligned} \int_{-\infty}^{\infty} e^{-kx^2} dx &= \lim_{t \rightarrow -\infty} \int_t^0 e^{-kx^2} dx + \lim_{t \rightarrow \infty} \int_0^t e^{-kx^2} dx \\ &= \lim_{t \rightarrow -\infty} \frac{1}{\sqrt{k}} \int_t^0 e^{-u^2} du + \lim_{t \rightarrow \infty} \frac{1}{\sqrt{k}} \int_0^t e^{-u^2} du \\ &= \frac{1}{\sqrt{k}} \left( \lim_{t \rightarrow -\infty} \int_t^0 e^{-u^2} du + \lim_{t \rightarrow \infty} \int_0^t e^{-u^2} du \right) \\ &= \frac{1}{\sqrt{k}} \int_{-\infty}^{\infty} e^{-u^2} du = \frac{1}{\sqrt{k}} \sqrt{\pi}. \end{aligned}$$

**[6 marks]**

**Q3.**

(a) Let  $u = \cos 3x$ . Then  $du = -3 \sin 3x \, dx$ . Then

$$\begin{aligned} \int \sin^5(3x) \cos^4(3x) \, dx &= \int (\sin^2(3x))^2 \cos^4(3x) (\sin 3x) \, dx \\ &= \int (1 - u^2)^2 u^4 \left(-\frac{1}{3}\right) du = -\frac{1}{3} \int (1 - 2u^2 + u^4) u^4 \, du = -\frac{1}{3} \int u^4 - 2u^6 + u^8 \, du \\ &= -\frac{1}{3} \left( \frac{u^5}{5} - \frac{2u^7}{7} + \frac{u^9}{9} \right) + C = -\frac{\cos^5(3x)}{15} + \frac{2\cos^7(3x)}{21} - \frac{\cos^9(3x)}{27} + C. \end{aligned}$$

**[6 marks]**

(b) Let  $u = (\ln x)^2$ ,  $v' = x^{-2}$ . Then  $u' = 2(\ln x)x^{-1}$ ,  $v = -x^{-1}$ . By integration-by-parts, we have

$$\int (\ln x)^2 x^{-2} \, dx = (\ln x)^2 (-x^{-1}) + \int 2(\ln x)x^{-1}(-x^{-1}) \, dx = (\ln x)^2 (-x^{-1}) + \int 2(\ln x)x^{-2} \, dx$$

Let  $U = 2 \ln x$ ,  $V' = x^{-2}$ . Then  $U' = 2x^{-1}$ ,  $V = -x^{-1}$ . By integration-by-parts again, we have

$$\begin{aligned} \int (\ln x)^2 x^{-2} \, dx &= (\ln x)^2 (-x^{-1}) + 2(\ln x)(-x^{-1}) + \int 2x^{-1}x^{-1} \, dx \\ &= (\ln x)^2 (-x^{-1}) + 2(\ln x)(-x^{-1}) - 2x^{-1} + C. \end{aligned}$$

**[6 marks]**

(c) Let  $I(a, b) = \int_0^1 x^a (1-x)^b \, dx$ . Let  $u = x^a$ ,  $v' = (1-x)^b$ . Suppose  $a > 0$ . Then by integration-by-parts, we have

$$\begin{aligned} I(a, b) &= \left[ x^a \cdot \frac{-(1-x)^{b+1}}{b+1} \right]_0^1 - \int_0^1 a x^{a-1} \frac{-(1-x)^{b+1}}{b+1} \, dx \\ &= 0 + \frac{a}{b+1} I(a-1, b+1). \end{aligned}$$

Keep iterating this recursive formula until we reach  $I(0, b+a)$ :

$$\begin{aligned} I(a, b) &= \frac{a}{b+1} I(a-1, b+1) = \frac{a}{b+1} \frac{a-1}{b+2} I(a-2, b+2) = \frac{a}{b+1} \frac{a-1}{b+2} \frac{a-2}{b+3} I(a-3, b+3) \\ &= \dots = \frac{a}{b+1} \frac{a-1}{b+2} \dots \frac{1}{b+a} I(0, b+a) = \frac{a!b!}{(a+b)!} I(0, b+a) = \frac{a!b!}{(a+b)!} \int_0^1 (1-x)^{b+a} \, dx \\ &= \frac{a!b!}{(a+b)!} \left[ -\frac{(1-x)^{b+a+1}}{b+a+1} \right]_0^1 = \frac{a!b!}{(a+b)!} \left( 0 - \frac{-1}{b+a+1} \right) = \frac{a!b!}{(a+b+1)!} \end{aligned}$$

**[6 marks]**