

NANYANG TECHNOLOGICAL UNIVERSITY  
SPMS/DIVISION OF MATHEMATICAL SCIENCES

2023/24 Semester 1

MH5100 Advanced Investigations into Calculus I

Week 10

**Problem 1.** Suppose  $f$  and  $g$  are differentiable functions such that  $f(g(x)) = x$  and  $f'(x) = 1 + [f(x)]^2$ . Show that  $g'(x) = \frac{1}{1+x^2}$ .

**Solution 1.** From  $f(g(x)) = x$ ,

$$\begin{aligned} f'(g(x)) g'(x) &= 1 \\ (1 + [f(g(x))]^2) g'(x) &= 1 \\ (1 + x^2) g'(x) &= 1 \\ g'(x) &= \frac{1}{1+x^2} \end{aligned}$$

□

**Problem 2.** Find  $y''(0)$  if  $e^y + xy = e$ .

**Solution 2.** Differentiating implicitly w.r.t  $x$ ,

$$\begin{aligned} y'e^y + xy' + y &= 0 \\ y' &= -\frac{y}{e^y + x} \\ y''e^y + (y')^2 e^y + xy'' + y' + y' &= 0 \\ y''(e^y + x) + (y')^2 e^y + 2y' &= 0 \\ y'' &= -\frac{(y')^2 e^y + 2y'}{(e^y + x)} \end{aligned}$$

When  $x = 0, y = 1, y' = -e^{-1}$

$$\begin{aligned} y'' &= \frac{-(-e^{-1})^2 e + 2e^{-1}}{e} \\ &= \frac{-e^{-1} + 2e^{-1}}{e} \\ &= -\frac{1}{e^2} \end{aligned}$$

**Problem 3.** Let

$$f(x) = \frac{\sin^2 x}{1 + \cot x} + \frac{\cos^2 x}{1 + \tan x}.$$

Find  $f'(x)$ .

**Solution 3.** We first simplify  $f(x)$ ,

$$\begin{aligned} f(x) &= \frac{\sin^2 x}{1 + \cot x} + \frac{\cos^2 x}{1 + \tan x} = \frac{\sin^2 x}{1 + \frac{\cos x}{\sin x}} + \frac{\cos^2 x}{1 + \frac{\sin x}{\cos x}} \\ &= \frac{\sin^3 x}{\sin x + \cos x} + \frac{\cos^3 x}{\cos x + \sin x} = \sin^2 x - \sin x \cos x + \cos^2 x = 1 - \frac{1}{2} \sin 2x. \end{aligned}$$

Thus,

$$f'(x) = -\cos 2x.$$

**Problem 4.** Find the derivative function of  $F(x)$ .

$$F(x) = \begin{vmatrix} x-1 & 1 & 2 \\ -3 & x & 3 \\ -2 & -3 & x+1 \end{vmatrix}$$

**Solution 4.** Here we use Jacobi's formula.

$$\begin{aligned} F'(x) &= \begin{vmatrix} \frac{d}{dx}(x-1) & \frac{d}{dx}(1) & \frac{d}{dx}(2) \\ -3 & x & 3 \\ -2 & -3 & x+1 \end{vmatrix} + \begin{vmatrix} x-1 & 1 & 2 \\ \frac{d}{dx}(-3) & \frac{d}{dx}(x) & \frac{d}{dx}(3) \\ -2 & -3 & x+1 \end{vmatrix} + \begin{vmatrix} x-1 & 1 & 2 \\ -3 & x & 3 \\ \frac{d}{dx}(-2) & \frac{d}{dx}(-3) & \frac{d}{dx}(x+1) \end{vmatrix} \\ &= \begin{vmatrix} 1 & 0 & 0 \\ -3 & x & 3 \\ -2 & -3 & x+1 \end{vmatrix} + \begin{vmatrix} x-1 & 1 & 2 \\ 0 & 1 & 0 \\ -2 & -3 & x+1 \end{vmatrix} + \begin{vmatrix} x-1 & 1 & 2 \\ -3 & x & 3 \\ 0 & 0 & 1 \end{vmatrix} \\ &= \begin{vmatrix} x & 3 \\ -3 & x+1 \end{vmatrix} + \begin{vmatrix} x-1 & 2 \\ -2 & x+1 \end{vmatrix} + \begin{vmatrix} x-1 & 1 \\ -3 & x \end{vmatrix} \\ &= (x)(x+1) + 9 + (x^2 - 1) + 4 + x(x-1) + 3 \\ &= 3x^2 + 15 \end{aligned}$$

Jacobi's formula: Given that  $f_{ij}(x)$  ( $i, j = 1, 2, \dots, n$ ) is a differentiable function, we have

$$\frac{d}{dx} \begin{vmatrix} f_{11}(x) & f_{12}(x) & \cdots & f_{1n}(x) \\ f_{21}(x) & f_{22}(x) & \cdots & f_{2n}(x) \\ \vdots & \vdots & & \vdots \\ f_{n1}(x) & f_{n2}(x) & \cdots & f_{nn}(x) \end{vmatrix} = \sum_{k=1}^n \begin{vmatrix} f_{11}(x) & f_{12}(x) & \cdots & f_{1n}(x) \\ f_{21}(x) & f_{22}(x) & \cdots & f_{2n}(x) \\ \vdots & \vdots & & \vdots \\ f'_{k1}(x) & f'_{k2}(x) & \cdots & f'_{kn}(x) \\ \vdots & \vdots & & \vdots \\ f_{n1}(x) & f_{n2}(x) & \cdots & f_{nn}(x) \end{vmatrix}$$

The proof is straightforward if one knows how to evaluate the determinant of a matrix.

**Problem 5.** Show that the sum of the  $x$ - and  $y$ -interceptors of any tangent line to the curve  $\sqrt{x} + \sqrt{y} = \sqrt{c}$  is equal to  $c$ .

**Solution 5.** Differentiating implicitly w.r.t  $x$ ,

$$\begin{aligned} \frac{1}{2\sqrt{x}} + \frac{y'}{2\sqrt{y}} &= 0 \\ y' &= -\frac{\sqrt{y}}{\sqrt{x}} \end{aligned}$$

The tangent line is given by

$$\begin{aligned} y - y_1 &= -\frac{\sqrt{y_1}}{\sqrt{x_1}}(x - x_1) \\ y &= -\frac{\sqrt{y_1}}{\sqrt{x_1}}x + \sqrt{x_1 y_1} + y_1 \end{aligned}$$

When  $x = 0$ ,

$$\begin{aligned} y &= \sqrt{x_1 y_1} + y_1 \\ &= \sqrt{y_1}[\sqrt{x_1} + \sqrt{y_1}] \end{aligned}$$

When  $y = 0$ ,


$$\begin{aligned}
 0 &= -\frac{\sqrt{y_1}}{\sqrt{x_1}}x + \sqrt{x_1 y_1} + y_1 \\
 \frac{\sqrt{y_1}}{\sqrt{x_1}}x &= \sqrt{x_1 y_1} + y_1 \\
 x &= \frac{\sqrt{x_1}}{\sqrt{y_1}} [\sqrt{x_1 y_1} + y_1] \\
 &= \sqrt{x_1} [\sqrt{x_1} + \sqrt{y_1}]
 \end{aligned}$$

Summing up,

$$\begin{aligned}
 \sqrt{y_1} [\sqrt{x_1} + \sqrt{y_1}] + \sqrt{x_1} [\sqrt{x_1} + \sqrt{y_1}] &= [\sqrt{x_1} + \sqrt{y_1}] [\sqrt{x_1} + \sqrt{y_1}] \\
 &= \sqrt{c} \cdot \sqrt{c} \\
 &= c
 \end{aligned}$$

□

**Problem 6.** Let  $P(x, y)$  be a point on the parabola  $y^2 = 4px$  with focus  $F(p, 0)$ . Let  $\alpha$  be the angle between the parabola and the line segment  $FP$ , and let  $\beta$  be the angle between the horizontal line  $y = y_1$  and the parabola as in the figure. Prove that  $\alpha = \beta$ . (Thus, by a principle of geometrical optics, light from a source placed at  $F$  will be reflected along a line parallel to the  $x$ -axis.)



**Solution 6.** We start by finding the derivative of the parabola through implicit differentiation.

$$2y \frac{dy}{dx} = 4p \Rightarrow \frac{dy}{dx} = \frac{2p}{y}$$

We notice that  $\frac{2p}{y} = \tan \beta$ . We let  $\gamma$  be the angle between  $FP$  and the  $x$ -axis.  $\tan \gamma = \frac{y}{x-p}$ . We further notice that  $\alpha = \pi - (\pi - \gamma) - \beta = \gamma - \beta$

$$\begin{aligned}
 \tan \alpha = \tan (\gamma - \beta) &= \frac{\tan \gamma - \tan \beta}{1 + \tan \gamma \tan \beta} \\
 &= \frac{\left(\frac{y}{x-p}\right) - \left(\frac{2p}{y}\right)}{1 + \left(\frac{y}{x-p}\right) \left(\frac{2p}{y}\right)} \\
 &= \frac{y^2 - 2p(x-p)}{y(x+p)} \\
 &= \frac{2px + 2p^2}{y(x+p)} \\
 &= \frac{2p}{y}
 \end{aligned}$$

Therefore,  $\alpha = \beta$

□